

Serum Levels of Interleukin-36 Alpha and Interleukin-36 Receptor Antagonist In Behcet's Syndrome

Die Serumspiegel von Interleukin-36 Alpha und Interleukin-36 Receptor Antagonist im Behcet-Syndrom

Authors

Pelin Ünsal¹ , Pamir Çerçi² , Şükrü Alper Açıkgöz², Gökşal Keskin², Ümit Ölmez²

Affiliations

- 1 Department of Internal Medicine, Ankara University Faculty of Medicine, Ankara, Turkey
- 2 Department of Internal Medicine, Division of Immunology and Allergy, Ankara University Faculty of Medicine, Ankara, Turkey

Key words

Interleukin-36, Immunologie, Pathogenesis, Vasculitis, Behcet's syndrome

Schlüsselwörter

Interleukin-36, Studienlage, Pathogenese, Vaskulitis, Behcet-Syndrom

Bibliography

Akt Rheumatol 2022; 47: 233–238

DOI 10.1055/a-1550-2069

ISSN 0341-051X

© 2022. Thieme. All rights reserved.

Georg Thieme Verlag, Rüdigerstraße 14, 70469 Stuttgart, Germany

Correspondence

Pelin Ünsal

Department of Internal Medicine
Ankara University Faculty of Medicine
06100 Ankara
Turkey

Tel.: +90 (312) 3051538, Fax: +90 (312) 3051538

pelin_saracoglu@hotmail.com

ABSTRACT

Background Behcet's syndrome (BS) is a systemic vasculitic disorder. This study aimed to investigate the levels of serum IL-36 α and IL-36Ra in patients with BS.

Material and Methods A total of 80 subjects (60 BS patients and 20 healthy controls [HC]) were included.

Results The median IL-36 α level was 0.11 ng/ml in the BS group and 0.09 ng/ml in the HC group ($p = 0.058$). The mean IL-36Ra level was 13.62 pg/ml in the BS group and 13.26 pg/ml in the HC group ($p = 0.348$). Serum IL-36Ra levels of the active group were significantly higher ($p = 0.037$). Patients with oral ulcers and central nervous system involvement had higher serum IL36Ra levels. In the BS group, a positive correlation was found between serum IL-36Ra and CRP. In a multivariate analysis, the IL-36Ra level (OR = 1.067; 95 % CI = 1.001–1.137; $p = 0.045$) was independently associated with disease activity. **Conclusion** According to these findings, it is not clear whether such a slight difference is clinically significant, but they suggest that the IL-36 cytokine family may play a role in the course of the disease.

ZUSAMMENFASSUNG

Hintergrund Das Behcet-Syndrom (BS) ist eine Form der systemischen Vaskulitiden. Ziel dieser Studie war es, die Serumspiegel von IL-36 α und IL36Ra bei Patienten mit BS zu untersuchen.

Material und Methoden Insgesamt 80 Probanden (60 BS-Patienten und 20 gesunde Kontrollpersonen [KP]) wurden in die Studie eingeschlossen.

Ergebnisse Der mittlere IL-36 α -Spiegel betrug im BS 0,11 ng/ml und im KP 0,09 ng/ml ($p = 0,058$). Der mittlere IL-36Ra-Spiegel betrug 13,62 pg/ml im BS und 13,26 pg/ml im KP ($p = 0,348$). Der IL-36Ra Serumspiegel der aktiven Gruppe waren signifikant höher ($p = 0,037$). Patienten mit oralen Geschwüren und Beteiligung des zentralen Nervensystems hatten höhere IL-36Ra-Serumspiegel. In der BS Gruppe wurde eine positive Korrelation zwischen IL-36Ra und CRP im Serum gefunden. In der multivariaten Analyse war der IL-36Ra-Spiegel (OR = 1067; 95 % CI = 1001–1137; $p = 0045$) unabhängig mit der Krankheitsaktivität assoziiert.

Schlussfolgerung Nach diesen Erkenntnissen ist nicht klar, ob ein derart geringfügiger Unterschied eine klinische Bedeutung hat. Die Daten legen jedoch nahe, dass die IL-36-Zytofamilie möglicherweise eine Rolle im Krankheitsverlauf spielt.

Introduction

Behcet's syndrome (BS), that affect multiple systems in a chronic course with exacerbations and remissions, is a vasculitic disease. BS is characterized by recurrent oral, genital ulcers, uveitis, arthritis and vascular gastrointestinal, neurological, pulmonary and cardiac systems involvement [1].

The etiopathogenesis of the syndrome is still unclear. Inflammation, infection, genetic factors, and complex changes in innate and acquired immunity play a role in the pathogenesis [2]. Previous studies reported higher serum levels of inflammatory cytokines in patients with Behcet's syndrome [3, 4]. Mostly, Interleukin (IL)-1 family cytokines are investigated to better understand the disease pathogenesis. The new members of the IL-1 cytokine family are IL-36 α , IL-36 β , IL-36 γ , and IL-36 receptor antagonists (Ra). These cytokines are located on human chromosome 2 like most of the other IL-1 family members [5]. IL-36 plays a vital role in innate and acquired immunity. IL-36 stimulates dendritic cells from the host cells of innate and acquired immunity. Human dendritic cells also induce IL-1, IL-6, IL-12, IL-18 and IL-23 secretion when stimulated with IL-36 and increase CD83, CD86, and MHC II expression on the cell surface. In addition, IL-36 stimulation increases allogeneic T cell proliferation and IFN- γ release [6–8]. Previous studies showed that IL-36 may important in the pathogenesis of inflammatory diseases such as psoriasis, rheumatoid arthritis (RA), Sjögren's syndrome, systemic lupus erythematosus and inflammatory bowel diseases [9–11]. IL-36Ra acts as a natural antagonist of IL-36 α by binding to the IL-36 receptor. Deficiency of the interleukin-36 receptor antagonist may lead to an inflammatory skin condition known as generalized pustular psoriasis [12].

This study aimed to investigate the levels of serum IL-36 α and IL-36Ra in Turkish patients with BS and to determine its correlation with disease activity and clinical manifestations.

Materials and Methods

Participants

Eighty patients were included in the study and 60 of them were diagnosed with Behcet's Disease according to the "International Study Group for Behcet's Disease". Behcet's patients were followed up at Ankara University Faculty of Medicine, multidisciplinary BS diagnosis and treatment unit between 2015 and 2017. Twenty age and sex matched healthy persons who had no acute or chronic diseases were enrolled to serve as the healthy control (HC) group. Medical histories of all HCs were recorded and venous blood samples were taken after systemic and rheumatological assesment. The demographic data and symptoms of the patients were questioned, and the systemic involvement since the onset of the disease was evaluated. Active disease was defined as the presence of oral ulcers, and at least two of the clinical findings (skin lesions, genital ulcers, active arthritis and recent vascular, ocular or neurological involvement) found during the taking of blood samples. BS was also accepted as active if the high erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP) levels were detected with clinical symptoms despite current treatment [13]. The Ethical Committee of Ankara University Faculty of Medicine approved research proto-

col (No: 08–346–16). Each participant provided written informed consent.

Assay for IL-36 α and IL-36Ra

Serum IL-36 α and IL-36Ra levels were measured by sandwich enzyme-linked immunosorbent assay (ELISA) kits following the manufacturer's instructions (My BioSource, San Diego, CA, USA). The obtained venous blood samples for IL36- α and IL-36Ra measurement were stored at -80°C after the centrifugation process. The IL-36 α kit sensitivity was <0.094 ng/ml and the IL-36Ra kit sensitivity was 1.95 pg/ml.

Statistical analysis

Visual and analytical procedures were used to evaluate whether numerical variables were normally distributed. T-test or Mann Whitney U test was applied according to whether the variables were normally distributed. Chi-square test for categorical variables was used to compare possible associated factors. Spearman correlation constants were used when the relationship between variables was examined. Multivariate analyses was "performed using a backward logistic regression model to predict disease activity. Possible factors that were indicated in univariate analyses were entered into the logistic regression analyses in order to determine the independent correlates for disease activity. Hosmer-Lemeshow goodness of fit statistics were used to asses model fit. All data were analyzed using IBM SPSS Statistics 23. p values <0.05 were considered statistically significant.

Results

Sixty patients with BS and twenty age and sex-matched healthy controls were included in the study. Thirty-one of the patients had active disease (51.7%). The median age was 42 (18–65) in all BS patients and 41 (21–64) in the HCs. Median disease duration of BS patients was 7.5 (1–28) years. Demographic and clinical parameters are shown in ► **Table 1**.

Median serum IL-36 α level was 0.1102 (0.09–10.08) ng/ml in the BS group and 0.0940 (0.09–0.42) ng/ml in the HC group ($p=0.058$). Serum IL-36 α levels was similar both in patients with active and inactive disease (median,IQR 0.1117 [0.25] and 0.0998 [0.15], respectively; $p=0.8$). There was no statistically significant difference between serum IL-36 α levels in active versus control and inactive versus control groups ($p=0.069$ and $p=0.1$, respectively) (► **Fig. 1**).

Mean serum IL-36Ra level was found to be 15.3 ± 9.3 pg/ml in the BS group and 17.90 ± 13 pg/ml in the HC group ($p=0.348$). When compared with the inactive group, serum IL-36Ra levels of the active group were significantly higher (17.80 ± 10.63 pg/ml vs 12.77 ± 7.12 pg/ml. respectively, $p=0.037$). There was no statistically significant difference between serum IL-36 α levels of the active versus control and inactive versus control groups ($p=0.976$ and $p=0.082$) (► **Fig. 2**).

Serum IL-36Ra levels in patients with oral ulcers was significantly higher from without oral ulcer ($p=0.018$). Also, serum IL-36Ra levels were higher in patients with central nervous system (CNS) involvement than those without neurological involvement (24.45 ± 11.58 pg/ml vs 14.45 ± 8.71 pg/ml, $p=0.011$). No statisti-

► **Table 1** Demographic and clinical characteristics of Behcet's patients and healthy controls.

Parameter	All BD patients (n:60)	Active BD patients (n:31)	Inactive BD patients (n:29)	Healthy Controls (n:20)	p
Age (year) (median,IQR)	42 (16)	38 (20)	44 (11.5)	41 (21.5)	0.274
Sex (F)	31 (51.7%)	17 (54.8%)	14 (48.3%)	10 (50%)	0.872
Oral ulcers	59 (98.3%)	31 (100%)	28 (96.6%) *	-	
Genital ulcers	46 (76.7%)	23 (74.2%)	23 (79.3%) *	-	
Papulopustular eruption	35 (58.3%)	15 (48.4%)	20 (69%) *	-	
Uveitis	30 (50%)	11 (35.5%)	19 (65.5%) *	-	
Deep vein thrombosis	18 (30%)	6 (19.4%)	12 (41.4%) *	-	
Thrombophlebitis	12 (20%)	7 (22.6%)	5 (17.2%) *	-	
Arthritis	17 (28.3%)	11 (35.5%)	6 (20.7%) *	-	
Pathergy test (+)	17 (28.3%)	11 (35.5%)	6 (20.7%) *	-	
Central Nervous System involvement	5 (8.3%)	4 (12.9%)	1(3.4%) *	-	
Epididymitis	2 (3.3%)	1 (3.2%)	1 (3.2%) *	-	

* Not in the previous 3 months but during the course of the disease.

cally significant difference was found between serum IL-36Ra and other systemic manifestations. There was no statistically significant difference between serum IL-36 α levels and systemic involvement such as oral ulcers, genital ulcers, papulopustular eruption, uveitis, deep vein thrombosis, arthritis, Pathergy test positivity, CNS involvement, gastrointestinal involvement and epididymitis in Behcet's syndrome.

In BS group, positive correlation was found between serum IL-36Ra and CRP levels ($r=0.261$; $p=0.044$). There were no significant correlations between serum IL-36 α levels, sedimentation rates and CRP levels ($r=0.069$; $p=0.601$ and $r=0.039$; $p=0.769$, respectively). Also there was also no statistically significant difference between IL-36Ra levels and erythrocyte sedimentation rates ($r=0.206$; $p=0.115$).

In logistic regression analysis, gender, serum IL-36 α and IL-36Ra levels were included in the model to detect the independent correlates of disease activity. Serum IL-36Ra level (OR = 1.067; 95% CI = 1.001–1.137; $p=0.045$) was independently associated with disease activity.

Discussion

In our study we found that serum IL-36Ra levels were higher in patients with oral ulcers and central nervous system involvement. Also a positive correlation was found between serum IL-36Ra and CRP levels in BS group. Another important result was that serum IL-36Ra levels were independently associated with disease activity.

Behcet's syndrome is a systemic inflammatory disorder of unknown etiology. The IL-1 family cytokines play an important role in regulating the immune system by expression of proinflammatory mediators and integrins. The importance of IL-1 has been shown in Behcet's syndrome and other autoinflammatory and autoimmune disease such as RA, Sjögren's disease, gout, chondrocalcinosis, Still's disease, type 2 DM and interstitial lung disease [14]. In some other studies serum IL-33 and IL-37 levels, which are also

members of the IL-1 family, were reported higher in active Behcet patients than inactive BS and the control groups [13, 15].

IL-36, a new member of the IL-1 family, plays an essential role in the immune system through pro-inflammatory cytokine release and T cell proliferation [10]. Previous studies suggested that IL-36 α plays a role in psoriasis and inflammatory arthritis, IL-36 γ plays a role in airway inflammation, while IL-36 β plays a role in obesity-related inflammation [16]. Ciccia F et al. found that in Sjögren disease serum IL-36 α levels were higher and IL-36Ra levels were lower than in the control group [9]. Another study showed that IL-36 α levels are correlated with disease activity in systemic lupus erythematosus patients [17].

In our study, we found that serum IL-36 α levels were slightly higher in the active BS group compared to the HCs and the inactive BS group. Despite being statistically insignificant we still believe this finding should be noted due to its concordance with previous studies suggesting that inflammation in BS is mostly mediated by IL-1 cytokines [18]. But the more considerable and also statistically significant result of our study was the detection of higher IL-36Ra levels in active BS group. We presume this elevation may be a healthy response that aims to limit the inflammation. Due to lack of detailed studies about IL-36Ra levels in BS pathogenesis we can not support our presumption with existing literature. But fortunately, in recent years several comprehensive reviews regarding regulation, expression and immune functions of IL-36 family cytokines were published [11, 19, 20]. Although the certain mechanisms of these cytokines in the pathogenesis of inflammatory conditions are not revealed yet, the one thing all authors agreed on was the anti-inflammatory capability of IL-36Ra which uses a similar mechanism with IL-1Ra for antagonizing IL-1 α . Anakinra, drug the recombinant IL-1 receptor antagonist, is used in rheumatoid arthritis and deficiency of IL-1 Receptor Antagonist (DIRA) [21]. Since the recognition of the deficiency of the IL-36-receptor antagonist (DITRA), characterized by an aggressive form of psoriasis, numerous reports have been published suggesting that drugs targeting IL-1 pathway




Fig. 1 Serum IL-36 α levels in Behcet's syndrome and control group BS: Behcet's syndrome; *Median (Interquartile Range); p¹ denotes the comparison of patients with All BS patients vs. Healthy Controls; p² denotes the comparison of Active BS patients vs. Inactive BS patients.




Fig. 2 Serum IL-36 Ra levels in Behcet's syndrome and control group BS: Behcet's syndrome; p¹ denotes the comparison of patients with All BS patients vs. Healthy Controls; p² denotes the comparison of Active BS patients vs. Inactive BS patients.

may be beneficial in treatment [22–24]. However, IL-36Ra is currently not used as therapeutics on the clinical use.

IL-36 α , IL-36 β , IL-36 γ , and IL-36 Ra are generally produced by keratinocytes in the skin. IL-36 cytokines are important in infectious and inflammatory skin diseases, especially psoriasis [25]. In patients with psoriasis, IL-36Ra, IL-36 α , IL-36 β and IL-36 γ levels have been shown to be higher than the control group [26]. We didn't find any differences in serum IL-36 α and IL-36Ra levels in Behcet's patients with cutaneous involvement compared to without involvement. In our study, only serum IL-36Ra levels were higher in patients with oral ulcers than without.

IL-36 is an also important factor for arthritis. Frey S et al. found that synovium IL-36 α levels were higher in patients with psoriasis and RA than osteoarthritis. In this study, there was no difference

between the three groups of IL-36 receptor and IL-36 Ra expression [27]. In patients with psoriasis and rheumatoid arthritis, IL-36 α , γ and IL-36Ra levels were elevated and correlated with IL-1 β , but IL-36 β levels were elevated only in patients with RA [28]. In another study; IL-36 receptor is expressed in human synovial fibroblast and human articular chondrocytes but there was no increase in IL36 β levels in the inflamed human and mouse joints. Serum IL-36 β levels were also similar in healthy controls with rheumatoid arthritis, osteoarthritis [29]. In another study which compared RA and psoriatic arthritis showed that IL-36 α is expressed at similar levels in the synovium while psoriatic arthritis synovium expressed less IL-36Ra compared with RA. And researchers speculated that this result may be associated with decreased sensitivity to DMARDs in psoriatic arthritis patients. This study also demonstrated significantly higher plasma IL-36 α levels in RA patients compared with psoriatic arthritis patients, but no difference for the IL-36Ra levels [30]. In the TNF-induced arthritic mice, IL-36 α and IL-36 receptor levels were higher than normal mice and IL-36Ra levels were found to be similar. No changes in disease severity were observed after administration of anti-IL-36 receptor antagonist to TNF-induced arthritic mice [31]. As a result, it was thought that IL-36 was not a mediator of synovial inflammation [32]. In our study, we found that serum IL-36 α and IL-36Ra levels were similar in patients with and without arthritis.

In a brain cell study, IL-1Rrp2 (IL-36 receptor) was expressed in astrocytes and microglia, but this expression was not observed in neurons [33]. Previous studies found that IL-36 levels elevated in patients with multiple sclerosis and neuromyelitis optica spectrum disorder patients [21]. Karumbaiah L. et al. found that in an animal model in which brain micro motion is simulated, IL-36Ra levels were increased in astrocytes and microglia. In the same study, IL-36Ra levels were elevated after microwave electrodes were placed in adult mice [34]. In our study, IL-36Ra levels in Behcet's patients with neurological involvement were higher than those without neurological involvement. This result is similar to the published literature.

IL-36 α and IL-36 γ levels were increased in inflammatory mucosal tissues of patients with inflammatory bowel disease [35]. A recent study identified that higher levels of IL-36 α in fibrotic intestinal tissues from patients with Crohn's disease and ulcerative colitis compared with control individuals [36]. In our study, no statistically significant difference was found in IL-36 α and IL-36 Ra levels between patients with and without gastrointestinal involvement.

We believe the small sample size of our study was an important limitation and further studies are needed to elucidate the relationship between elevated IL-36Ra levels and activity of BS. Also we did not measure the level of IL-36Ra and IL-36 α in long term follow-ups of BS patients. This makes us unable to determine the possible changes of serum levels of these cytokines in a particular BS patient over time regarding to the activity of the disease. IL-36 expression is enhanced upon response to a number of stimuli including cytokines, lipopolysaccharides, infections and smoke. We assessed the disease activity with presence of clinical findings and acute phase reactants, thus we did not use a standart Behcet's syndrome activity scores because there is no gold standart test to evaluate disease activity and these scores had some limitations such as inconsistency between patients and physicians assesment, and

cultural differences [37]. Another limitation of our study is that this confounding factors was not evaluated [19]. Lastly, we performed our measurements on serum samples only. If obtained simultaneously, tissue samples from skin lesions or synovial fluids would have been valuable.

Conclusion

In conclusion, we found significantly higher serum levels of IL-36Ra in BS patients. Moreover this finding was associated with disease activity, serum CRP levels and clinical manifestations such as oral ulcers and neurological involvement. It is not clear whether such a slight difference has any clinical meaning, and should carefully consider the confounding factors which may affect the level of IL-36 α and IL-36Ra. This study supports that IL-36 cytokine family plays a role in the process of the disease, but further studies with larger groups are needed to confirm whether these cytokines can be used as disease-activity markers.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- [1] Alpsoy E. Behcet's disease: A comprehensive review with a focus on epidemiology, etiology and clinical features, and management of mucocutaneous lesions. *J Dermatol* 2016; 43: 620–632. doi:10.1111/1346-8138.13381
- [2] Zeidan MJ, Saadoun D, Garrido M et al. Behçet's disease physiopathology: a contemporary review. *Autoimmunity Highlights* 2016; 7: 4
- [3] Zhou Z, Chen S, Shen N et al. Cytokines and Behcet's disease. *Autoimmunity reviews* 2012; 11: 699–704
- [4] de Chambrun MP, Wechsler B, Geri G et al. New insights into the pathogenesis of Behcet's disease. *Autoimmunity reviews* 2012; 11: 687–698
- [5] Taylor SL, Renshaw BR, Garka KE et al. Genomic organization of the interleukin-1 locus. *Genomics* 2002; 79: 726–733. doi:10.1006/geno.2002.6752
- [6] Gabay C, Towne JE. Regulation and function of interleukin-36 cytokines in homeostasis and pathological conditions. *Journal of leukocyte biology* 2015; 97: 645–652
- [7] Foster AM, Baliwag J, Chen CS et al. IL-36 promotes myeloid cell infiltration, activation, and inflammatory activity in skin. *The Journal of Immunology* 2014; 192: 6053–6061
- [8] Mutamba S, Allison A, Mahida Y et al. Expression of IL-1Rrp2 by human myelomonocytic cells is unique to DCs and facilitates DC maturation by IL-1F8 and IL-1F9. *European journal of immunology* 2012; 42: 607–617
- [9] Ciccia F, Accardo-Palumbo A, Alessandro R et al. Interleukin-36 α axis is modulated in patients with primary Sjögren's syndrome. *Clinical & Experimental Immunology* 2015; 181: 230–238
- [10] Gresnigt MS, van de Veerdonk FL. Biology of IL-36 cytokines and their role in disease. In: *Seminars in immunology*. Elsevier; 2013: 458–465
- [11] Walsh PT, Fallon PG. The emergence of the IL-36 cytokine family as novel targets for inflammatory diseases. *Ann N Y Acad Sci* 2018; 1417: 23–34. doi:10.1111/nyas.13280
- [12] Marrakchi S, Guigue P, Renshaw BR et al. Interleukin-36-receptor antagonist deficiency and generalized pustular psoriasis. *N Engl J Med* 2011; 365: 620–628. doi:10.1056/NEJMoa1013068
- [13] Cerci P, Altiner S, Inal A et al. Investigating the role of IL-33 in the pathogenesis of Behcet's Disease. *Acta Clin Belg* 2017; 72: 434–438. doi:10.1080/17843286.2017.1314241
- [14] Lopalco G, Cantarini L, Vitale A et al. Interleukin-1 as a common denominator from autoinflammatory to autoimmune disorders: premises, perils, and perspectives. *Mediators of inflammation* 2015; 2015
- [15] Ozguclu S, Duman T, Ates FSO et al. Serum interleukin-37 level and interleukin-37 gene polymorphism in patients with Behcet disease. *Clinical rheumatology* 2019; 38: 495–502. doi:10.1007/s10067-018-4288-7
- [16] Hahn M, Frey S, Hueber AJ. The novel interleukin-1 cytokine family members in inflammatory diseases. *Current opinion in rheumatology* 2017; 29: 208–213
- [17] Chu M, Wong CK, Cai Z et al. Elevated expression and pro-inflammatory activity of IL-36 in patients with systemic lupus erythematosus. *Molecules* 2015; 20: 19588–19604
- [18] Cho S, Kim J, Cho SB et al. Immunopathogenic characterization of cutaneous inflammation in Behcet's disease. *Journal of the European Academy of Dermatology and Venereology : JEADV* 2014; 28: 51–57. doi:10.1111/jdv.12054
- [19] Bassoy EY, Towne JE, Gabay C. Regulation and function of interleukin-36 cytokines. *Immunol Rev* 2018; 281: 169–178. doi:10.1111/imr.12610
- [20] Boutet MA, Nerviani A, Pitzalis C. IL-36, IL-37, and IL-38 Cytokines in Skin and Joint Inflammation: A Comprehensive Review of Their Therapeutic Potential. *Int J Mol Sci* 2019; 20. doi:10.3390/ijms20061257
- [21] Zhou L, Todorovic V. Interleukin-36: Structure, Signaling and Function. *Adv Exp Med Biol* 2020. doi:10.1007/5584_2020_488
- [22] Bonekamp N, Caorsi R, Viglizzo GM et al. High-dose ustekinumab for severe childhood deficiency of interleukin-36 receptor antagonist (DITRA). *Ann Rheum Dis* 2018; 77: 1241–1243. doi:10.1136/annrheumdis-2017-211805
- [23] Gomez-Garcia F, Sanz-Cabanillas JL, Viguera-Guerra I et al. Scoping Review on Use of Drugs Targeting Interleukin 1 Pathway in DIRA and DITRA. *Dermatol Ther (Heidelb)* 2018; 8: 539–556. doi:10.1007/s13555-018-0269-7
- [24] Hospach T, Glowatzki F, Blankenburg F et al. Scoping review of biological treatment of deficiency of interleukin-36 receptor antagonist (DITRA) in children and adolescents. *Pediatr Rheumatol Online J* 2019; 17: 37. doi:10.1186/s12969-019-0338-1
- [25] Buhl AL, Wenzel J. Interleukin-36 in Infectious and Inflammatory Skin Diseases. *Front Immunol* 2019; 10: 1162. doi:10.3389/fimmu.2019.01162
- [26] Johnston A, Xing X, Guzman AM et al. IL-1F5,-F6,-F8, and-F9: a novel IL-1 family signaling system that is active in psoriasis and promotes keratinocyte antimicrobial peptide expression. *The Journal of Immunology* 2011; 186: 2613–2622
- [27] Frey S, Derer A, Messbacher M-E et al. The novel cytokine interleukin-36 α is expressed in psoriatic and rheumatoid arthritis synovium. *Annals of the rheumatic diseases* 2013; 72: 1569–1574
- [28] Boutet MA, Bart G, Penhoat M et al. Distinct expression of interleukin (IL)-36 α , β and γ , their antagonist IL-36Ra and IL-38 in psoriasis, rheumatoid arthritis and Crohn's disease. *Clin Exp Immunol* 2016; 184: 159–173. doi:10.1111/cei.12761
- [29] Magne D, Palmer G, Barton JL et al. The new IL-1 family member IL-1F8 stimulates production of inflammatory mediators by synovial fibroblasts and articular chondrocytes. *Arthritis research & therapy* 2006; 8: R80

- [30] Boutet MA, Nerviani A, Lliso-Ribera G et al. Interleukin-36 family dysregulation drives joint inflammation and therapy response in psoriatic arthritis. *Rheumatology (Oxford)* 2020; 59: 828–838. doi:10.1093/rheumatology/kez358
- [31] Derer A, Groetsch B, Harre U et al. Blockade of IL-36 receptor signaling does not prevent from TNF-induced arthritis. *PloS one* 2014; 9
- [32] Dietrich D, Gabay C. Inflammation: IL-36 has proinflammatory effects in skin but not in joints. *Nature Reviews Rheumatology* 2014; 10: 639
- [33] Berglöff E, Andre R, Renshaw BR et al. IL-1Rrp2 expression and IL-1F9 (IL-1H1) actions in brain cells. *Journal of neuroimmunology* 2003; 139: 36–43
- [34] Karumbaiah L, Norman SE, Rajan NB et al. The upregulation of specific interleukin (IL) receptor antagonists and paradoxical enhancement of neuronal apoptosis due to electrode induced strain and brain micromotion. *Biomaterials* 2012; 33: 5983–5996. doi:10.1016/j.biomaterials.2012.05.021
- [35] Nishida A, Hidaka K, Kanda T et al. Increased Expression of Interleukin-36, a Member of the Interleukin-1 Cytokine Family, in Inflammatory Bowel Disease. *Inflammatory bowel diseases* 2016; 22: 303–314. doi:10.1097/mib.0000000000000654
- [36] Scheibe K, Kersten C, Schmied A et al. Inhibiting Interleukin 36 Receptor Signaling Reduces Fibrosis in Mice With Chronic Intestinal Inflammation. *Gastroenterology* 2019; 156: 1082–1097 e1011. doi:10.1053/j.gastro.2018.11.029
- [37] Floris A, Espinosa G, Serpa Pinto L et al. Discordance between patient and physician global assessment of disease activity in Behcet's syndrome: a multicenter study cohort. *Arthritis Res Ther* 2020; 22: 278. doi:10.1186/s13075-020-02362-1