

Screening of Antimicrobial Activity of Essential Oils against Bovine Respiratory Pathogens – Focusing on *Pasteurella multocida*



Authors

Doris Bismarck¹, Jens Becker², Elisabeth Müller¹, Vera Becher³, Lisa Nau³, Philipp Mayer³

Affiliations

- 1 Laboklin GmbH & Co.KG, Bad Kissingen, Germany
- 2 Clinic for Ruminants, Vetsuisse-Faculty, University of Bern, Bern, Switzerland
- 3 SaluVet GmbH, Bad Waldsee, Germany

Key words

essential oils, *Pasteurella multocida*, *Mannheimia haemolytica*, bovine respiratory disease

received

February 28, 2021

accepted after revision

December 13, 2021

published online

February 18, 2022

Bibliography

Planta Med 2022; 88: 274–281

DOI 10.1055/a-1726-9291

ISSN 0032-0943

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

Correspondence

Dr. Doris Bismarck

Laboklin GmbH & CO.KG

Steubenstraße 4, 97688 Bad Kissingen, Germany

Phone: + 49 97 17 20 20, Fax: + 49 97 16 85 46

bismarck@laboklin.com



Supplementary material is available under
<https://doi.org/10.1055/a-1726-9291>

ABSTRACT

Administration of essential oils as natural plant products with antimicrobial activity might be an alternative to antibiotic treatment of bovine respiratory disease. The aim of this study was to analyse the *in vitro* antimicrobial activity of 11 essential oils against *Pasteurella multocida* isolated from the respiratory tract of calves using microdilution with determination of minimum inhibitory and bactericidal concentration as well as agar disc diffusion. Additionally, antimicrobial activity against *Mannheimia haemolytica* and bacteria in the *Mannheimia* clade was assessed by agar disc diffusion. Seven essential oil mixtures were also tested against all bacterial isolates. *P. multocida* was strongly inhibited by cinnamon cassia and lemongrass oil followed by coriander, winter savory, thyme, clove, and peppermint oil in the microdilution assays. Eucalyptus, wintergreen, spruce, and star anise oil showed lower activity. Comparison of both methods revealed an underestimation of cinnamon cassia oil activity by agar disc diffusion and conflicting results for wintergreen oil in microdilution, which precipitated in broth. Cinnamon cassia, thyme, wintergreen, lemongrass, and winter savory oil all showed strong antimicrobial activity against *M. haemolytica*. Bacteria in the *Mannheimia* clade were mostly inhibited by cinnamon cassia and thyme oil. *Pasteurella* isolates were more susceptible to inhibition by essential oils than *Mannheimia* isolates. Essential oil mixtures did not show stronger antibacterial activity than single essential oils. In conclusion, cinnamon cassia and lemongrass as well as coriander, winter savory, and thyme oil are promising candidates for treatment of *P. multocida*-associated bovine respiratory infections.

Introduction

Pasteurella multocida and *Mannheimia haemolytica* are commensals and also opportunistic pathogens in multifactorial BRD [1,2]. In cattle, BRD is a leading cause of morbidity and mortality [3]. Pathogenesis of BRD depends on multiple factors, such as the animal host, the environment, and involved pathogens [1,2]. Of these, *P. multocida* is an important causative agent in the development of BRD [4]. *M. haemolytica* is a major cause of severe fibrin-

ous pleuropneumonia [5]. Clinical signs of infection involve respiratory distress, cough, fever, nasal discharge, and inappetence associated with weight loss [1,4,6].

Often, antimicrobials are used for metaphylactic or therapeutic treatment of clinical BRD. A European study observed a still high susceptibility of *P. multocida* and *M. haemolytica* isolated from diseased, antimicrobial non-treated cattle against licensed antibiotics [7]. However, in North America, the prevalence of non-susceptible respiratory tract isolates in diseased animals is steadily increasing

ABBREVIATIONS

ANOVA	analysis of variance
BRD	bovine respiratory disease
CAMHB	cation-adjusted Mueller-Hinton broth
EO	essential oil
IZR	inhibition zone radius
MALDI-TOF MS	matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry
MBC	minimum bactericidal concentration
MIC	minimum inhibitory concentration

[8]. Several studies analysed the association of antimicrobial treatment and development of bacterial resistances. For example, multiple antibacterial treatments are associated with a higher probability of antimicrobial resistances in cattle [9]. Lower respiratory tract isolates of cattle receiving metaphylactic tulathromycin and chlortetracycline treatment showed a high macrolide resistance level [10]. Stanford et al. [11] found an increased antimicrobial resistance for macrolides, higher MICs across antimicrobial classes, and more resistance-related gene determinants in BRD isolates from cattle managed with antimicrobials in comparison to non-treated animals. This emphasises the importance of prudent antimicrobial use and the necessity of ongoing research on alternatives for prophylaxis, metaphylaxis, and treatment of BRD.

Medicinal plants containing EOs bear great potential for combined or alternative therapy of respiratory conditions in cattle [12]. EOs like thyme, eucalyptus, tea tree, and ravintsara are already used by some practicing veterinarians for the treatment of respiratory infections via oral, topic, or rectal application [13]. The inhalation of EOs is also recommended [14]. EOs are natural plant products and complex mixtures of volatile lipophilic compounds like monoterpenes, sesquiterpenes, and/or phenylpropanoids [15]. They are mainly produced by water or steam distillation or mechanical processing of plant material. As an *in vitro* screening method for antibacterial activity of EOs, agar disc diffusion can be used [16], which provides an affordable method for susceptibility testing of EOs in veterinary routine diagnostics. Agar disc diffusion has disadvantages due to volatility and different solubility of EO components in the agar. The more elaborate broth dilution is the preferred method for analyses of EO activity, but hardly feasible in the daily laboratory routine.


Reports describing *in vitro* antibacterial activity of EOs against bacterial bovine respiratory pathogens are scarce [17–19]. Hence, further knowledge about the potential of EOs as an alternative treatment of BRD is needed. The aim of this study was (i) to analyse the *in vitro* antimicrobial activity of 11 EOs against *P. multocida* isolates from the respiratory tract of calves by microdilution, determining both the minimum inhibitory and bactericidal concentrations as well as by agar disc diffusion for comparison purposes. Additionally, (ii) to analyse antimicrobial activity of EOs against *M. haemolytica* and bacteria in the *Mannheimia* clade and (iii) antibacterial activity of seven EO mixtures against all bacterial isolates by agar disc diffusion. Thus, EOs for further research on phytotherapeutic prophylaxis and treatment of BRD should be identified.

Results

In microdilution, cinnamon cassia and lemongrass oil showed the strongest antibacterial activity against *P. multocida* with median MICs of 0.0078 and 0.0313%, respectively (► Fig. 1 a). Coriander, winter savory, and thyme oil had an MIC of 0.125%, followed by clove (MIC of 0.1875%), peppermint (MIC of 0.25%), and eucalyptus and wintergreen oil (both: MIC of 0.5%). The weakest EOs were spruce (MIC of 1.0%) and star anise oil (MIC > 1.0%). It is important to mention that wintergreen oil immediately precipitated in the broth at the bottom of the well. Therefore, the visual determination of the MIC for wintergreen oil was not distinct. The MBC differed from the MIC, except for winter savory, thyme, and eucalyptus oil (► Fig. 1 a). For several bacterial isolates, the MBC and MIC differed by one dilution step. This applies for spruce (10% of isolates), coriander, clove, peppermint oil (20–26% of isolates), lemongrass (43% of isolates), and cinnamon cassia oil (67% of isolates). For wintergreen and star anise oil, the MIC and MBC were not always possible to determine. MICs that did not have bactericidal effects led to a reduction of bacterial growth on plated agar plates, in contrast to growth controls. In agar disc diffusion, lemongrass oil (median IZR: 30.13 mm, interquartile range: 7.31) most effectively inhibited the growth of *P. multocida*, followed by cinnamon cassia (22.25 mm, 2.38), thyme (18.5 mm, 2.38), winter savory (16.75 mm, 0.94), coriander (15.00 mm, 2.88), wintergreen (14.5 mm, 1.25), peppermint (13.5 mm, 2.38), and clove oil (11.25 mm, 1.88) (► Fig. 1 b). Comparison of the results of agar disc diffusion and microdilution of *P. multocida* (► Fig. 2) revealed two major differences in EO activity. Cinnamon cassia oil was much less effective than expected from the microdilution results. Wintergreen oil showed better activity in agar disc diffusion than expected from the microdilution results, where precipitation of the EO occurred. The other EOs showed concordant results of both methods. Thyme, winter savory, and coriander oil had similar activity in agar disc diffusion (IZR of 18.5–15.00 mm) and the same MIC (0.125%) in microdilution. Peppermint and clove oil also showed similar activity in agar disc diffusion (IZR of 13.5–11.25 mm) and microdilution (MIC values of 0.25 and 0.1875%). Star anise, spruce, and eucalyptus oil had the weakest activity in both agar disc diffusion (IZR of 7.50–5.00 mm) and microdilution (MIC of 0.5% – ≥ 1.0%).

Isolates of *M. haemolytica* and bacteria in the *Mannheimia* clade were most efficiently inhibited by cinnamon cassia and thyme oil in agar disc diffusion (► Fig. 3). *M. haemolytica* was also inhibited by wintergreen, lemongrass, winter savory, clove, coriander, peppermint, and star anise oil in descending order. Bacteria in the *Mannheimia* clade were also inhibited by winter savory, coriander, wintergreen, lemongrass, peppermint, clove, and star anise oil. Eucalyptus and spruce oil only yielded small or no inhibition zones against all bacterial isolates. A two-way ANOVA was conducted to examine the effect of bacterial isolates and EOs on the size of the IZR. There was a statistically significant interaction between the EO and bacterial isolate ($F = 12.37$, $df = 20, 279$, $p < 0.001$). Predominantly, *P. multocida* isolates were more susceptible than *Mannheimia* isolates, except for star anise and eucalyptus oil (► Fig. 3).

Additionally, agar disc diffusion of EO mixtures, composed of the most potent single EOs, was performed. All EO mixtures showed antibacterial activity against all bacterial isolates (Fig. 1S,




► **Fig. 1** Depicted is (a) the minimum inhibitory and bactericidal concentration (MIC and MBC) in % of EOs and (b) the inhibition zone radius (IZR) in mm against *P. multocida* ($n = 10$). Statistically significant results of comparison of MICs or IZR of EOs are given. Conducted was a Kruskal-Wallis test (chi square = 100.94, $df = 10$, $p < 0.001$) and post hoc Dunn test with Benjamini-Hochberg adjustment for pairwise comparison. P values of statistically significant differences and indicating asterisks (* p value ≤ 0.05 , ** p value ≤ 0.01 , *** p value ≤ 0.001 , -not significant) are stated within the figure. (1) MIC and MBC $> 1\%$

Supporting Information). A two-way ANOVA was conducted that examined the effect of bacterial isolates and EO mixtures on the size of the IZR. There was no interaction between the EO and bacterial isolate ($F = 1.14$, $df = 12$, 189 , $p = 0.328$), but the bacterial isolate ($F = 266.44$, $df = 2$, 189 , $p < 0.001$) as well as the EO mixture ($F = 8.01$, $df = 6$, 189 , $p < 0.001$) had an effect on the size of the IZR. Multiple comparison of the main effects revealed that *P. multocida* was more susceptible against the EO mixtures than *M. haemolytica* ($p < 0.001$) and bacteria in the *Mannheimia* clade ($p < 0.001$). *M. haemolytica* was also more susceptible than bacteria in the *Mannheimia* clade ($p < 0.001$) (Fig. 1S, Supporting Information). The EO mixtures did not exceed antibacterial activity of the single EOs (Fig. 2S, Supporting information).

Discussion

The analysis of antimicrobial activity of EOs is lacking standardised methods and is demanding due to their complexity, water insolubility, and volatility [20]. In agar disc diffusion, the size of the inhibition

zone of bacterial growth depends on the diffusion of the EO components into the agar, which could be impaired by their hydrophobicity and volatility [16]. However, inhibition of bacterial growth proves antibacterial activity. Broth dilution enables direct contact between all EO components and the pathogen and allows for the determination of the MIC and subsequent determination of the MBC. Bacteriostatic activity has been defined as a ratio of MBC to MIC of > 4 [21], lower ratios indicate a bactericidal activity. For the EOs analysed in this study, a bactericidal mode of action can be assumed, because their MBC-MIC ratio is 1–2. Broth dilution is considered the more reliable method for analysis of antimicrobial activity of EOs. Nevertheless, agar disc diffusion is used in veterinary routine diagnostics to predict EO activity before therapeutic use (aromatogram). For the specific EOs analysed in this study, agar disc diffusion is generally valid, as shown by the comparison of the results of both methods. Two exceptions are the underestimation of cinnamon cassia oil activity and the conflicting results for wintergreen oil. Wintergreen oil immediately precipitates in the broth despite the presence of an emulsifier. In this case, the microdilution has to be interpreted with cau-



► **Fig. 2** Plotted is the minimum inhibitory concentration (MIC) and the inhibition zone radius (IZR) of EOs against *P. multocida* and the regression line with 0.95 confidence interval of prediction. Cinnamon cassia oil, which shows largest divergence, is encircled.

tion. The limitations of agar disc diffusion account for *Pasteurella* and *Mannheimia* isolates in the same manner and comparison of differences in their susceptibility are admissible. Predominantly, *P. multocida* isolates are more susceptible than *Mannheimia* isolates, which was also observed by others [17, 19]. Moreover, agar disc diffusion of seven EO mixtures was performed. The good antimicrobial activity of all EO mixtures against *P. multocida* most likely depends on the high content of lemongrass oil (60–70%), which showed strong antimicrobial activity against *P. multocida* in agar disc diffusion. For determination of the synergistic effects of EO mixtures, determination of the fractional inhibitory concentration using checkerboard assays could be performed in future studies.


Only a few studies investigated the antimicrobial activity of EOs against BRD pathogens. Two studies analysed the effect of winter savory and thyme oil against *P. multocida* [17, 19]. Amat et al. [17] observed good activity of thyme oil (*Thymus zygis* L.) against one bovine clinical respiratory tract isolate of *P. multocida*, with an MIC of 0.013%. They [17] also found the MIC equalled the MBC for this *Pasteurella* isolate, similar to the results obtained in our study. In a study from Salzmann [19], winter savory and thyme oil (*Thymus vulgaris* L.) showed strong antibacterial activity against bovine *P. multocida* and *M. haemolytica* isolates using agar disc diffusion and microdilution, with MICs of 0.39 and 0.63%, respectively. In the same study [19], eucalyptus oil had a weak antibacterial effect against *P. multocida* in agar disc diffusion and inhibited bacterial growth in microdilution, with an MIC of 0.23%. Different MICs of thyme oil obtained in these studies could be explained by different plant species of the thyme used (*T. zygis* L. and *T. vulgaris* L.), but all belonging to *Thymi herba*. EOs as natural plant products can vary in their composition of single chemical compounds due to the plant parts used, extraction methods, region of growth, or harvest season [22]. The antimicrobial activity of an EO can differ depending on its composition [23]. Thus, analysed antimicrobial activities apply for the specific batch of the tested EO within a

study. A study by Kissels et al. [24] analysed the antimicrobial activity of the EO components thymol, carvacrol, and 1,8-cineole against bovine type strains of *P. multocida* and *M. haemolytica* using microdilution: thymol and carvacrol, which are major components of thyme and winter savory oil, exhibited strong antimicrobial activity. The monoterpenoid 1,8-cineole, which is a major component of eucalyptus oil, had no antibacterial effect [24]. This strong and weak activity of major EO components [24] equates the antimicrobial activity of their EOs found in our study.

Toxic side effects are not expected, as low MICs obtained for most EOs implicate a possible low dose application of EOs. EO preparations could be applied orally or by inhalation, direct nasal, or rectal application. *In vitro* and *in vivo* studies demonstrating EO safety and clinical efficiency on respiratory symptoms in animals and humans encourage their future use in veterinary medicine [17, 25–28]. Fararh et al. [26] showed that an orally applied EO mixture of eucalyptus, lemon, cornment, rosemary, and *Echinacea purpurea* (0.015–0.03%) in combination with an injection of tulathromycin reduced respiratory manifestation and improved blood gas values in calves with respiratory disease caused by *P. multocida* and *Haemophilus* spp. Inhalation or direct nasal application of diluted EO preparations would enable specific targeting of the respiratory tract. In the literature, it has been reported [17] that *in vitro*, no cytotoxic effect against bovine turbinate cells could be detected by the use of diluted thyme or eucalyptus oil in concentrations of $\leq 0.4\%$. The MIC values for thyme and eucalyptus oil found in our study are below this value, thus safe application of those EOs should be possible while still exerting antibacterial activity. Further, only minimal or no cytotoxic effect could be observed *in vitro* by LeBel et al. [25] for diluted thyme and winter savory oil ($\leq 0.078\%$) against porcine tracheal epithelial cells. Human studies showed a short-term beneficial effect of eucalyptus oil inhalation (final air concentration of 0.056%) [27] or nasal application of an EO spray mixture (*Eucalyptus citriodora* and *globulus*, *Mentha piperita*, *Origanum syriacum*, *Rosmarinus officinalis*) [28].

Besides antimicrobial activity, other positive effects of EOs might contribute to the host defence against respiratory tract infections. Wu et al. [29] showed that linalool, a major compound of coriander and thyme oil (chemotype linalool), activated the Nrf-2 signalling pathway in a human lung cell line and impaired the expression of pro-inflammatory cytokines of *P. multocida*. Subcutaneous treatment with linalool also led to enhanced clearance of *P. multocida*, decreased lung neutrophil accumulation, and improved survival of mice [29]. Additionally, Amat et al. [30] observed only minimal *in vitro* effects of thyme and eucalyptus oil on commensal microbiota like Lactobacilli, Bacilli, and Staphylococci, which might be able to maintain respiratory health by inhibition of pathogen colonisation.

Overall, cinnamon cassia and lemongrass oil showed the best antimicrobial activity against bovine respiratory tract isolates of *P. multocida*, followed by coriander, winter savory, and thyme oil. Comparison of microdilution and agar disc diffusion of *P. multocida* isolates showed reliable results obtained in agar disc diffusion for the specific EOs analysed in this study, except for an underestimation of cinnamon cassia oil activity and inconsistent results of wintergreen oil. *Mannheimia* isolates were strongly inhibited by cinnamon cassia and thyme oil in agar disc diffusion. Further re-



► **Fig. 3** Depicted is the inhibition zone radius (IZR) in mm of EOs against *P. multocida* (n = 10), *M. haemolytica* (n = 10), and *Mannheimia* clade (n = 10). Only statistically significant results of comparison of differences in susceptibility of *P. multocida* vs. *M. haemolytica* vs. *Mannheimia* clade against the same EO are given. Conducted was a two-way ANOVA (interaction between EO and bacterial isolate: $F = 12.37$, $df = 20, 279$, $p < 0.001$) and a post hoc pairwise comparison with Benjamini-Hochberg adjustment. P values of statistically significant differences and indicating asterisks (*p value ≤ 0.05 , **p value ≤ 0.01 , ***p value ≤ 0.001) are stated with the significance bars.

search is needed to validate the clinical potential, especially of cinnamon cassia, lemongrass, and thyme oil in BRD treatment.

Material and Methods

Bacterial isolates

Clinical bacterial isolates of *P. multocida* (n = 10), *M. haemolytica* (n = 10), and *Mannheimia* clade (n = 10) were analysed in this study. All *P. multocida*, three *M. haemolytica*, and all *Mannheimia* clade isolates were derived from deep nasopharyngeal swabs of veal calves, sampled within the “outdoor veal calf” project [31]. Nasopharyngeal swabs of the calves were streaked onto *Pasteurella* selective agar purchased from Thermo Scientific Oxoid. Plates were incubated at 37 °C for 24 h. One single colony per plate was selected for species identification with MALDI-TOF MS using Microflex LT from Bruker Daltonik GmbH. Colonies identified as *P. multocida*, *M. haemolytica*, and bacteria in the *Mannheimia* clade (identification on genus level) were purified on trypticase soy agar plates containing 5% sheep blood purchased from Becton Dickinson and incubated at 37 °C for 24 h. Identification was confirmed by MALDI-TOF MS and the isolates were frozen in 30% glycerol stocks at –80 °C. Seven *M. haemolytica* isolates derived from bovine nose swabs which were sent in for veterinary routine diagnostics to Laboklin GmbH & Co.KG. Since these isolates originated from routine diagnostic samples, no further information on the sampled animals was available and isolates were chosen randomly from the diagnostic pool. Isolates were identified by colony morphology and MALDI-TOF MS with Microflex LT/SH by Bruker Daltonik GmbH. A confidence score was used to rate agreement between the created spectrum and reference spectra provided by the Bruker database. Isolates had confidence scores > 2

allowing for genus and species identification. Bacterial isolates were stored frozen with the CRYOBANK system from Mast Diagnostica at –20 °C without additional additives.

Essential oils and essential oil mixtures

The tested EOs (► Table 1) and EO mixtures (► Table 2) were provided by SaluVet GmbH. Spruce oil was obtained from two different plant types. EO mixtures were composed of EOs that showed the strongest antibacterial activity tested individually. Lemongrass oil, according to results of previously carried out internal examinations, was expected to have a very strong antibacterial effect against all tested microorganisms. Therefore, it was chosen as the main component of all EO mixtures, with a content of at least 60%. Other oils were added in such a concentration that possible synergistic effects were expected to be detectable, whilst lemongrass oil still mainly determined the effect.

Agar disc diffusion

All bacterial isolates [*P. multocida* (n = 10), *M. haemolytica* (n = 10), and *Mannheimia* clade (n = 10)] were analysed by agar disc diffusion. Cryopreserved bacteria were recovered at 36 °C on Columbia agar with 5% sheep blood purchased from Becton Dickinson 24 h before performing agar disc diffusion. Agar disc diffusion of EOs or EO mixtures was performed on Mueller-Hinton agar with 5% sheep blood from Becton Dickinson as previously described by us [32]. Agar plates were inoculated with 100 µL of bacterial suspension adjusted to 0.5 McFarland in sterile 0.9% NaCl solution. Afterwards, a sterile 6-mm diameter blank filter paper disc purchased from Becton Dickinson was placed in the middle of the inoculated agar plate and loaded with 10 µL of the EO or EO mixture to be analysed. After overnight incubation at 36 °C, the inhibition zones

► **Table 1** The 11 analysed EOs with their common name, scientific name, plant species, used plant part, origin of plants, extraction method, and chemical composition according to manufacturer or literature.

EO common name	Scientific name	Plant species	Plant part	Origin of used plants	Extraction method	Chemical composition given by the [A] manufacturer/[B] literature
Cinnamon cassia	<i>Cinnamomi cassiae aetheroleum</i>	<i>Cinnamomum cassia</i> (L.) J.Presl	leaves and young branches	China	steam distillation	[A] cinnamaldehyde 77.9%, <i>trans</i> -2-methoxycinnamaldehyde 9.1%, cinnamyl acetate 3.4%, coumarin 2.1%, eugenol 0.1%
Clove	<i>Caryophylli aetheroleum</i>	<i>Syzygium aromaticum</i> (L.) Merr. et L.M. Perry	leaves	Indonesia	steam distillation	[B] eugenol (60–95%) [39]; (75–88%) [40], α - and β -caryophyllene (5–10%) [39]; (75–88%) [40], acetyleugenol (2–27%) [39], (4–15%) [40], < 1%: 2-heptanone, ethyl hexanoate, humuleneol; sesquiterpenes, α -humulene, α -humulene epoxide, β -humulene, β -caryophyllene oxide, α -cubebene, α -copaene, α -cadinene, δ -cadinene, α -ylangene, calacorene, calamenene [40]
Coriander	<i>Coriandri aetheroleum</i>	<i>Coriandrum sativum</i> L.	fruits	Russia	steam distillation	[A] linalool 71.04%, α -pinene 6.12%, γ -terpinene 4.91%, camphor 4.53%, geranyl acetate 3.74%, limonene 2.59%, geraniol 1.78%, <i>p</i> -cymene 0.65%, α -terpineol 0.28%
Eucalyptus	<i>Eucalypti aetheroleum</i>	<i>Eucalyptus globulus</i> Labill.	leaves and branches	China	steam distillation	[B] 1,8-cineole (54–95%), α -pinene (2.6%), <i>p</i> -cymene (2.7%) [39]
Lemongrass	<i>Cymbopogoni aetheroleum</i>	<i>Cymbopogon flexuosus</i> Nees ex Wats.	aerial parts	India	steam distillation	[B] citral (75%), geraniol (5.6%) [41]
Peppermint	<i>Menthae piperitae aetheroleum</i>	<i>Mentha x piperita</i> L.	aerial parts	India	steam distillation	[A] menthol 40.32%, menthone 25.22%, menthyl acetate 5.33%, cineole 4.6%, isomenthone 4.16%, menthofuran 2.79%, limonene 2.14%, pulegone 1.00%, isopulegol 0.19%, carvone 0.07%
Spruce	<i>Piceae aetheroleum</i>	<i>Abies sibirica</i> Ledeb.; <i>Abies alba</i> Mill.	needles and branches	Russia	steam distillation	[B] <i>Abies sibirica</i> : bornylacetate (30–40%), camphene (ca. 10%), santene, α - and β -pinene, α -phellandrene, limonene; <i>Abies alba</i> : bornylacetate (37–49%), camphene (10–17%), α - and β -pinene [38]
Star anise	<i>Anisi stellati aetheroleum</i>	<i>Illicium verum</i> Hook. fil.	fruits	China	steam distillation	[B] <i>trans</i> -anethole (80–90%), methylchavicol, foeniculin, <i>cis</i> -anethol, anisaldehyde, limonene, linalool, α -pinene [38]
Thyme	<i>Thymus zygis aetheroleum</i>	<i>Thymus zygis</i> L.	aerial parts	Spain	steam distillation	[A] thymol 46.69%, γ -terpinene 9.65%, <i>p</i> -cymol 19.56%, linalool 4.51%, carvacrol 3.04%, myrcene 1.87%, α -terpinene 1.72%, α -thujene 1.39%, terpinene-4-ol 1.01%, carvacrol methyl ether 0.34%
Wintergreen	<i>Gaultheriae aetheroleum</i>	<i>Gaultheria procumbens</i> L.	leaves	China	steam distillation	[A] methyl salicylate 98.90%
Winter savory	<i>Saturejae montanae aetheroleum</i>	<i>Satureja montana</i> L.	aerial parts	Southern Europe	steam distillation	[B] carvacrol (30–60%), <i>p</i> -cymene (10–20%), γ -terpinene (5–15%), borneol, carvone, β -caryophyllene [38]

were measured. Growth controls with only sterile blank filter paper discs and negative controls of EOs without bacterial inoculum were performed. Agar disc diffusion was run in duplicate. Since the blank filter paper discs had a radius of 3 mm, this was the smallest IZR to be determined.

Microdilution

Microdilution for determination of MIC and MBC was conducted for all 11 EOs against *P. multocida* isolates ($n = 10$) as described by Cermelli et al. [33] with broth and inoculum according to CLSI [34]. Cryopreserved bacteria were recovered at 36 °C on Columbia agar

► **Table 2** The composition of the seven EO mixtures.

EO mixtures		
2 EOs	1	70% lemongrass + 30% cinnamon cassia
	2	70% lemongrass + 30% thyme
	3	70% lemongrass + 30% wintergreen
	4	70% lemongrass + 30% winter savory
3 EOs	5	70% lemongrass + 30% cinnamon cassia + 10% thyme
	6	65% lemongrass + 20% winter savory + 15% cinnamon cassia
4 EOs	7	60% lemongrass + 15% cinnamon cassia + 15% winter savory + 10% thyme

with 5% sheep blood 24 h before performing microdilution. A two-fold serial dilution of EOs from 1.0–0.0039% was prepared in 96-well plates using CAMHB from Becton Dickinson with 0.5% Tween 20 from Merck Millipore as the emulsifier. Bacterial inoculum adjusted to 0.5 McFarland in sterile NaCl prediluted in CAMHB broth with 0.5% Tween 20 was seeded in prepared 96-well plates. Bacterial inoculum had a final dilution of 1 : 200 in 100 µL broth per well. A positive growth control with only CAMHB with 0.5% Tween 20 but no EO, and a negative control of the twofold serial dilution of EOs without bacterial inoculum was run alongside. The 96-well plates were sealed with a plastic sheet and incubated at 36 °C for 24 h. The MIC was determined visually as the lowest concentration of EO inhibiting visible bacterial growth in 96-well plates.

For determination of MBC, 100 µL of suitable wells were plated on Columbia agar with 5% sheep blood along with the positive and negative controls. The MBC was determined as the lowest concentration of EO inhibiting bacterial growth on inoculated agar plates.

Experimental design

In this study, activity of 11 EOs against *P. multocida* isolates (n = 10) was analysed using microdilution for the determination of MIC and MBC. Microdilution was performed in triplicate and the mean of each triplicate was used for further statistical analyses. Additionally, the 11 EOs and 7 EO mixtures were screened for their antimicrobial activity against bacterial isolates of *P. multocida* (n = 10), *M. haemolytica* (n = 10), and bacteria in the *Mannheimia* clade (n = 10). Agar disc diffusion was performed in duplicate, and the mean of each duplicate was used for further statistical analyses.

Statistical analyses

Analysis was conducted in “R” [35] and figures were produced using the package ggplot2 [36]. The Shapiro-Wilk test was used to test for normality. Each group of bacterial isolate-EO combination was analysed individually for normality. The IZR of *P. multocida* and spruce oil [W(9) = 0.830, p = 0.033], star anise oil [W(9) = 0.771, p = 0.006], and wintergreen oil [W(9) = 0.792, p = 0.012], as well as *M. haemolytica* and coriander oil [W(9) = 0.843, p = 0.048], eucalyptus oil [W(9) = 0.833, p = 0.036], spruce oil [W(9) = 0.366, p < 0.001], and star anise oil [W(9) = 0.836, p = 0.040] as well as

Mannheimia clade and spruce oil [W(9) = 0.837, p = 0.041] were not normally distributed.

Differences of EO activity for agar disc diffusion and microdilution of *P. multocida* were analysed with the Kruskal-Wallis test for global comparison of differences, followed by Dunn’s post hoc test with Benjamini-Hochberg adjustment for multiple pairwise comparison. Differences of susceptibility of *P. multocida* vs. *M. haemolytica* vs. *Mannheimia* clade against the same EO in agar disc diffusion were analysed with the ARTool of “R” [37] as follows: after the alignment rank transformation to allow for analyses of nonparametric data, a two-way ANOVA was performed for global comparison of differences, followed by a post hoc pairwise comparison with ART-C and Benjamini-Hochberg adjustment.

Supporting information

Supporting Information includes graphics on the EO mixture activity against *P. multocida*, *M. haemolytica*, and *Mannheimia* clade (Fig. 15) and on the activity of EO mixtures and their single EO constituents against *P. multocida*, *M. haemolytica*, and *Mannheimia* clade (Fig. 25).

Contributors’ Statement

Data collection: D. Bismarck, J. Becker, E. Müller, V. Becher, L. Nau, P. Mayer; design of the study: D. Bismarck, J. Becker, E. Müller, V. Becher, L. Nau, P. Mayer; statistical analysis: D. Bismarck; interpretation of the data: D. Bismarck, J. Becker, E. Müller, V. Becher, L. Nau, P. Mayer; drafting the manuscript: D. Bismarck, J. Becker; revision of the manuscript: E. Müller, V. Becher, L. Nau, P. Mayer.

Acknowledgements

We thank Prof. Dr. Mireille Meylan from the Clinic for Ruminants, and Prof. Dr. Vincent Perreten from the Institute of Veterinary Bacteriology, Vetsuisse-Faculty, University Bern, for support and supplying bovine bacterial isolates. Bacterial isolates were sampled within the “outdoor veal calf” project [financed by the National Research Programme 72 (NRP72) of the Swiss National Science Foundation, project 407240_167083] and isolated at the Institute of Veterinary Bacteriology, Vetsuisse-Faculty, University Bern. This study was funded by SaluVet GmbH and Laboklin GmbH&Co.KG.

Conflict of Interest

Philipp Mayer and Lisa Nau are employees of SaluVet GmbH, which provided the essential oils. Vera Becher was an employee of SaluVet GmbH when the study was conducted. The other authors declare that they have no conflict of interest.

References

- [1] Rice JA, Carrasco-Medina L, Hodgins DC, Shewen PE. *Mannheimia haemolytica* and bovine respiratory disease. Anim Health Res Rev 2007; 8: 117–128
- [2] Caswell JL. Failure of respiratory defenses in the pathogenesis of bacterial pneumonia of cattle. Vet Pathol 2014; 51: 393–409
- [3] Murray GM, O’Neill RG, McElroy MC, Earley B, Cassidy JP. Evolving views on bovine respiratory disease: An appraisal of selected key pathogens – Part 1. Vet J 2016; 217: 95–102

- [4] Dabo SM, Taylor JD, Confer AW. *Pasteurella multocida* and bovine respiratory disease. Anim Health Res Rev 2007; 8: 129–150
- [5] Confer AW, Ayalew S. *Mannheimia haemolytica* in bovine respiratory disease: Immunogens, potential immunogens, and vaccines. Anim Health Res Rev 2018; 19: 79–99
- [6] Friend SC, Wilkie BN, Thomson RG, Barnum DA. Bovine pneumonic pasteurellosis: Experimental induction in vaccinated and nonvaccinated calves. Can J Comp Med 1977; 41: 77–83
- [7] de Jong A, Thomas V, Simjee S, Moyaert H, El Garch F, Maher K, Morrissey I, Butty P, Klein U, Marion H, Rigaut D, Vallé M. Antimicrobial susceptibility monitoring of respiratory tract pathogens isolated from diseased cattle and pigs across Europe: The VetPath study. Vet Microbiol 2014; 172: 202–215
- [8] Portis E, Lindeman C, Johansen L, Stoltman G. A ten-year (2000–2009) study of antimicrobial susceptibility of bacteria that cause bovine respiratory disease complex—*Mannheimia haemolytica*, *Pasteurella multocida*, and *Histophilus somni*—in the United States and Canada. J Vet Diagn Invest 2012; 24: 932–944
- [9] Coetzee JF, Magstadt DR, Sidhu PK, Follett L, Schuler AM, Krull AC, Cooper VL, Engelken TJ, Kleinhenz MD, O'Connor AM. Association between antimicrobial drug class for treatment and retreatment of bovine respiratory disease (BRD) and frequency of resistant BRD pathogen isolation from veterinary diagnostic laboratory samples. PLoS One 2019; 14: e0219104
- [10] Timsit E, Hallewell J, Booker C, Tison N, Amat S, Alexander TW. Prevalence and antimicrobial susceptibility of *Mannheimia haemolytica*, *Pasteurella multocida*, and *Histophilus somni* isolated from the lower respiratory tract of healthy feedlot cattle and those diagnosed with bovine respiratory disease. Vet Microbiol 2017; 208: 118–125
- [11] Stanford K, Zaheer R, Klima C, McAllister T, Peters D, Niu YD, Ralston B. Antimicrobial resistance in members of the bacterial bovine respiratory disease complex isolated from lung tissue of cattle mortalities managed with or without the use of antimicrobials. Microorganisms 2020; 8: 288
- [12] Ayrlé H, Mevissen M, Kaske M, Nathues H, Gruetzner N, Melzig M, Walkenhorst M. Medicinal plants—prophylactic and therapeutic options for gastrointestinal and respiratory diseases in calves and piglets? A systematic review. BMC Vet Res 2016; 12: 89
- [13] Bouy M. Ätherische Öle für Wiederkäuer – Bewährte Indikationen und Fallbeispiele aus der französischen Nutztierpraxis. 33. Schweizerische Jahrestagung für Phytotherapie 2018 – Ätherische Öle und ihr therapeutisches Potential. Baden, Switzerland; 2018
- [14] Brendieck-Worm C, Klarer F, Stöger E. Heilende Kräuter für Tiere: Pflanzliche Hausmittel für Heim- und Nutztiere. 2nd ed. Bern: Haupt Verlag; 2018
- [15] Reichling J, Schnitzler P, Suschke U, Saller R. Essential oils of aromatic plants with antibacterial, antifungal, antiviral, and cytotoxic properties—an overview. Forsch Komplementmed 2009; 16: 79–90
- [16] Kalembe D, Kunicka A. Antibacterial and antifungal properties of essential oils. Curr Med Chem 2003; 10: 813–829
- [17] Amat S, Baines D, Timsit E, Hallewell J, Alexander TW. Essential oils inhibit the bovine respiratory pathogens *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni* and have limited effects on commensal bacteria and turbinate cells *in vitro*. J Appl Microbiol 2019; 126: 1668–1682
- [18] Amat S, Baines D, Alexander TW. A vapour phase assay for evaluating the antimicrobial activities of essential oils against bovine respiratory bacterial pathogens. Lett Appl Microbiol 2017; 65: 489–495
- [19] Salzmann BA. Antimikrobielle *In-vitro*-Wirkung von acht ätherischen Ölen auf *Pasteurella multocida* und *Mannheimia haemolytica* aus Nasentupferproben von Kälbern [Master Thesis]. Bern: Vetsuisse-Faculty; 2019
- [20] Janssen AM, Scheffer JJ, Baerheim Svendsen A. Antimicrobial activity of essential oils: A 1976–1986 literature review. Aspects of the test methods. Planta Med 1987; 53: 395–398
- [21] Pankey GA, Sabath LD. Clinical relevance of bacteriostatic versus bactericidal mechanisms of action in the treatment of Gram-positive bacterial infections. Clin Infect Dis 2004; 38: 864–870
- [22] Do TKT, Hadji-Minaglou F, Antonioti S, Fernandez X. Authenticity of essential oils. TrAC Trends Anal Chem 2015; 66: 146–157
- [23] Lis-Balchin M, Deans SG, Eaglesham E. Relationship between bioactivity and chemical composition of commercial essential oils. Flavour Fragr J 1998; 13: 98–104
- [24] Kissels W, Wu X, Santos RR. Short communication: Interaction of the isomers carvacrol and thymol with the antibiotics doxycycline and tilmicosin: *In vitro* effects against pathogenic bacteria commonly found in the respiratory tract of calves. J Dairy Sci 2017; 100: 970–974
- [25] LeBel G, Vaillancourt K, Bercier P, Grenier D. Antibacterial activity against porcine respiratory bacterial pathogens and *in vitro* biocompatibility of essential oils. Arch Microbiol 2019; 201: 833–840
- [26] Fararh KM, Farid SA, Abd EL-Hamied SS, El-Sharkawy RB. Clinicopathological changes in calves with respiratory diseases after treatment with essential volatile oil and other drugs. BVIMJ 2017; 33: 237–247
- [27] Ben-Arye E, Dudai N, Eini A, Torem M, Schiff E, Rakover Y. Treatment of upper respiratory tract infections in primary care: a randomized study using aromatic herbs. Evid Based Complement Alternat Med 2011; 2011: 690346
- [28] Cohen BM, Dressler WE. Acute aromatics inhalation modifies the airways. Effects of the common cold. Respiration 1982; 43: 285–293
- [29] Wu Q, Yu L, Qiu J, Shen B, Wang D, Soromou LW, Feng H. Linalool attenuates lung inflammation induced by *Pasteurella multocida* via activating Nrf-2 signaling pathway. Int Immunopharmacol 2014; 21: 456–463
- [30] Amat S, Timsit E, Baines D, Yanke J, Alexander TW. Development of bacterial therapeutics against the bovine respiratory pathogen *Mannheimia haemolytica*. Appl Environ Microbiol 2019; 85: e01359–01419
- [31] Becker J, Schüpbach-Regula G, Steiner A, Perreten V, Wüthrich D, Hausherr A, Meylan M. Effects of the novel concept 'outdoor veal calf' on antimicrobial use, mortality and weight gain in Switzerland. Prev Vet Med 2020; 176: 104907
- [32] Bismarck D, Schneider M, Müller E. Antibakterielle *In-vitro*-Wirksamkeit ätherischer Öle gegen veterinärmedizinisch relevante Keime klinischer Isolate von Hunden, Katzen und Pferden. Complement Med Res 2017; 24: 153–163
- [33] Cermelli C, Fabio A, Fabio G, Quaglio P. Effect of eucalyptus essential oil on respiratory bacteria and viruses. Curr Microbiol 2008; 56: 89–92
- [34] CLSI. Performance Standards for antimicrobial Disk and Dilution Susceptibility Tests for Bacteria isolated from Animals. 4th ed. CLSI supplement VET08. Wayne, PA: Clinical and Laboratory Standards Institute; 2018
- [35] R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria. 2021. Accessed November 2, 2021 at: <https://www.R-project.org/>
- [36] Wickham H. ggplot2: Elegant graphics for data analysis. New York: Springer; 2009
- [37] Elkin LA, Kay M, Higgins JJ, Wobbrock JO. An Aligned Rank Transform Procedure for Multifactor Contrast Tests. In: Nichols J, Kumar R, Nebeling M, eds. The 34th Annual ACM Symposium on User Interface Software and Technology. New York: ACM; 2021: 754–768
- [38] Hiller K, Melzig MF. Lexikon der Arzneipflanzen und Drogen, 2nd ed. Heidelberg: Spektrum Akademischer Verlag; 2010
- [39] World Health Organization. WHO monographs on selected medicinal plants, Volume 2. Geneva: World Health Organization; 2002
- [40] European Scientific Cooperative on Phytotherapy. ESCOP Monographs Online Series, Caryophylli Aetheroleum – clove oil. Exeter: European Scientific Cooperative on Phytotherapy; 2014
- [41] Nath SC, Saha BN, Bordoloi DN, Mathur RK, Leclercq PA. The chemical composition of the essential oil of *Cymbopogon flexuosus* (Steud) Wats. Growing in Northeast India. J Essent Oil Res 1994; 6: 85–87