Digital cholangioscopy-guided removal of an Ascaris worm from the biliary tree





▶ Fig. 1 Endoscopic ultrasound images showing: a a long, linear hyperechoic strip without any acoustic shadow within the common bile duct, consistent with the "strip" sign (dotted red line outlines the worm); b a central, longitudinal anechoic shadow, consistent with the "innertube" or "double-tube" sign (red arrows), within the common bile duct (dotted green line).



Fig. 3 Photographs of the 14-cm Ascaris that was pulled alive from the bile duct.



► Fig.2 Endoscopic ultrasound showing the papilla that had been opened up by the previous sphincterotomy.

A-37-year woman, who had undergone endoscopic retrograde cholangiopancreatography (ERCP) and sphincterotomy for common bile duct (CBD) stones followed by cholecystectomy 3 years ago, presented to us complaining of right upper quadrant pain for 3 days. Laboratory investigations showed raised liver enzymes (alanine transaminase 100U/L and alkaline phosphatase 320U/L), with normal bilirubin levels. An ultrasound of the abdomen showed a mildly dilated CBD with aerobilia. Endoscopic ultrasound (EUS) was performed, which showed long, moving, linear hyperechoic strips, without any acoustic shadow within the CBD, consistent with the "strip" sign and a central, longitudinal anechoic shadow, consistent with the "inner-tube" or "double-tube" sign, which suggests a diagnosis of biliary ascariasis (> Fig. 1). EUS examination of the ampulla showed an open biliary orifice (> Fig. 2) owing to the previous sphincterotomy, with flow of water within the CBD on ingestion and aerobilia.

The patient underwent ERCP with a therapeutic duodenoscope (TJF-180F; Olympus, Japan), which showed the previous papillary sphincterotomy, but no worm was seen at the papillary orifice. Contrast



Video 1 Endoscopic ultrasound showing the features of a live *Ascaris* within the bile duct and its cholangioscopy-guided removal.

was not injected as the patient had a history of contrast allergy. Digital single-operator cholangioscopy (DSOC; SpyGlass; Boston Scientific, USA) of the CBD was performed through the duodenoscope. DSOC showed a long, live, linear tubular worm occupying the whole CBD and piercing into the right anterior hepatic duct (**Video 1**). The worm was removed with forceps (SpyBite; Boston Scientific) under direct visualization (**Fig. 3**). Following the procedure, the patient was stable and albendazole was given as deworming therapy.

Biliary ascariasis is a common cause of pancreaticobiliary disease in tropical countries. Risk factors for biliary ascariasis include a history of cholecystectomy, choledocholithotomy, sphincteroplasty, or endoscopic sphincterotomy, and pregnancy [1]. Our patient had a history of both cholecystectomy and biliary sphincterotomy (▶ Fig. 2). ERCP plays an important role in the diagnosis of pancreaticobiliary ascariasis, as well as in its therapy by direct extraction of the worm [2]. In this case, biliary ascariasis was suspected on EUS and DSOC confirmed the diagnosis and also assisted with removal of the worm under direct visualization.

Endoscopy_UCTN_Code_CCL_1AZ_2AI

Competing interests

The authors declare that they have no conflict of interest.

The authors

Radhika Chavan Rushi C. Pipavat, Ravi Bokarvadia, Sanjay L. Rajput Department of Gastroenterology, Ansh Clinic, Ahmedabad, Gujarat, India

Corresponding author

Radhika Chavan, MD

Department of Gastroenterology, Ansh Clinic, Near Divine School and Hirabhai Tower, Maninagar, Ahmedabad, Gujarat, 380008, India drradhikachavan@gmail.com



- Khuroo MS, Rather AA, Khuroo NS et al. Hepatobiliary and pancreatic ascariasis. World J Gastroenterol 2016; 22: 7507–7517
- [2] Chavan R, Ramchandani M, Nabi Z et al. Recurrent acute pancreatitis due to pancreatic duct ascariasis in a young man. Endoscopy 2018; 50: E292–E293

Bibliography

Endoscopy 2022; 54: E999–E1000 DOI 10.1055/a-1887-5888 ISSN 0013-726X published online 4.8.2022 © 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (https:// creativecommons.org/licenses/by-nc-nd/4.0/) Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS https://eref.thieme.de/e-videos



Endoscopy E-Videos is an open access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and wavers acc. to HINARI are available.

This section has its own submission website at

https://mc.manuscriptcentral.com/e-videos