A. Deepthi et al.



270

#### Review

# Recent Progress on the [3+2] Cycloaddition Route for the Synthesis of All-Carbon Quaternary Stereocentres

Ani Deepthi<sup>\*</sup> Maneesh Mohan Meenakshy C. Balachandran

Department of Chemistry, University of Kerala, Kariavattom campus, Thiruvananthapuram 695581, Kerala state, India anideepthi@gmail.com





#### License terms: (cc)(i) = (s)

© 2022. The Author(s). This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes or adapted, remixed, transformed or built upon. (https://creativecommons.org/licenses/by-nc-nd/4.0/)

**Abstract** Construction of all-carbon quaternary centres is an important task in organic synthesis. In spite of the challenges associated with Csp<sup>3</sup>–Csp<sup>3</sup> bond construction in a sterically constrained environment, significant advances have been made in this area. Among the latter, both catalytic and noncatalytic [3+2] cycloaddition approaches have gained wide attention recently. This short review summarizes the [3+2] cycloaddition reactions reported during the period 2016–2022 for the synthesis of molecules possessing one or more all-carbon quaternary stereocentres.

**Key words** all-carbon, quaternary, stereocentres, [3+2] cycloaddition, cyclization

### 1 Introduction

All-carbon quaternary centres refer to stereogenic carbon centres attached to four different neighbouring carbons.<sup>1,2</sup> Such carbon centres have a special importance owing to their ubiquitous presence in natural products and to their specificity in biological systems due to their structural diversity and enhanced conformational constraints, which makes them suitable targets for protein binding. For example, spirocyclic compounds have found increased utility in drug discovery because of their three-dimensional structural uniqueness.<sup>3</sup> Figure 1 shows some examples of biologically active molecules containing all-carbon quaternary spiro-centres. Fredericamycin A,<sup>4a</sup> first isolated from *Streptomyces griseus* ATCC49344, is known for its potent activity against P388 mouse leukaemia, CD8F mammary tumours and B16 melanoma *in vivo*; whereas coleophomone D,<sup>4b</sup> isolated from *Stachybotrys parvispora Hughes*, is known for its antibacterial and antihypertensive properties. Elatol displays antibiofouling, antibacterial and antifungal properties and is also cytotoxic against HeLa and Hep-2-human carcinoma cell lines.<sup>5a</sup> Colletoic acid is an inhibitor of human 11β-hydroxy steroid dehydrogenase type I (11β-HSD1 inhibition is a new therapeutic approach for type I diabetes mellitus),<sup>5b</sup> and horsfiline, an oxindole alkaloid, is a therapeutic agent.<sup>5c</sup>

Given the importance of generating quaternary stereogenic carbon centres, especially as spirocyclic molecules, numerous efforts have been made for developing efficient methods for their stereocontrolled synthesis.<sup>6-11</sup> Asymmetric [3+2] cycloadditions have created a niche in organic chemistry.<sup>12</sup> Recently Wang and Liu catalogued all-carbon [3+2] cycloaddition methods for natural product synthesis in which they invoke the importance of the strategy to create complex scaffolds of biological interest.<sup>13</sup> In this review, we present selected [3+2] cycloaddition reactions that lead to formation of one or more all-carbon quaternary stereocentres reported during the period 2016-2022. These have been classified into different categories (i) based on the substrate used to generate the 1,3-dipolar species, (ii) those involving allyl-palladium intermediates, (iii) those involving phosphine-allenoate zwitterionic intermediates, and (iv) those involving radical pathways, nitrones, nitrile oxides and nitronate salts, although certain examples can be included in more than one group in this classification.



#### **Biographical Sketches**



Ani Deepthi was born in Kerala, India. After securing an MSc degree in Applied Chemistry from Cochin University of Science and Technology, she undertook her PhD degree working under the guidance of Dr Vijay Nair in CSIR-NIIST, Thiruvananthapuram and carried out post-doctoral studies at the National University of Singapore with Prof. Suresh Valiyaveettil. Currently, she is an Assistant Professor at the Department of Chemistry, University of Kerala. Her research interests include synthesis of small molecules

Government Guest Lecturer at Sree Narayana College, Kollam and later worked as a Project Assistant Ш at CSIR-NIIST. Thiruvananthapuram, Kerala. He secured his MPhil. degree from possessing bioactive or photoactive properties, chemosensors, and isolation and semisynthetic modification of natural products. She has 32 publications in-peer reviewed journals and two students have secured their PhD under her guidance.



Maneesh Mohan was born in Alappuzha, Kerala, India. He received his BSc. degree in 2011 from Sree Narayana College, Cherthala and MSc. degree in 2016 from Sree Narayana College, Chengannur. He then worked as a

the Department of Chemistry,

University of Kerala and is currently pursuing his Ph.D under the guidance of Dr. Ani Deepthi. His research interests include synthesis and in vitro anticancer evaluation of beta carboline-based spiroheterocycles.



Meenakshy Chandrika Balachandran was born in Kuzhithurai, Tamil Nadu, India. She received her B.Sc. degree in 2018 from University College, Palayam, Thiruvananthapuram and M.Sc. Degree in 2020 from H.H. The Maharaja's College for Women, Vazhuthacaud, Thiruvananthapuram. She qualified in the joint CSIR-UGC NET in 2020 and is currently carrying out her doctoral research under the quidance of Dr. Ani Deepthi at the Department of Chemistry, University of Kerala, Kariavattom. Her research focuses on the synthesis of dispiro-heterocyclic compounds and their anti-cancer properties.

## 2 Classification based on Substrate used to Generate the 1,3-Dipole

#### 2.1 Suitably Substituted Cyclopropanes

SvnOpen

Previous studies have shown that vinyl cyclopropanes (VCPs), when catalytically activated by palladium or by other transition metals, generate a 1,3-dipolar intermediate that can, in turn, form five-membered rings with various dipolarophiles via [3+2] cycloaddition reactions.<sup>14</sup> However, due to the challenges associated with the short lifetime of the intermediate in the presence of phosphine ligands. asymmetric versions of these reactions have great significance.<sup>15</sup> In this context, recently developed synergistic catalvsis proved to be advantageous, wherein the vinvl cvclopropane is activated by a palladium catalyst while the dipolarophile is activated by a second catalyst. For instance, Jørgensen and co-workers in 2016 developed a protocol in which the activated vinyl cyclopropane reacts with the dipolarophile that is, in turn, activated by an organocatalyst. The methodology resulted in the synthesis of densely substituted cyclopentanes with up to four contiguous stereocentres in high yields and with excellent stereoselectivities. For example, vinyl cyclopropane 1 reacted with cinnamaldehyde 2 in the presence of benzoic acid in acetonitrile to yield the product 3 (Scheme 1). Here, the vinyl cyclopropane is activated by palladium while the  $\alpha,\beta$ -unsaturated aldehyde is activated as the iminium ion formed by reaction with the organocatalyst 4. More specifically, oxidative addition of Pd(0) catalyst facilitates ring opening of the vinyl cyclopropane, yielding a  $\pi$ -allyl palladium intermediate **5** and, in parallel, condensation of catalyst **4** and cinnamaldehyde **2** leads to the formation of the iminium ion **6**. Intermediates **5** and **6** combine to generate the [3+2] cycloadduct **3**.<sup>16</sup>

Concurrently, there were also successful efforts by Vitale and co-workers to employ iminium/enamine organocatalysis also with Pd(0) activation of vinyl cyclopropanes for the enantioselective synthesis of polysubstituted cyclopentanes by the formal [3+2] cycloaddition of vinyl cyclopropanes with enals,<sup>17</sup> even though generation of molecules containing all-carbon quaternary centres was not reported.

In 2020, a synergistic bimetallic catalytic system comprising palladium and rhodium catalysts was used by Du and co-workers for the enantioselective synthesis of multisubstituted spirocyclopentane oxindoles containing an allcarbon quaternary stereocentre. In this reaction,  $\alpha$ , $\beta$ -unsaturated 2-acyl imidazole **8** underwent [3+2] cycloaddition with spirovinyl cyclopropanyl-2-oxindole **7** in the presence of palladium and chiral rhodium catalysts to yield the 3-spirocyclopentane-2-oxindole derivative **9** in high yield (Scheme 2). Here, the zwitterionic  $\pi$ -allyl palladium intermediate formed from **7** attacks the *Re*-face of the bidentate *N*,*O*-coordinated intermediate generated by the coordination of rhodium to compound **8**, facilitating the formation of the product **9** as the major isomer (Scheme 2).<sup>18</sup>

Enantioselective ring-openings of other suitably substituted cyclopropanes have also emerged as a powerful strategy for five-membered ring construction.<sup>19</sup> An intramolecular [3+2] annulation reaction of cyclopropanes was report-



272

THIEME



		273	
		THIEME	
SynOpen	A. Deepthi et al.	OPEN ACCESS	Review

ed by Chen, Yang and co-workers in 2016 (Scheme 3).<sup>20</sup> In this reaction, an amide linker was used for the annulation and  $TiCl_4$  was used as the Lewis acid. The coordination of the two carbonyl groups of substrate **10** to titanium(IV) probably facilitates the ring-opening of the cyclopropane, generating a dicarbonyl anion that, in turn, attacks the enone moiety to furnish the six-membered lactam ring and finally leads to dihydroquinolinone **11**.



Donor–acceptor cyclopropane **12** was used for the synthesis of pharmaceutically relevant azabicyclo[3.2.1]octane **15**, bearing two all-carbon quaternary stereogenic centres at the bridgehead positions, by reduction of imine **14** and subsequent deprotection and intramolecular amidation. The imine was, in turn, synthesized by MgI<sub>2</sub> catalysed [3+2] cycloaddition of donor–acceptor cyclopropane **12** with azadiene **13** in dichloromethane (Scheme 4).<sup>21</sup> Wang and co-workers in 2019 used the [3+2] annulation strategy to construct an all-carbon stereocentre by conducting the 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) mediated reaction of cyanocyclopropane carbonate **16** and (*E*)-3-aryl-2-cyano acrylate **17**. The reaction takes place by the initial deprotonation of cyclopropane **16**, which ring opens to form an ylide intermediate. The [3+2] annulation of the ylide to acrylate **17** leads to the cycloadduct **18** after a tautomerization step (Scheme 5).<sup>22</sup>

Aminocyclopropanes have recently emerged as suitable reaction partners that can undergo annulation with alkenes and cyclopropenes under photocatalysis. In 2019, Waser and co-workers reported the [3+2] annulation reaction of cyclopropenes with cyclopropyl aniline **20** using photocatalvsis. The catalvst used was 2.4.5.6-tetrakis(diphenvlamino)isophthalonitrile (4DPAIPN). The transformation proved to be highly efficient, and diastereoselectivity was further improved by choosing bulky cyclopropyl aniline and difluorocyclopropenes (Scheme 6).<sup>23</sup> Subsequently, in 2020, a highly diastereoselective and enantioselective [3+2] cycloaddition of cyclopropyl amine with  $\alpha$ -alkyl styrenes was reported by Ooi and co-workers using photocatalysis to yield cyclopentanes containing all-carbon quaternary stereocentres. Here, iridium-polypyridyl complexes were used as photocatalysts and the reaction is initiated by capture of the anionic component of the photocatalyst by the cyclopropyl urea 22,







generating a chiral supramolecular ion pair. Under irradiation, single electron transfer (SET) from the substrate to the excited-state cationic iridium generates a radical cation that undergoes stereoselective bond formation with the  $\alpha$ alkyl styrene 23 within the restrictions of the asymmetric environment created by the chiral anion.



A representative reaction for synthesis of the cyclopentane 24 is depicted in Scheme 7.<sup>24</sup> The Waser group also reported the use of the amino cyclopropane monoester 25 for [3+2] annulation with suitably substituted indoles using silyl bistriflimide as catalyst (Scheme 8). The method was successfully applied for the construction of a non-symmetrical all-carbon quaternary centre at the acceptor position of the cyclopropane in good yield and diastereoselectivity, as exemplified in Scheme 8.25

Construction of an all-carbon quaternary stereocentre at the  $\alpha$ -position of aza-arenes was reported by Jiang and co-workers by synergistic photoredox and Brönsted acid catalysis. Scheme 9 shows a representative reaction of cyclopropyl aniline 20 with 2-(1-phenylvinyl)pyridine 28 that leads to 29. It was found that there were remarkable differences in enantioselectivity when using electron-withdrawing or electron-donating substituents on the aromatic ring





of the 2-(1-arylyinyl)pyridine, which led to the proposition of a ternary transition state with the chiral Brönsted acid acting as the bifunctional catalyst. The chiral Brönsted acid catalyst used was an iminodiphosphoric acid that co-catalyses the reaction along with the photocatalyst, dicyanopyrazine (DPZ).26



Scheme 9 Photocatalysed enantioselective pyridine tethered cyclopentane synthesis



Scheme 7 Photocatalysed enantioselective urea-tethered cyclopentane synthesis

	275		
SynOpen A. Deep	pthi et al.	R	leview

#### 2.2 3-Homoacyl Coumarins

Lin and co-workers have utilized the all-carbon 1,3-dipole precursor 3-homoacyl coumarin **30**. in stereoselective [3+2] cycloaddition reactions in the presence of a squaramide catalyst. Initially indandione alkylidene 31 was used as the dipolarophilic partner, which led to the synthesis of coumarin/indandione fused spirocyclopentanes 32, bearing four contiguous stereocentres.<sup>27a</sup> Later these workers used alkylidene oxindoles as the dipolarophilic partner to yield a cycloadduct containing five contiguous stereocentres of which one is an all-carbon guaternary spirocentre.<sup>27b</sup> The reaction takes place via initial activation of the homoacvl coumarin **30** by deprotonation to provide a conjugate acidbase pair with H-bonding interactions. The Re-face of activated coumarin then adds to the Re-face of the alkylidene oxindole **33** to generate a Michael adduct that ultimately leads to product 34 (Scheme 10). Lin's group has also employed the strategy for the enantioselective synthesis of spiropyrazolone fused cyclopenta[c]chromen-4-ones.<sup>27c</sup> The reaction occurs by [3+2] cycloaddition of 3-homoacyl coumarin **30** with  $\alpha$ , $\beta$ -unsaturated pyrazolone **35** in the

presence of a cinchona alkaloid-derived hydrogen-bonding catalyst and benzoic acid. Excellent yields of spiropyrazolones of the type **36** were obtained in a highly enantioselective manner. All these reactions are summarized in Scheme 10.

#### 2.3 Isatin-Based (Trifluoromethyl)imines

Isatin-based (trifluoromethyl)imines were reported to undergo [3+2] annulations with alkenes efficiently in the presence of metal catalysts in conjunction with organocatalysts. A Brönsted base and Lewis acid co-operatively catalysed asymmetric 1,3-dipolar cycloaddition reaction was reported to yield a series of trifluoromethyl substituted 2,3pyrrolidinyl dispiro-oxindoles with high enantioselectivity. The reaction was initiated by deprotonation of **37** by the complex generated from the organocatalyst and  $Et_2Zn$ , accompanied by release of ethane. This was followed by coordination of methylene indolinone **38** to the zinc from the less-hindered face. Michael addition followed by Mannich reaction led to an intermediate complex, from which the final product **39** was released by proton exchange and the catalytic cycle continued (Scheme 11).<sup>28</sup>





In another report, isatin-based (trifluoromethyl)imines were reported to undergo [3+2] cycloaddition with unsaturated 4-benzylidene chromanones **40** promoted by the organocatalyst DABCO. The authors could also achieve up to 69% enantiopurity for the spirooxindole-chromanone hybrid **41** using Takemoto's bifunctional thiourea catalyst **42**, via formation of an '*exo*' transition state (Scheme 12).<sup>29</sup> Concurrently Knipe and co-workers reported the use of cinchona-derived thiourea catalyst **43** for the synthesis of spiropyrrolidine oxindoles with excellent enantioselectivities. Some of the examples reported contained molecules with all-carbon quaternary stereocentres, as exemplified in the synthesis of **45** and **47** (Scheme 13).<sup>30</sup>

#### 2.4 Iminoesters and Isocyanoesters

Zhang and co-workers in 2017 reported a copper(I) catalysed asymmetric *exo*-selective [3+2] cycloaddition of  $\beta$ trifluoromethyl  $\beta$ , $\beta$ -disubstituted enone **49** with azomethine ylides (generated from glycine ketinimine 48), leading to the synthesis of chiral pyrrolidines bearing a trifluoromethylated quaternary carbon centre. The chiral ligand used in the reaction was (S)-MeO-DTBM-Biphep 53 and the copper salt used was Cu(CH<sub>3</sub>CN)<sub>4</sub>BF<sub>4</sub> (Scheme 14). The reaction was found to be general for a wide range of enones and acyclic azomethine ylides. Moreover, the products formed could be oxidized to 3H-pyrroles using 2,3-dichloro-5,5-dicyano-1,4-benzoquinone (DDQ) and converted into *N*-hydroxy pyrroles and nitrones using *m*-CPBA in varying efficiencies.<sup>31</sup> Later, the same group reported the Cu(I)-Ming-Phos-catalysed enantioselective [3+2] cycloaddition of glycine ketinimine with  $\beta$ -trifluoromethyl enone **50**, which yielded the highly functionalized pyrrolidine **52** containing an all-carbon guaternary stereocentre in 95% enantiomeric excess (Scheme 14).<sup>32</sup>

Subsequently, ligand-controlled [3+2] cycloaddition of iminoesters leading to chiral pyrrolidines with adjacent or discrete quaternary stereocentres with at least one all-car-



Scheme 12 DABCO mediated all-carbon spirocentre formation



Scheme 13 Use of thiourea catalyst





bon stereocentre was reported by the same group. Scheme 15 depicts a representative reaction of  $\alpha$ -methyl-iminoester **55** and  $\beta$ , $\beta$ -disubstituted enone **56** in the presence of a copper catalyst and chiral ligands L1 or L2, yielding 57a or 57b as the major product, depending on the ligand. Computational studies provided insights into the regioselective control. When ligand L1 was used, the phosphorus and nitrogen atoms of the ligand remained coordinated to the Cu(I) throughout the process, yielding 57a in higher amounts, while a switch in regioselectivity was observed when L2 was used due to formation of a Cu-O<sup>enone</sup> bond, with the amine nitrogen atom of L2 dissociating from Cu(I) centre.<sup>33</sup> The same group has also reported the synthesis of optically active dihydropyrroles containing an all-carbon guaternary stereocentre by copper-catalysed [3+2] cycloaddition of 56 with isocyanoesters of the type 58, and maximum yields and enantioselectivities were obtained using (R)-DTBM-Seg-Phos as catalyst. Scheme 16 shows a representative example for the synthesis of the dihydropyrrole **59**.<sup>34</sup>

#### 2.5 Aldehydes and Isatins

Aldehydes and isatins are versatile substrates to generate azomethine ylides. For instance, construction of an allcarbon spiro quaternary centre was reported by Boudriga and co-workers in 2019 during the diastereoselective synthesis of dispiropyrrolo[2,1-a]isoquinoline-fused pyrrolidine-2,5-diones by [3+2] cycloaddition of the cyclic diketone-based tetrahydroisoquinolinium N-ylide. In this reaction, the azomethine ylide formed from isatin 60 and 1,2,3,4-tetrahydroisoquinoline **61** approaches the (E)-3arylidene-1-phenyl-pyrrolidine-2,5-dione 62 in an exomanner to yield products 63 and 64, both containing a spiro quaternary carbon centre, as depicted in Scheme 17.<sup>35</sup> Concurrently, Yan and co-workers reported that acetic acid can act as a catalyst to facilitate the formation of azomethine vlides from aromatic aldehvdes and pyrrolidine, which, in turn, undergo [3+2] cycloaddition with 3-arylidene indolin-2-one or with 2-arylidene-1,3-indanedione to yield the corresponding functionalised pyrrolidines containing a spiro all-carbon quaternary stereocentre. In this reaction, the iminium ion **70**, formed from aromatic aldehyde and pyrrolidine, undergoes a [1,3]-hydride shift vielding an enamine intermediate 72. The latter then undergoes an aldol type reaction with a second molecule of aldehyde 65 followed by dehydration and deprotonation, vielding a conjugated azomethine ylide 67 that finally participates in [3+2] cycloaddition with the alkene 68 to yield cycloadduct 69, as shown in Scheme 18.36











#### 3 Reaction of Allyl Palladium Intermediates

Palladium-catalysed [3+2] cycloaddition of 5-vinyl oxazolidinones and trisubstituted alkenes using chiral ammonium-phosphine ligands was reported by Takashi and coworkers in 2016. The reaction takes place by an initial intermolecular addition of an allyl palladium species to the alkene, generating a zwitterionic intermediate. Subsequent ring closure and bond formation between the two reactive sites of the intermediate yields a product possessing three contiguous stereogenic centres, including two all-carbon quaternary centres. A phosphine ligand incorporating a quaternary ammonium halide component was used to assist Pd-halide contact. A representative reaction of oxazolidinone **73** with (E)-ethyl 2-cyano-3-phenylacrylate **74** leading to product **75** is shown in Scheme 19.<sup>37</sup>

Nucleophilic 1,3-dipolar  $\pi$ -allyl palladium intermediates generated from vinyl ethylene carbonates have been found to be efficient reaction partners. For instance, synthesis of spirooxindoles with two contiguous all-carbon stereocentres was reported by Hu and co-workers by the palladium-catalysed [3+2] cycloaddition of methylene indolinone with vinvl ethylene carbonate. The reaction takes place through the initial formation of the Pd- $\pi$ -allyl intermediate 79, which undergoes [3+2] cycloaddition with methylene indolinone 76 by intramolecular nucleophilic attack to yield 3,3'-tetrahydrofuryl spirooxindole 78 (Scheme 20).<sup>38</sup> Recently, the 1,3-dipolar  $\pi$ -allyl palladium intermediate generated from vinyl ethylene carbonate was also reacted with chalcones under rhodium catalysis, leading to formation of tetrahydrofuran derivatives possessing an all-carbon quaternary stereocentre. The asymmetric bimetallic catalysis was achieved using a chiral rhodium complex catalyst along with Pd<sub>2</sub>(dba)<sub>3</sub>·CHCl<sub>3</sub> and was demonstrated using  $\alpha$ , $\beta$ -unsaturated 2-acyl imidazole **81** and racemic phenyl vinyl carbonate **77** as cycloaddition partners. The reaction takes place by the activation of the  $\alpha$ , $\beta$ -unsaturated 2acyl imidazole substrate **81** by the chiral rhodium complex through bidentate coordination. The zwitterionic  $\pi$ -allyl intermediate approaches the double bond of the rhodium complex through its *Si*-face. Subsequent coordination-dissociation and electron neutralization generates an intermediate that undergoes substitution with **81**, releasing the target 1,2,3,4-tetrahydrofuran molecule **82** (Scheme 21).<sup>39</sup>











#### SynOpen 2022, 6, 270–285



Very recently, Zi and co-workers used vinyl methylene cyclic carbonates of the type **83** to generate vinyl-substituted palladium-oxyallyl species that undergo enantioselective inverse electron-demand [3+2] cycloaddition with electron-deficient nitroalkenes of type **84**. A hydrogenbond-donating ligand was used to construct the cyclopentanone **86** containing an all-carbon quaternary stereocentre in a highly stereoselective manner. The optimised ligand for achieving maximum enantiomeric excess was found to be Fe-Ur-Phos **85**, which contains a urea moiety. The latter facilitates the hydrogen-bond formation between the chiral catalyst and nitro group of **84**, which imparts the stereocontrol of the reaction (Scheme 22).<sup>40</sup>

Liu and co-workers recently reported cooperative catalysis mediated by a palladium catalyst and urea-tertiary amine organocatalyst for the [3+2] cycloaddition between  $\gamma$ -methylidene- $\delta$ -valerolactones of type **87** and  $\beta$ -nitrostyrene **88**. The reaction occurs by the initial generation of a 1,4-dipole from **87** by oxidative addition of palladium, facilitated by CO<sub>2</sub> release and followed by intermolecular Michael addition to the species generated from the nitroolefin. The strong hydrogen bonding between the urea-tertiary amine catalyst **89** and the nitro-olefin helps in the *Re*face attack of the zwitterionic  $\pi$ -allyl intermediate onto the nitro-olefin, providing good control of the stereochemistry of the all-carbon quaternary stereocentre. Subsequent intramolecular cyclisation affords the final product **90** (Scheme 23).<sup>41</sup>

### 4 Reaction of Phosphine-Allenoate Zwitterionic Species

It is well known that phosphines can add to alkynoates and allenoates to generate zwitterions that, in turn, can undergo [3+2] cycloaddition with alkenes and alkynes.<sup>42</sup> Asymmetric versions of such reactions using chiral phosphine catalysts have been widely explored<sup>43</sup> and the strategy has been utilised for the synthesis of all-carbon quaternary stereocentres.<sup>44</sup> An enantioselective [3+2] annulation of  $\alpha$ -substituted allenoates of the type **91** with  $\beta$ , $\gamma$ -unsaturated *N*-tosyl imine **92** in the presence of a phosphine catalyst was reported by Lu and co-workers. The reaction resulted in the formation of cyclopentene 93, containing an all-carbon guaternary centre. The reaction takes place via initial activation of the allenoate by attack of the phosphine, generating intermediate 94 that then undergoes [3+2] annelation with the *N*-tosyl imine. leading to product **93** after proton shift (Scheme 24).45 Recently Liu and co-workers used this strategy for the regioselective synthesis of CF<sub>3</sub>substituted guaternary carbon-centred molecules in moderate to good yields. The reaction occurs through initial nucleophilic addition of the phosphine onto the allenoate, vielding the zwitterionic intermediate **98** that undergoes  $\alpha$ addition to the double bond of 96 to yield the anionic intermediate 99, which, in turn, cyclises to 100. Final 1,2-proton transfer and  $\beta$ -elimination of the catalyst yields the target cyclopentene 97 (Scheme 25).46







Scheme 24 [3+2] Annulation of N-tosylimines with allenoates



Yao and co-workers reported a stereoselective and enantioselective trimerization of  $\gamma$ -aryl-3-butynoates catalysed by L-isoleucine-derived amide phosphine to yield cyclopentenones bearing an all-carbon quaternary stereocentre. The reaction occurs by addition of the phosphine to the allenoate (formed by isomerisation of the alkyne 102) to form zwitterionic intermediate 105, which, in turn, reacts with another molecule of allenoate to vield intermediate 106 followed by intramolecular Michael addition to yield **107.** After a 1,2-*H* shift and expulsion of the catalyst, **109** is obtained, leading eventually to the final product 104 containing an all-carbon quaternary stereocentre (Scheme 26).47 Very recently Takao and co-workers also utilised this strategy for the synthesis of cyclopentene molecules possessing all-carbon quaternary stereocentres using Oppolzer's camphor sultam 113 as one of the reaction partners.<sup>48</sup> The camphor sultam acts as a chiral auxiliary in this reaction, which is subsequently removed by chemoselective hydrolysis using alkaline hydrogen peroxide to furnish carboxylic acid 115 in 94% yield (Scheme 27). The high





regioselectivity is probably due to the bulkiness of the camphor sultam. The authors have also applied this strategy for the formal synthesis of R-(–)-puraquinonic acid.

In another earlier report, Morita–Baylis–Hillman (MBH) carbonates were used instead of allenoates to generate zwitterionic intermediates using nucleophilic phosphines. The latter reacted with MBH carbonate **117** to yield the salt **119** that, in turn, was deprotonated by the *tert*-butoxide to form species **120**. The dispirobisoxindole **118**, containing two all-carbon quaternary stereocentres, was formed by [3+2] cycloaddition of isoindigo **116** with **120** through formation of intermediates **121** and **122** (Scheme 28).<sup>49</sup>

#### 5 Miscellaneous Cycloaddition Reactions

In 2017, Li and co-workers reported that an allene-tethered electron-deficient olefin underwent intramolecular [3+2] cycloaddition in the presence of benzene thiol to yield [3.3.0]bicyclic systems containing *vicinal* quaternary carbon stereocentres. The intramolecular cascade reaction is initiated by an electrophilic benzene thiol radical **123**, affording a thermodynamically stable tertiary radical **124**, by adding to the central sp-carbon atom of the allene **121**. Subsequent attack of this radical to the  $\alpha$ , $\beta$ -unsaturated double bond via a 5-*exo-trig*-cyclisation led to radical intermediate **125** that then underwent a 5-*endo-trig*-cyclisation to generate **126**. The latter in the next step abstracts a hydrogen atom from the benzene thiol and is converted into the final product **122** (Scheme 29).<sup>50</sup>

An intramolecular silvl-nitronate cycloaddition was reported by Veselovsky and co-workers for the enantioselective synthesis of substituted cyclopentanones containing an all-carbon guaternary stereocentre. Henry reaction of 4methyl pentenal **127** with nitromethane in the presence of the chiral catalyst 128 yields the unsaturated nitro alcohol **129.** The latter is then converted into silvl ether **130** that, treated with 1,1,1,3,3,3-hexamethyldisilazane when (HMDS), yielded cyclopentaisoxazolidinone 132 through formation of the nitronate 131 by a stereoselective intramolecular cycloaddition with trans-disposition of the OTBS substituent relative to the annulated isoxazolidinone ring. Further transformation of 132 by exposure to sodium methoxide led to ring opened oximes 134a and 134b (anti/syn ratio 12:1 by <sup>1</sup>H NMR analysis) through nitroso tautomerism. Final deoximation of the oximes by tritura-





tion with periodic acid resulted in the cycloalkanone **135** (Scheme 30).<sup>51</sup>

A formal 1,3-dipolar cycloaddition of 3-ylidene oxindoles of the type **136** with aryl diazomethane **140** (generated from benzaldehyde **65** and tosyl hydrazine **137**) was reported by Babu and co-workers in 2018.<sup>52</sup> In this reaction, the intermediate cycloadduct **141** underwent decomposition in the aprotic solvent to deliver 3-spirocyclopropyl-2oxindole **138**, containing an all-carbon quaternary stereocentre. It was also observed that, in the presence of protic solvent, the cycloadduct **141** tautomerizes to form **142** which, in turn, is oxidised to the corresponding spiropyrazole oxindole. The latter then undergoes spontaneous rearrangement to pyrazoloquinazolinone **139**. Thus, a solventcontrolled switchable product selectivity was demonstrated, as depicted in Scheme 31.

Gao and co-workers, in 2019, reported the application of a [3+2] cycloaddition strategy for the construction of the core skeleton of calyciphylline A alkaloids. Herein, nitroneinduced [3+2] cycloaddition was used for the construction of the *cis*-hydroindole A–C rings containing an all-carbon quaternary centre at C5. Scheme 32 shows a representative construction of the stereogenic C5 centre of an important precursor to himalensine  $A^{53,54}$  in which a cycloheptene substrate **143** is treated with *N*-benzylhydroxylamine resulting in a nitrone intermediate that, on intramolecular 1,3-dipolar cycloaddition through 6-*exo*-cyclisation with the  $\alpha$ , $\beta$ -unsaturated ester, results in cycloadduct **144**, an important precursor containing the all-carbon quaternary centre of himalensine A **145**.

Gao and co-workers, in 2019, reported the asymmetric total synthesis of (-)-viridin and (-)-viridiol (Figure 2), both containing a quaternary all-carbon stereocentre at C-10. The strategy involved the synthesis of compound 146 from L-ribose, which was then transformed into the unsaturated ester 147 by ruthenium-catalysed cross-metathesis. Treatment of 147 with hydroxylamine hydrochloride generated oxime 148, which was converted into nitrile oxide 149 with chloramine-T. Subsequent intramolecular [3+2] cycloaddition generated isooxazoline 150, which, on reductive hydrogenolysis and hydrolysis, yielded ketone 151. Wittig reaction of the latter produced compound 152. which, on reaction with dimethyl hydroxylamine hydrochloride (DMHH), yielded the Weinrib amide 153. This reacted with the anion of dihydroindenol 154 to yield 155 as a mixture of diastereomers at C17 (see Figure 2). Compound 155 was







converted into the desired tetracyclic core 156, containing an all-carbon quaternary stereocentre at C10, using a metal-catalysed hydrogen atom transfer (MHAT) strategy with the help of cobalt-salen catalyst 158 and phenyl silane. Selective removal of the TES group using Dowex resin followed by oxidation with Dess-Martin periodinane (DMP) vielded compound 157, which was converted into viridin and viridiol (Scheme 33).55

CO2E



Figure 2 Structures of viridin and viridiol

#### Conclusion 6

144

Bn

= CO<sub>2</sub>Et

Effective ways of constructing all-carbon stereocentres have been reported in the past five years using [3+2] cycloaddition strategies involving metal-catalysed, organocatalysed, photocatalysed, base/acid-mediated, and thermal [3+2] cycloadditions, leading to products containing one or more all-carbon stereocentres. We believe that this review along with other reviews in this period, 56-61 will provide insight for organic chemists wishing to construct molecules containing all-carbon quaternary stereocentres.

(–)-himalensine A

#### **Conflict of Interest**

The authors declare no conflict of interest.

#### **Funding Information**

M.M. and M.C.B. thank the University of Kerala for Research Fellowships.

#### Acknowledgment

The authors thank the University of Kerala for provision of facilities.



284

Scheme 33 Sequential synthesis of viridiol precursor

#### References

- (1) Fuji, K. Chem. Rev. 1993, 93, 2037.
- (2) Christoffers, J.; Mann, A. Angew. Chem. Int. Ed. 2001, 40, 4591.
- (3) Bora, D.; Kaushal, A.; Shankaraiah, N. Eur. J. Med. Chem. 2021, 215, 113263.
- (4) (a) Chen, Y.; Luo, Y.; Ju, J.; Wendt-Pienkowski, E.; Rajski, S. R.; Shen, B. J. Nat. Prod. 2008, 71, 431. (b) Nicolaou, K. C.; Montagnon, T.; Vassilikogiannakis, G. Chem. Commun. 2002, 2478.
- (5) (a) White, D. E.; Stewart, I. C.; Grubbs, R. H.; Stoltz, B. M. J. Am. Chem. Soc. 2008, 130, 810. (b) Sawada, T.; Nakada, M. Org. Lett. 2013, 15, 1004. (c) Jossang, A.; Jossang, P.; Hadi, H. A.; Sevenet, T.; Bodo, B. J. Org. Chem. 1991, 56, 6527.
- (6) Riant, O.; Hannedouche, J. Org. Biomol. Chem. 2007, 5, 873.
- (7) Cozzi, P. G.; Hilgraf, R.; Zimmermann, N. Eur. J. Org. Chem. 2007, 5969.
- (8) Bella, M.; Gasperi, T. Synthesis 2009, 1583.
- (9) Quasdorf, K. W.; Overman, L. E. Nature 2014, 516, 181.
- (10) Corey, E. J.; Guzman-Perez, A. Angew. Chem. Int. Ed. **1998**, 37, 388.
- (11) Willis, M. C. J. Chem. Soc., Perkin Trans. 1 1999, 1765.
- (12) Hashimoto, T.; Maruoka, K. Chem. Rev. 2015, 115, 5366.
- (13) Wang, Z.; Liu, J. Beilstein J. Org. Chem. 2020, 16, 3015.

- (14) For reviews on the synthesis and reactivity of VCPs, see:
  (a) Ganesh, V.; Chandrasekharan, S. Synthesis 2016, 48, 4347.
  (b) Brownsey, D. K.; Gorobets, E.; Derksen, D. J. Org. Biomol. Chem. 2018, 16, 3506.
- (15) Some examples include: (a) Trost, B. M.; Morris, P. J.; Sprague, S. J. J. Am. Chem. Soc. 2012, 134, 17823. (b) Mei, L.-Y.; Wei, Y.; Shi, M.; Xu, Q. Organometallics 2013, 32, 3544. (c) Xie, M.-S.; Wang, Y.; Li, J.-P.; Du, C.; Zhang, Y.-Y.; Hao, E.-J.; Zhang, Y.-M.; Qu, G.-R.; Guo, H.-M. Chem. Commun. 2015, 51, 12451.
- (16) Halskov, K. S.; N?sborg, L.; Tur, F.; Jørgensen, K. A. Org. Lett. 2016, 18, 2220.
- (17) Laugeois, M.; Ponra, S.; Ratovelomanana-Vidal, V.; Michelet, V.; Vitale, M. R. *Chem. Commun.* **2016**, *52*, 5332.
- (18) Wan, Q.; Chen, L.; Li, S.; Kang, Q.; Yuan, Y.; Du, Y. Org. Lett. **2020**, 22, 9539.
- (19) Pirenne, V.; Muriel, B.; Waser, J. Chem. Rev. 2021, 121, 227.
- (20) Xiao, J.-A.; Xia, P. J.; Zhang, X.-Y.; Chen, X.-Q.; Ou, G.-C.; Yang, H. *Chem. Commun.* **2016**, *52*, 2177.
- (21) Verma, K.; Banerjee, P. Adv. Synth. Catal. 2017, 359, 3848.
- (22) Dai, C.; Li, M.; Chen, M.; Luo, N.; Wang, C. J. Chem. Res. 2019, 43, 43.
- (23) Muriel, B.; Gagnebin, A.; Waser, J. Chem. Sci. 2019, 10, 10716.
- (24) Uraguchi, D.; Kimura, Y.; Ueoka, F.; Ooi, T. J. Am. Chem. Soc. 2020, 142, 19462.
- (25) Pirenne, V.; Robert, E. G. L.; Waser, J. Chem. Sci. 2021, 12, 8706.

		THIEME	
SynOpen A	A. Deepthi et al.	<b>OPEN</b> ACCESS	Review

- (26) Yin, Y.; Li, Y.; Gonçalves, T. P.; Zhan, Q.; Wang, G.; Zhao, X.; Qiao,
   B.; Huang, K.-W.; Jiang, Z. J. Am. Chem. Soc. 2020, 142, 19451.
- (27) (a) Chen, Y.-R.; Ganapuram, M. R.; Hsieh, K.-H.; Chen, K.-H.; Karanam, P.; Vagh, S. S.; Liou, Y.-C.; Lin, W. Chem. Commun. **2018**, 54, 12702. (b) Vagh, S. S.; Karanam, P.; Liao, C.-C.; Liu, T.-H.; Liou, Y.-C.; Edukondalu, A.; Chen, Y.-R.; Lin, W. Adv. Synth. Catal. **2020**, 362, 1679. (c) Khairnar, P. V.; Su, Y.-H.; Edukondalu, A.; Lin, W. J. Org. Chem. **2021**, 86, 12326.
- (28) Yi, Y.; Hua, Y.-Z.; Lu, H.-J.; Liu, L.-T.; Wang, M.-C. Org. Lett. **2020**, 22, 2527.
- (29) Li, Z.; Lu, Y.; Tian, Y. P.; Hao, X.-X.; Liu, X.-W.; Zhou, Y.; Liu, X.-L. *Tetrahedron* **2021**, *98*, 132297.
- (30) Duffy, C.; Roe, W. E.; Harkin, A. H.; McNamee, R.; Knipe, P. C. New J. Chem. 2021, 45, 22034.
- (31) Xu, B.; Zhang, Z.-M.; Liu, B.; Xu, S.; Zhou, L.-J.; Zhang, J. Chem. Commun. **2017**, 53, 8152.
- (32) (a) Xu, B.; Zhang, Z.-M.; Xu, X.; Liu, B.; Xiao, Y.; Zhang, J. ACS Catal. 2017, 7, 210. (b) Liu, B.; Zhang, Z.-M.; Xu, B.; Xu, S.; Wu, H.-H.; Zhang, J. Adv. Synth. Catal. 2018, 360, 2144.
- (33) Xu, S.; Zhang, Z.-M.; Xu, B.; Liu, B.; Liu, Y.; Zhang, J. J. Am. Chem. Soc. **2018**, 140, 2272.
- (34) Xu, B.; Zhang, Z.-M.; Zhou, L.; Zhang, J. Org. Lett. 2018, 20, 2716.
- (35) Boudriga, S.; Haddad, S.; Askri, M.; Soldera, A.; Knorr, M.; Strohmann, C.; Golz, C. *RSC Adv.* **2019**, 9, 11082.
- (36) Huang, Y.; Fang, H.-L.; Huang, Y.-X.; Sun, J.; Yan, C.-G. J. Org. Chem. **2019**, 84, 12437.
- (37) Imagawa, N.; Nagato, Y.; Ohmatsu, K.; Ooi, T. *Bull. Chem. Soc. Jpn.* **2016**, *89*, 649.
- (38) Wang, J.; Zhao, L.; Rong, Q.; Lv, C.; Lu, Y.; Pan, X.; Zhao, L.; Hu, L. Org. Lett. **2020**, *22*, 5833.
- (39) Ming, S.; Qurban, S.; Du, Y.; Su, W. Chem. Eur. J. 2021, 27, 12742.
- (40) Zhang, Y.; Qin, T.; Zi, W. J. Am. Chem. Soc. 2021, 143, 1038.
- (41) Gao, C.; Zhang, T.; Li, X.; Wu, J.-D.; Liu, J. Org. Chem. Front. **2022**, 9, 2121.

- (42) Zhang, C.; Lu, X. J. Org. Chem. 1995, 60, 2906.
- (43) (a) Wei, Y.; Shi, M. Org. Chem. Front. 2017, 4, 1876. (b) Ni, H.; Chan, W.-L.; Lu, Y. Chem. Rev. 2018, 118, 9344. (c) Guo, H.; Fan, Y. C.; Sun, Z.; Wu, Y.; Kwon, O. Chem. Rev. 2018, 118, 10049.
- (44) Li, H.; Lu, Y. Asian J. Org. Chem. 2017, 6, 1130.
- (45) Ni, H.; Yao, W.; Lu, Y. Beilstein J. Org. Chem. 2016, 12, 343.
- (46) Chen, G.-S.; Zhang, J.-W.; Li, Z.-D.; Zhao, Y.-L.; Liu, Y.-L. Org. Chem. Front. 2020, 7, 3399.
- (47) Gao, Y.; Zhang, J.; Shan, W.; Fei, W.; Yao, J.; Yao, W. Org. Lett. 2021, 23, 6377.
- (48) Oga, M.; Takamatsu, Y.; Ogura, A.; Takao, K. J. Org. Chem. **2022**, 87, 8788.
- (49) Ren, H.-X.; Peng, L.; Song, X.-J.; Liao, L.-G.; Zou, Y.; Tian, F.; Wang, L.-X. Org. Biomol. Chem. 2018, 16, 1297.
- (50) Li, S.; Zhang, P.; Li, Y.; Lu, S.; Gong, J.; Yang, Z. Org. Lett. 2017, 19, 4416.
- (51) Lozanova, A. V.; Stepanov, A. V.; Zlokazov, M. V.; Veselovsky, V. V. ARKIVOC 2017, (iii), 217.
- (52) Ramu, G.; Krishna, N. H.; Pawar, G.; Sastry, K. N. V.; Nanubolu, J. B.; Babu, B. N. ACS Omega **2018**, 3, 12349.
- (53) Zhong, J.; He, H.; Gao, S. Org. Chem. Front. 2019, 6, 3781.
- (54) Shi, H.; Michaelides, I. N.; Derses, B.; Jakubee, P.; Nguyen, Q. N. N.; Paton, R. S.; Dixon, D. J. J. Am. Chem. Soc. **2017**, 139, 17755.
- (55) Ji, Y.; Xin, Z.; He, H.; Gao, S. J. Am. Chem. Soc. 2019, 141, 16208.
- (56) Ling, T.; Rivas, F. Tetrahedron 2016, 72, 6729.
- (57) Zeng, X.-P.; Cao, Z.-Y.; Wang, Y.-H.; Zhou, F.; Zhou, J. Chem. Rev. 2016, 116, 7330.
- (58) Feng, J.; Holmes, M.; Krische, M. J. Chem. Rev. 2017, 117, 12564.
- (59) Xu, P.-W.; Yu, J.-S.; Chen, C.; Cao, Z.-Y.; Zhou, F.; Zhou, J. ACS *Catal.* **2019**, *9*, 1820.
- (60) Wang, Z.; Liu, J. Beilstein J. Org. Chem. 2020, 16, 3015.
- (61) Wang, Z. Org. Chem. Front. 2020, 7, 3815.