



Infected Pseudoaneurysm of the Radial Artery

Haruki Mizuta, MD¹ Heishiro Fujikawa, MD¹ Hisashi Motomura, MD, PhD¹

¹ Department of Plastic and Reconstructive Surgery, Graduate School of Medicine, Osaka Metropolitan University, Osaka, Japan

Address for correspondence Heishiro Fujikawa, MD, Department of Plastic and Reconstructive Surgery, Graduate School of Medicine, Osaka Metropolitan University, 1-4-3 Asahi-machi, Abeno-ku, Osaka, 545-8586, Japan (e-mail: fuji.heishiro@gmail.com).

Arch Plast Surg 2023;50:325–328.

The incidence rates of pseudoaneurysms of the radial artery, local infections, and bacteremia as complications of arterial cannulation procedures are relatively low, affecting only 0.048, 0.4, and 0.6% patients, respectively.^{1,2} The current report describes the successful treatment of an infected pseudoaneurysm of the radial artery, an extremely rare complication of arterial cannulation,³ by early surgical intervention.

A 65-year-old male patient was diagnosed with cardiac arrest caused by anaphylactic shock after undergoing contrast-enhanced computed tomography (CT). The patient received intensive care and underwent careful examination for autoimmune polyneuropathy. His previous medical history included renal transplantation for gouty nephropathy.

Twenty-five days later, the patient exhibited progressive erythema at the site of radial artery cannulation and, upon suspecting cellulitis, intravenous administration of cefazolin and vancomycin was initiated. Blood culture examination revealed methicillin-resistant *Staphylococcus aureus*, and antibiotic treatment was commenced. The affected area was seen to be markedly swollen, blistered, and exhibited visibly abnormal pulsations 3 days after commencement of treatment (► Fig. 1).

As the patient had a history of chronic renal failure, a CT scan was performed which revealed an infected pseudoaneurysm (► Fig. 2), necessitating emergency surgery. The preoperative Allen test was found to be negative and, following surgical debridement, the radial artery was ligated and the pseudoaneurysm was excised. The radial artery was seen to be partially ruptured (► Fig. 3). The radial and median nerves were preserved, and negative pressure wound therapy (NPWT) (V.A.C., KCI Co., Tokyo, Japan) was carried out.

We started the administration of NPWT at a pressure of –75 mm Hg in a continuous mode. Four days after NPWT, granulation appeared gradually. However, palmaris longus and flexor carpi radialis tendon remained exposed. We increased the pressure to –100 mm Hg. Using basic fibroblast

growth factor (bFGF, Fiblast spray), we applied the OASIS collagen matrix (Cook Biotech Incorporated, United States) to tendon-exposed areas. The conditions were favorable for enhanced granulation; therefore, we increased the pressure to –125 mm Hg and modified the pressure every 3 days. Three weeks after NPWT, tendon-exposed areas showed complete granulation; thus, we considered this as the time required to complete NPWT.

The patient exhibited complete wound healing after 2 months, and no complications other than a slight sensory disorder of the dorsal thumb was observed (► Fig. 4).

Staphylococcus aureus are the most commonly observed etiological agent in infected pseudoaneurysms of the radial artery.⁴ Risk factors for the development of infections associated with radial artery catheters include prolonged catheterization (> 4 days) and hospitalization,² while *Staphylococcus aureus* infections, persistent bacteremia for at least 48 hours after catheter removal, and antibiotic therapy have been shown to affect prognosis.⁴ Conditions of both our cases are matched.⁵

Infected pseudoaneurysms are clinically characterized by rapidly swelling wine-red tumors with pulsation. Diagnosis can be made based on clinical observation of a lesion that shrinks upon compression and rapidly refills when released, recognition of an aneurysm through CT or ultrasound examination, and detection of bacteria using blood culture examination.⁴

However, not all cases may present with the typical symptoms. Some patients (including that presented in the current case report) do not exhibit pulsation in the early stages of the disease, making it difficult to distinguish the lesion from cellulitis caused by arterial cannulation.

Infected pseudoaneurysms of the radial artery have a high risk of rupture and require immediate arterial ligation. In case of ischemia in the peripheral region caused by ligation of the radial artery, revascularization using a vein graft should be considered. Therefore, an Allen test should also be performed preoperatively in all cases.

received

July 20, 2022

accepted after revision

February 7, 2023

accepted manuscript online

February 21, 2023

DOI <https://doi.org/>

10.1055/a-2039-3563.

eISSN 2234-6171.

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)

Thieme Medical Publishers, Inc., 333 Seventh Avenue, 18th Floor, New York, NY 10001, USA



Fig. 1 Clinical examination observed at the time of initial presentation of the patient at our department showed a dark-red tumor with blisters and abnormal pulsation on the left wrist.

Noninfectious pseudoaneurysms that are small in size (< 3 cm), stable, and asymptomatic often thrombose within 4 weeks, and follow-up can be recommended in these cases.³ However, this is only applicable for noninfectious pseudoaneurysms that occur after catheter removal or trauma, and

those occurring during indwelling of a radial artery catheter require more careful management.

Previous evidence suggests that approximately 40% of the patients with local infections caused by radial artery catheterization went on to develop a pseudoaneurysm,² suggesting that




Fig. 2 Clinical imaging (computed tomography [CT]): CT revealed low-echoic lesion at his left wrist, suggesting an infected pseudoaneurysm (yellow arrowhead).




Fig. 3 Partially ruptured radial artery was observed intraoperatively, and arterial ligation was performed immediately.



Fig. 4 Clinical observation after 6 months showed complete wound healing.

these lesions can be expected to evolve into infected pseudoaneurysms, even in the absence of an infection initially.⁵

Relatively early diagnosis, timely surgical intervention, careful wound management using NPWT, and adequate debridement contributed to postoperative success in our patient.

To the best of our knowledge, there are a limited number of case reports focusing on infected pseudoaneurysms of the radial artery in the field of plastic surgery. The findings of this study suggest that these lesions require careful postoperative wound management and should, therefore, be given due importance by plastic surgeons.

Author's Contributions

Conceptualization: H.M. and H.F. Writing-original draft: H.M. Writing-review and editing: H.F., H.M. Approval of final manuscript: all authors.

Ethical Approval

The study was performed in accordance with the principles of the Declaration of Helsinki. Written informed consent was obtained.

Patient Consent

The patient provided written informed consent for the publication and the use of her images.

Conflict of Interest

None declared.

References

- 1 Falk PS, Scuderi PE, Sherertz RJ, Motsinger SM. Infected radial artery pseudoaneurysms occurring after percutaneous cannulation. *Chest* 1992;101(02):490–495
- 2 Swanson E, Freiberg A, Salter DR. Radial artery infections and aneurysms after catheterization. *J Hand Surg Am* 1990;15(01):166–171
- 3 Wolf S, Mangano DT. Pseudoaneurysm, a late complication of radial-artery catheterization. *Anesthesiology* 1980;52(01):80–81
- 4 Patel KN, Gandhi SP, Sutariya HC. Radial artery pseudoaneurysm: a rare complication after a single arterial puncture for blood-gas analysis. *Indian J Crit Care Med* 2016;20(10):622–626
- 5 Garg K, Howell BW, Saltzberg SS, et al. Open surgical management of complications from indwelling radial artery catheters. *J Vasc Surg* 2013;58(05):1325–1330