## E-Videos

Total stricture of liver transplant biliary anastomosis resolved with peroral digital cholangioscopy and the non-flexible end of a 0.018-inch biliary guidewire



A 64-year-old man underwent orthotopic liver transplantation due to hepatocellular carcinoma on liver cirrhosis diagnosed 2 years earlier. Three months after transplantation surgery, he presented with jaundice, and a cholangio magnetic resonance imaging scan revealed a biliary anastomosis stricture with marked kinking and retrograde dilatation. A multidisciplinary committee decided on an endoscopic approach.

Initially, retrograde biliary cannulation by endoscopic retrograde cholangiopancreatography was attempted, but advancement of different guidewires through the biliary anastomosis was not possible. The use of peroral digital single-operator cholangioscopy helped us to visualize a complex biliary stricture, which explained why it had not been possible to advance any kind of guidewire into the intrahepatic ducts (> Fig. 1). A 0.018-inch guidewire (Novagold; Boston Scientific, Marlborough, Massachusetts, USA) was inverted with the purpose of using the non-flexible end as a fine needle. This stiff end was successfully inserted under cholangioscopy/fluoroscopy quidance through the closed anastomosis, similarly to a needle (> Fig. 2, ▶ Fig. 3). After this maneuver, the guidewire was restored to its standard position, with the flexible end now able to advance through the anastomosis following the needle-like puncture (> Video 1).

Finally, endoscopic maximal stent therapy (MST) was initiated. First, balloon dilation up to 8 mm was performed, followed by successful placement of three plastic stents (Advanix 10 Fr, 8.5 Fr, and 7 Fr × 12 cm; Boston Scientific) as the first session of a 1-year MST program (**> Fig. 4**).

This case of post-liver transplant biliary stricture was resolved using cholangioscopy/fluoroscopy guidance and a guidewire technical variant. The utility of digital single-operator cholangioscopy



**Fig.1** Cholangioscopy images. **a** The post-liver transplant biliary stenosis. **b** The flexible end of a 0.018-inch guidewire pushing against the biliary stricture.



▶ Fig. 2 Fluoroscopy images. a Failed endoscopic retrograde cholangiopancreatography attempt. b Successful use of peroral digital single-operator cholangioscopy with an inverted 0.018-inch guidewire.

for the endoscopic management of biliary strictures after liver transplantation has been published previously [1,2]. However, use of the non-flexible end of a 0.018-inch guidewire as a method of crossing a complete biliary stenosis has not been previously reported. Endoscopy\_UCTN\_Code\_TTT\_1AR\_2AZ

#### Acknowledgment

We thank the CERCA Programme/Generalitat de Catalunya for institutional support.



► Fig. 3 Cholangioscopy image of the non-flexible end of a 0.018-inch guidewire puncturing through the biliary stricture.



► Fig.4 Three biliary plastic stents placed through the biliary stricture, as the first session of a 1-year program of maximal stent therapy.

#### **Competing interests**

J. B. Gornals is consultant for Boston Scientific. M. Puigcerver-Mas, A. Garcia-Sumalla, J. Lopez-Dominguez, and L. Llado Garriga declare that they have no conflict of interest.



**Video 1** Total stricture of liver transplant biliary anastomosis resolved with peroral digital cholangioscopy and the non-flexible end of a 0.018-inch biliary guidewire.

#### The authors

Maria Puigcerver-Mas<sup>1,2,3</sup>, Albert Garcia-Sumalla<sup>1,2,3</sup>, Josefina Lopez-Dominguez<sup>2,3,4</sup>, Laura Llado Garriga<sup>2,3,4</sup>, Joan B. Gornals<sup>1,2,3</sup>

- 1 Endoscopy Unit, Department of Digestive Diseases, Hospital Universitari de Bellvitge, Barcelona, Catalonia, Spain
- 2 Bellvitge Biomedical Research Institute (IDIBELL), Barcelona, Catalonia, Spain
- 3 Universitat de Barcelona, Barcelona, Catalonia, Spain
- 4 Liver transplant Unit, Department of General Digestive Surgery, Hospital Universitari de Bellvitge, Barcelona, Catalonia, Spain

## Corresponding author

#### Joan B. Gornals, MD, PhD

Endoscopy Unit, Department of Digestive Diseases, Hospital Universitari de Bellvitge, Bellvitge Biomedical Research Institute (IDIBELL), Universitat de Barcelona, Feixa Llarga s/n, 08907 L'Hospitalet de Llobregat, Barcelona, Catalonia, Spain jgornals@bellvitgehospital.cat

#### References

- Bokemeyer A, Lenze F, Stoica V et al. Utility of digital single-operator video cholangioscopy for the endoscopic management of biliary complications after liver transplantation – a single-center experience. Transplantation 2023; 107: 925–932
- [2] Bokemeyer A, Gross D, Brückner M et al. Digital single-operator cholangioscopy: a useful tool for selective guidewire placements across complex biliary strictures. Surg Endosc 2019; 33: 731–737

# Bibliography

Endoscopy 2023; 55: E781–E782 DOI 10.1055/a-2092-0246 ISSN 0013-726X © 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

