

# Exploring Growth, Maturity, and Age as Injury Risk Factors in High-Level Youth Football



## Authors

Eirik Halvorsen Wik<sup>1, 2, 3, 4</sup> , Karim Chamari<sup>1</sup>, Montassar Tabben<sup>1</sup> , Valter Di Salvo<sup>5, 6</sup>, Warren Gregson<sup>5, 7</sup>, Roald Bahr<sup>1, 2</sup>

## Affiliations

- 1 FIFA Medical Centre of Excellence, Aspetar Orthopaedic and Sports Medicine Hospital, Doha, Qatar
- 2 Oslo Sports Trauma Research Center, Department of Sports Medicine, Norwegian School of Sport Sciences, Oslo, Norway
- 3 Institute of Sport and Exercise Medicine, Department of Exercise, Sport and Lifestyle Medicine, Faculty of Medicine and Health Sciences, Stellenbosch University, Stellenbosch, South Africa
- 4 Division of Physiotherapy, Department of Health and Rehabilitation Sciences, Faculty of Medicine and Health Sciences, Stellenbosch University, Stellenbosch, South Africa
- 5 Football Performance and Science Department, Aspire Academy, Doha, Qatar
- 6 Department of Movement, Human and Health Sciences, University of Rome 'Foro Italico', Roma, Italy
- 7 Department of Sport and Exercise Sciences, Institute of Sport, Manchester Metropolitan University, Manchester, United Kingdom of Great Britain and Northern Ireland

## Key words

soccer, growth and development, epidemiology, sports medicine

received 08.03.2023

revised 02.09.2023

accepted 18.09.2023

accepted manuscript online 11.10.2023

published online 2024

## Bibliography

Sports Medicine International Open 2024; 8: a21804594

DOI 10.1055/a-2180-4594

ISSN 2367-1890

© 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purpose, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag, Rüdigerstraße 14,  
70469 Stuttgart, Germany

## Correspondence

Dr. Eirik Halvorsen Wik  
Stellenbosch University  
Faculty of Medicine and Health Sciences  
Francie van Zijl Drive  
7505 Stellenbosch  
South Africa  
Tel.: + 27 761426814  
[eirik.h.wik@gmail.com](mailto:eirik.h.wik@gmail.com)

## ABSTRACT

Rapid somatic growth and biological maturity status may affect injury patterns in youth football, yet firm conclusions cannot be drawn from the existing research. We aimed to explore growth velocity, maturity, and age as injury risk factors in 95 academy players (11.9–15.0 years), using anthropometric (height and body mass), maturity (skeletal age), injury, and football exposure data collected prospectively over three seasons (2016/17–2018/19). We compared the relative quality of mixed-effects logistic regression models with growth velocity for 223 growth intervals (average 113 days) included as fixed effects and adjusted for age (chronological or skeletal) plus load (hours/week). Associations were considered practically relevant based on the confidence interval for odds ratios, using thresholds of 0.90 and 1.11 to define small beneficial and harmful effects, respectively. We observed harmful effects of older age on overall (OR: 2.61, 95% CI: 1.15–5.91) and sudden onset (1.98, 1.17–3.37) injury risk. Significant associations ( $p < 0.05$ ) were observed for higher body mass change and greater maturity on sudden onset injuries, and for higher hours/week on gradual onset, bone tissue, and physis injuries. Future studies should include larger samples, monitoring athletes from pre-adolescence through maturation, to enable within-subject analyses and better understand the relationship between growth, maturation, and injuries.

## Introduction

Elite football (soccer) academies guide players through structured and intensive training programs to optimize long-term development. Injuries impact these opportunities and identification of risk factors is an important step towards reducing injury occurrence and severity by informing targeted injury risk reduction strategies [1]. In youth football, emerging evidence suggests an association between growth, maturation, and injury risk [2, 3]. However, further research is needed to better understand these relationships and how the impact of injuries may be reduced during periods of rapid changes to an athlete's body.

Growth represents an increase in the size of the body as a whole or of a specific body part, assessed using anthropometric measures [4]. An accelerated period of somatic growth is observed during the adolescent years, with peak height velocity (PHV) and peak weight velocity (PWV) for an average boy occurring around the age of 13 to 14 years and 14 to 15 years, respectively [4]. There is, however, a wide range in both timing (age at PHV from 12 to 17 years) and intensity (PHV from 5 to 12 cm/year) [5, 6]. Studies in high-level football have indicated that phases characterized by rapid growth are associated with greater overall, overuse, acute, and non-contact injury risk [7–14], and that players with faster PHV have a greater overall and growth-related injury burden compared to players with average or slower peak growth rates [15]. Still, it is difficult to provide clear recommendations as the associations are inconsistent and methodological differences make direct comparisons or pooling of results impossible.

Maturation is defined by Malina et al. [4] as the process of becoming mature or progressing towards a mature (adult) state (e. g., a fully ossified skeleton, adult height, or a functional reproductive system). Timing and tempo vary greatly between individuals (onset of puberty from 9 to 14 years in boys) [4], and the maturity status (where an individual is in the process at a given point) of two players who train and compete in the same age group can therefore differ substantially [4]. Only a few studies have related skeletal maturity – considered the single best marker of biological maturity [16] – to injury risk in high-level youth football [17–20]. While suggesting that maturity plays a role in the occurrence of certain injury types (e. g., apophyseal or osteochondral injuries, muscle injuries, joint/ligament injuries) [18–20], results are not consistent. Observed age-related injury patterns do, however, indicate that older players are at greater risk and that different pathology types are more prominent in younger, compared to older, age groups [21–23].

Since firm conclusions cannot be drawn from the current pool of literature, the aim of this study was to explore growth velocity, skeletal maturity, and chronological age as injury risk factors in male academy players. More specifically, our main research question was whether changes in height or body mass between assessments at the start, middle, and end of a football season were related to the occurrence of specific injury types when also taking age (chronological or skeletal), growth  $\times$  age/maturity interaction effects, and football exposure into consideration. As an exploratory study, no a priori hypothesis was stated.

## Materials and Methods

### Study design and participants

We used injury, individual training and match exposure, anthropometric and maturity data collected prospectively over three seasons (2016/17 through 2018/19) in the U13, U14, and U15 age groups at one elite national football academy in Qatar (2016/17: 64 players in the program, 2017/18: 77 players, 2018/19: 91 players). Participants were boys aged 11 to 15 years and full-time players typically participated in eight sessions during the school week, in addition to local club games on weekends. Part-time players participated in five sessions in addition to weekend club games. Written informed consent to use routinely collected monitoring data for research purposes was obtained from the players' guardians and ethics approval was granted from the Anti-Doping Lab Qatar Institutional Review Board (IRB Application #2014000012).

### Recording of injuries and football exposure

Training and match injuries were recorded by the designated team physiotherapist (i. e., one physiotherapist per team) who was present at all team sessions, supervised by two researchers. Recording procedures followed the recommendations from Fuller et al. [24], including only time-loss injuries (to reduce bias associated with using several clinical recorders over multiple seasons [25]), i. e., any physical complaint leading to the medical staff partially or fully restricting participation in future football activities. Diagnoses were confirmed by a sports medicine physician (employed full-time at the academy) in collaboration with the treating physiotherapist and reported based on the Sports Medicine Diagnostic Coding System (SMDCS) [26], alongside details about the date of injury and mechanism. No inter-rater reliability data is available regarding the diagnosis or classification of injuries during the observation period. Each team's designated sports scientist recorded individual training and match exposure.

Following the completion of the data collection, a researcher converted injury diagnoses to the updated SMDCS categories for tissue and pathology types [27, 28], and retrospectively allocated onset based on the reported mechanism and diagnosis following the consensus recommended definitions by Bahr et al. ("sudden": resulting from a specific identifiable event, or "gradual": lack of definable sudden, precipitating event) [28]. Only index injuries were considered for this study while recurrent injuries were excluded; these were defined as a time-loss injury to the same location of the same type as a previous injury recorded during the observation period [28].

### Anthropometric and maturity assessments

Measures of standing height and body mass were obtained at the start, middle, and end of each academy season by trained sports scientists following the recommendations outlined by Stewart et al. [29]. Measures were taken in the morning, prior to any activities to minimize diurnal variations. Standing height was measured to the nearest 0.1 cm applying the stretch stature method using a wall-mounted stadiometer (Holtain Ltd, Crymch, UK) and body mass was measured to the nearest 0.1 kg using digital scales (Adam Equipment, Milton Keynes, UK). Previously published test-retest data in a subsample of 17 academy players revealed a standard error of measurement (SEM) of 0.34 cm (95% confidence interval

(CI): 0.25 to 0.52 cm) for standing height [30]. This corresponds to a minimal detectable change (MDC) of approximately 1 cm.

Skeletal maturity was assessed at the beginning of the season using x-ray images of the player's left hand/wrist complex taken at Aspetar Orthopaedic and Sports Medicine Hospital. Skeletal age was determined using the Fels method [31] by one trained assessor. Intra-rater reliability for this method has previously been reported (intraclass correlation coefficient (ICC): 0.998, 95% CI: 0.996 to 0.999) [17].

## Inclusion and exclusion of growth intervals

It has been suggested that researchers examine growth over shorter periods of time to better account for non-linear growth patterns related to saltatory (episodic) growth [32]. To minimize the impact of measurement error on estimations and allow for detection of meaningful changes [32, 33], we calculated growth velocity per academy semester, defined as the two intervals from season start (August/September) to mid-season (January) and from mid-season to season end (May/June). The absolute change (cm or kg) was divided by the number of days between measurements and converted to expressions equivalent to cm/year and kg/year, respectively [12, 13]. For a growth interval to be included in the final analyses, a skeletal maturity assessment had to be available for the given player and season (assessed maximum 91 days within the start of the season).

## Statistical analyses

Descriptive statistics are presented as means with standard deviation (SD). Four separate mixed-effects logistic regression models (*xtnlogit* command) estimated associations for the effects of changes in height and body mass on the occurrence of overall, gradual onset, sudden onset, bone tissue and physis injury. Growth velocity for height and body mass were specified as distinct growth-related predictor variables (fixed effects). Models were adjusted for


chronological or skeletal age, and growth × age/maturity interaction, with player specified as a random effect plus a random intercept. The average weekly load (hours of training/match exposure per week) during the growth interval (or until the event if an injury occurred) was added as a covariate.

The Akaike Information Criterion (AIC) assessed the relative quality of each mixed-effects logistic regression model in the set of candidate models. The Akaike difference ( $\Delta$ AIC) from the estimated best model (i. e., the model with the lowest AIC value;  $\Delta$ AIC = 0) was evaluated according to the following scale: 0–2, essentially equivalent; > 2–7, plausible alternative; > 7–14, weak support; > 14, no empirical support [34]. Akaike weights ( $w_i$ ) provided a scaled interpretation about the relative quality of each competing model as the probability that a given model is the best in the set of four candidate models per endpoint. Thresholds for the adjusted odds ratios (OR) of 0.9, 0.7, 0.5, 0.3, and 0.1 and their reciprocals 1.11, 1.43, 2.0, 3.3, and 10 defined small, moderate, large, very large, and extremely large beneficial and harmful effects, respectively [35]. In the absence of an established anchor defining practically relevant associations between growth velocity and injury occurrence, we considered OR = 0.90 or OR = 1.11 to define substantially beneficial and substantially harmful effects, respectively [35]. Associations were declared practically relevant based on the location of the confidence interval for the estimated ORs to these thresholds. Outcome statistics are reported as point estimates and 95% CI. Statistical analyses were performed using Stata (StataBE v17.0; StataCorp, College Station, TX, USA).

## Results

### Inclusion of players and growth intervals

The inclusion of player-seasons and growth intervals are shown in ► **Fig. 1**, with an overview of exclusions due to incomplete assess-



► **Fig. 1** Inclusion of player seasons and growth intervals, with an overview of exclusions due to missing data. Player season: One player taking part in one season. Eligible player season: Player season with minimum two anthropometric measures. Eligible growth interval: An interval spanning from either the start to the middle of the season or the middle to the end of the season, with complete anthropometric assessments on both sides. Included growth intervals: An interval (Start-Mid or Mid-End) with a skeletal age assessment for the given season for a player.

► **Table 1** Overview of the 161 index injuries sustained within the growth periods included in the analyses, structured by location, onset, and pathology type.

| Location<br>Onset<br>Pathology type | Injuries<br>(count) | Incidence<br>(Inj. per 1000 h, 95% CI) |                       |
|-------------------------------------|---------------------|--|-----------------------|
| <b>Head and neck</b>                | <b>6</b>            | <b>0.28</b>                            | <b>(0.10 to 0.60)</b> |
| Sudden onset                        | 6                   | 0.28                                   | (0.10 to 0.60)        |
| Brain/spinal cord                   | 5                   | 0.23                                   | (0.07 to 0.54)        |
| Laceration                          | 1                   | 0.05                                   | (0.00 to 0.26)        |
| <b>Upper limb</b>                   | <b>18</b>           | <b>0.83</b>                            | <b>(0.49 to 1.31)</b> |
| Sudden onset                        | 18                  | 0.83                                   | (0.49 to 1.31)        |
| Fracture                            | 13                  | 0.60                                   | (0.32 to 1.02)        |
| Contusion (superficial)             | 4                   | 0.18                                   | (0.05 to 0.47)        |
| Joint sprain                        | 1                   | 0.05                                   | (0.00 to 0.26)        |
| <b>Trunk</b>                        | <b>7</b>            | <b>0.32</b>                            | <b>(0.13 to 0.66)</b> |
| Sudden onset                        | 4                   | 0.18                                   | (0.05 to 0.47)        |
| Contusion (superficial)             | 2                   | 0.09                                   | (0.01 to 0.33)        |
| Fracture                            | 1                   | 0.05                                   | (0.00 to 0.26)        |
| Non-specific                        | 1                   | 0.05                                   | (0.00 to 0.26)        |
| Gradual onset                       | 3                   | 0.14                                   | (0.03 to 0.40)        |
| Bone stress injury                  | 2                   | 0.09                                   | (0.01 to 0.33)        |
| Non-specific                        | 1                   | 0.05                                   | (0.00 to 0.26)        |
| <b>Hip/groin</b>                    | <b>26</b>           | <b>1.20</b>                            | <b>(0.78 to 1.75)</b> |
| Sudden onset                        | 4                   | 0.18                                   | (0.05 to 0.47)        |
| Contusion (superficial)             | 2                   | 0.09                                   | (0.01 to 0.33)        |
| Muscle injury                       | 1                   | 0.05                                   | (0.00 to 0.26)        |
| Non-specific                        | 1                   | 0.05                                   | (0.00 to 0.26)        |
| Gradual onset                       | 22                  | 1.01                                   | (0.64 to 1.53)        |
| Physis injury                       | 20                  | 0.92                                   | (0.56 to 1.42)        |
| Bone stress injury                  | 1                   | 0.05                                   | (0.00 to 0.26)        |
| Bursitis                            | 1                   | 0.05                                   | (0.00 to 0.26)        |
| <b>Thigh</b>                        | <b>35</b>           | <b>1.61</b>                            | <b>(1.12 to 2.24)</b> |
| Sudden onset                        | 31                  | 1.43                                   | (0.97 to 2.03)        |
| Muscle injury                       | 20                  | 0.92                                   | (0.56 to 1.42)        |
| Muscle contusion                    | 5                   | 0.23                                   | (0.07 to 0.54)        |
| Non-specific                        | 5                   | 0.23                                   | (0.07 to 0.54)        |
| Cartilage                           | 1                   | 0.05                                   | (0.00 to 0.26)        |
| Gradual onset                       | 4                   | 0.18                                   | (0.05 to 0.47)        |
| Physis injury                       | 3                   | 0.14                                   | (0.03 to 0.40)        |
| Non-specific                        | 1                   | 0.05                                   | (0.00 to 0.26)        |
| <b>Knee</b>                         | <b>19</b>           | <b>0.88</b>                            | <b>(0.53 to 1.37)</b> |
| Sudden onset                        | 11                  | 0.51                                   | (0.25 to 0.91)        |
| Contusion (superficial)             | 7                   | 0.32                                   | (0.13 to 0.66)        |
| Joint sprain                        | 2                   | 0.09                                   | (0.01 to 0.33)        |
| Fracture                            | 1                   | 0.05                                   | (0.00 to 0.26)        |
| Non-specific                        | 1                   | 0.05                                   | (0.00 to 0.26)        |
| Gradual onset                       | 8                   | 0.37                                   | (0.16 to 0.73)        |
| Physis injury                       | 5                   | 0.23                                   | (0.07 to 0.54)        |
| Synovitis/capsulitis                | 2                   | 0.09                                   | (0.01 to 0.33)        |
| Cartilage                           | 1                   | 0.05                                   | (0.00 to 0.26)        |
| <b>Lower leg</b>                    | <b>12</b>           | <b>0.55</b>                            | <b>(0.29 to 0.97)</b> |
| Sudden onset                        | 11                  | 0.51                                   | (0.25 to 0.91)        |
| Muscle injury                       | 5                   | 0.23                                   | (0.07 to 0.54)        |

► **Table 1** Continued.

| Location<br>Onset<br>Pathology type | Injuries<br>(count) | Incidence<br>(Inj. per 1000 h, 95% CI) |                       |
|-------------------------------------|---------------------|--|-----------------------|
| Contusion (superficial)             | 3                   | 0.14                                   | (0.03 to 0.40)        |
| Non-specific                        | 2                   | 0.09                                   | (0.01 to 0.33)        |
| Fracture                            | 1                   | 0.05                                   | (0.00 to 0.26)        |
| Gradual onset                       | 1                   | 0.05                                   | (0.00 to 0.26)        |
| Bone stress injury                  | 1                   | 0.05                                   | (0.00 to 0.26)        |
| <b>Ankle</b>                        | <b>25</b>           | <b>1.15</b>                            | <b>(0.75 to 1.70)</b> |
| Sudden onset                        | 20                  | 0.92                                   | (0.56 to 1.42)        |
| Joint sprain                        | 11                  | 0.51                                   | (0.25 to 0.91)        |
| Contusion (superficial)             | 9                   | 0.41                                   | (0.19 to 0.79)        |
| Gradual onset                       | 5                   | 0.23                                   | (0.07 to 0.54)        |
| Synovitis / capsulitis              | 5                   | 0.23                                   | (0.07 to 0.54)        |
| <b>Foot</b>                         | <b>13</b>           | <b>0.60</b>                            | <b>(0.32 to 1.02)</b> |
| Sudden onset                        | 9                   | 0.41                                   | (0.19 to 0.79)        |
| Contusion (superficial)             | 5                   | 0.23                                   | (0.07 to 0.54)        |
| Fracture                            | 2                   | 0.09                                   | (0.01 to 0.33)        |
| Joint sprain                        | 2                   | 0.09                                   | (0.01 to 0.33)        |
| Gradual onset                       | 4                   | 0.18                                   | (0.05 to 0.47)        |
| Physis injury                       | 2                   | 0.09                                   | (0.01 to 0.33)        |
| Non-specific                        | 2                   | 0.09                                   | (0.01 to 0.33)        |

ments. The final sample included 95 unique players contributing to 223 growth intervals (17 players with one growth interval, 48 with two, 14 with three, 12 with four, and 4 with five intervals), with a mean duration of 113 days (SD 24). A total of 161 index injuries (93 from training sessions, 68 from matches) and 21 712 exposure hours (18 642 training hours and 3070 match hours) were recorded within the growth intervals included (overall incidence: 7.4 injuries per 1000 h, training incidence: 5.0 per 1000 h; match incidence: 22.1 per 1000 h). The most common injury locations were the thigh (22%), hip/groin (16%), and ankle (16%), while the most common pathology types were superficial contusions (20%), physis injuries (19%), and muscle injuries (16%). A detailed injury overview is included in ► **Table 1**.

### Age, skeletal maturity, and growth velocity

The mean age at the start of a growth interval was 13.5 years (SD 0.8; range 11.9 to 15.0). Considering each player-season only once (a player could have two growth intervals per season but only one maturity assessment), the mean skeletal age at the start of the season was 14.4 years (SD 1.6) with skeletal ages ranging from 10.7 to 14.9 years in the U13 age group, 11.8 to 18.0 in U14, and 13.7 to 17.8 in U15. On average, players were 1.0 year (1.1; -1.5 to 4.7) advanced in skeletal age relative to chronological age. One player was skeletally mature (skeletal age 18 years), while 62 (45%) could be considered early maturing (skeletal age minimum one year in advance of chronological age), 70 (51%) as on time (skeletal age within one year) and four (3%) as late maturing (skeletal age minimum one year delayed) [36]. The mean semester growth velocity

► **Table 2** Relative model quality for each injury category.

| Model                           | AIC   | Δ AIC | $w_i$ | Inference              |
|---------------------------------|-------|-------|-------|------------------------|
| Overall (119 events)            |       |       |       |                        |
| Δ Body mass & skeletal age      | 306.8 | 0.0   | 0.59  | Best                   |
| Δ Height & skeletal age         | 308.0 | 1.2   | 0.33  | Essentially equivalent |
| Δ Height & chronological age    | 311.6 | 4.8   | 0.05  | Plausible alternative  |
| Δ Body mass & chronological age | 312.9 | 6.1   | 0.03  | Plausible alternative  |
| Sudden onset (90 events)        |       |       |       |                        |
| Δ Height & skeletal age         | 300.3 | 0.0   | 0.52  | Best                   |
| Δ Body mass & skeletal age      | 300.8 | 0.5   | 0.41  | Essentially equivalent |
| Δ Body mass & chronological age | 305.2 | 4.8   | 0.05  | Plausible alternative  |
| Δ Height & chronological age    | 306.5 | 6.1   | 0.02  | Plausible alternative  |
| Gradual onset (42 events)       |       |       |       |                        |
| Δ Body mass & skeletal age      | 216.4 | 0.0   | 0.43  | Best                   |
| Δ Height & skeletal age         | 216.7 | 0.3   | 0.37  | Essentially equivalent |
| Δ Height & chronological age    | 219.3 | 2.9   | 0.10  | Plausible alternative  |
| Δ Body mass & chronological age | 219.4 | 3.0   | 0.10  | Plausible alternative  |
| Bone tissue (49 events)         |       |       |       |                        |
| Δ Height & skeletal age         | 238.1 | 0.0   | 0.51  | Best                   |
| Δ Height & chronological age    | 239.7 | 1.7   | 0.22  | Essentially equivalent |
| Δ Body mass & skeletal age      | 240.4 | 2.3   | 0.16  | Plausible alternative  |
| Δ Body mass & chronological age | 241.1 | 3.0   | 0.11  | Plausible alternative  |
| Physis injury (27 events)       |       |       |       |                        |
| Δ Height & skeletal age         | 166.2 | 0.0   | 0.51  | Best                   |
| Δ Height & chronological age    | 168.2 | 2.0   | 0.19  | Plausible alternative  |
| Δ Body mass & skeletal age      | 168.6 | 2.4   | 0.15  | Plausible alternative  |
| Δ Body mass & chronological age | 168.7 | 2.5   | 0.15  | Plausible alternative  |

AIC, Akaike Information Criteria; Δ AIC, Akaike difference;  $w_i$ , Akaike weights.

was 6.3 cm/year (3.5; 0.0 to 17.8) for height and 5.3 kg/year (5.5; -14.6 to 19.3) for body mass.

### Relative model quality

The relative model quality of the four model combinations within the five injury categories is presented in ► **Table 2**. Growth velocity for body mass combined with skeletal age best explained the overall and gradual onset injury risk, while change in height combined with skeletal age best explained the risk of sudden onset, bone tissue, and physis injuries. Other model combinations were, however, considered equivalent or plausible alternatives.

### Effects of growth velocity, age, and skeletal maturity

► **Table 3** gives a complete overview of all model outcomes. Effects for load and growth × age/maturity interaction were not practically relevant for any injury categories. Practically relevant harmful effects of older age were observed for overall and sudden onset injury risk in the models adjusting for changes in height and body mass, respectively. Significant associations ( $p < 0.05$ ) between greater change in body mass (in the model with chronological age) and more advanced maturity (in the model with body mass change) were seen for sudden onset injury risk. These results were not practically relevant given our predefined thresholds (95% CI for OR  $< 0.9$  or  $> 1.1$ ). Significant, but not practically relevant, associations were also found between higher football load and risk of gradual onset

(all model combinations), bone tissue (all model combinations), and physis injuries (only for the model including body mass change and chronological age).

### Discussion

This study explored growth velocity and age (skeletal and chronological) as injury risk factors, accounting for growth × age/maturity interaction effects and individual training and match exposure. Based on prospective data from 95 unique players between 11 and 15 years over three seasons, we observed harmful effects of older age on overall and sudden onset injury risk. Significant associations were also found for greater change in body mass and more advanced maturity on sudden onset injury risk, and for greater football load on gradual onset, bone tissue, and physis injuries; however, these were not considered practically relevant based on our pre-defined thresholds. No significant growth × age/maturity interaction effects were seen.

### Associations between growth velocity and injury risk remain unclear

A potential link between growth velocity and injury risk is typically attributed to underlying mechanisms such as tissues adapting at different rates, increased tension on apophyses, or decreased neuromuscular control [3, 37–39]. In support of such a link, associa-

► **Table 3** Odds ratios for the four model combinations within each of the five injury outcomes. Numbers in italics indicate significant associations ( $p < 0.05$ ), while asterisks indicate practically relevant findings based on our predefined thresholds (95% CI for OR  $< 0.9$  or  $> 1.1$ ).

| Outcome                      | Model   | Odds ratio (95% CI)  | p-value |
|------------------------------|---|----------------------|---------|
| Overall<br>(119 events)      | $\Delta$ Height (cm/year)                     | 3.78 (0.79 to 18.03) | 0.10    |
|                              | Chronological age                             | 2.61 (1.15 to 5.91)  | 0.022*  |
|                              | $\Delta$ Height $\times$ chronological age    | 0.91 (0.81 to 1.02)  | 0.10    |
|                              | Hours per week                                | 1.06 (0.94 to 1.21)  | 0.35    |
|                              | $\Delta$ Height (cm/year)                     | 1.43 (0.59 to 3.43)  | 0.43    |
|                              | Skeletal age                                  | 1.40 (0.97 to 2.03)  | 0.07    |
|                              | $\Delta$ Height $\times$ skeletal age         | 0.98 (0.92 to 1.04)  | 0.44    |
|                              | Hours per week                                | 1.06 (0.93 to 1.20)  | 0.38    |
|                              | $\Delta$ Body mass (kg/year)                  | 1.82 (0.67 to 4.92)  | 0.24    |
|                              | Chronological age                             | 1.65 (0.99 to 2.73)  | 0.05    |
|                              | $\Delta$ Body mass $\times$ chronological age | 0.96 (0.89 to 1.03)  | 0.29    |
|                              | Hours per week                                | 1.08 (0.95 to 1.22)  | 0.26    |
|                              | $\Delta$ Body mass (kg/year)                  | 0.92 (0.54 to 1.58)  | 0.77    |
|                              | Skeletal age                                  | 1.21 (0.97 to 1.51)  | 0.09    |
|                              | $\Delta$ Body mass $\times$ skeletal age      | 1.01 (0.97 to 1.05)  | 0.59    |
|                              | Hours per week                                | 1.06 (0.94 to 1.20)  | 0.36    |
| Sudden onset<br>(90 events)  | $\Delta$ Height (cm/year)                     | 2.26 (0.53 to 9.58)  | 0.27    |
|                              | Chronological age                             | 1.95 (0.93 to 4.10)  | 0.08    |
|                              | $\Delta$ Height $\times$ chronological age    | 0.94 (0.85 to 1.05)  | 0.26    |
|                              | Hours per week                                | 1.06 (0.94 to 1.21)  | 0.33    |
|                              | $\Delta$ Height (cm/year)                     | 1.36 (0.59 to 3.13)  | 0.47    |
|                              | Skeletal age                                  | 1.38 (0.98 to 1.94)  | 0.07    |
|                              | $\Delta$ Height $\times$ skeletal age         | 0.98 (0.92 to 1.04)  | 0.45    |
|                              | Hours per week                                | 1.06 (0.93 to 1.20)  | 0.40    |
|                              | $\Delta$ Body mass (kg/year)                  | 2.81 (1.02 to 7.79)  | 0.046   |
|                              | Chronological age                             | 1.98 (1.17 to 3.37)  | 0.011*  |
|                              | $\Delta$ Body mass $\times$ chronological age | 0.93 (0.86 to 1.00)  | 0.06    |
|                              | Hours per week                                | 1.07 (0.94 to 1.22)  | 0.29    |
|                              | $\Delta$ Body mass (kg/year)                  | 1.18 (0.69 to 2.02)  | 0.54    |
|                              | Skeletal age                                  | 1.30 (1.04 to 1.63)  | 0.021   |
|                              | $\Delta$ Body mass $\times$ skeletal age      | 0.99 (0.96 to 1.03)  | 0.69    |
|                              | Hours per week                                | 1.06 (0.94 to 1.21)  | 0.35    |
| Gradual onset<br>(42 events) | $\Delta$ Height (cm/year)                     | 1.93 (0.26 to 14.11) | 0.52    |
|                              | Chronological age                             | 1.38 (0.48 to 4.00)  | 0.55    |
|                              | $\Delta$ Height $\times$ chronological age    | 0.96 (0.83 to 1.11)  | 0.56    |
|                              | Hours per week                                | 1.22 (1.01 to 1.48)  | 0.035   |
|                              | $\Delta$ Height (cm/year)                     | 1.35 (0.44 to 4.16)  | 0.60    |
|                              | Skeletal age                                  | 1.13 (0.70 to 1.84)  | 0.61    |
|                              | $\Delta$ Height $\times$ skeletal age         | 0.98 (0.91 to 1.06)  | 0.69    |
|                              | Hours per week                                | 1.23 (1.01 to 1.48)  | 0.035   |
|                              | $\Delta$ Body mass (kg/year)                  | 1.13 (0.30 to 4.24)  | 0.85    |
|                              | Chronological age                             | 0.91 (0.44 to 1.88)  | 0.79    |

► **Table 3** Continued.

| Outcome                         | Model                           | Odds ratio (95% CI)  | p-value             |
|---------------------------------|---------------------------------|----------------------|---------------------|
|                                 | Δ Body mass × chronological age | 1.00 (0.90 to 1.10)  | 0.94                |
|                                 | Hours per week                  | 1.24 (1.02 to 1.50)  | 0.027               |
|                                 | Δ Body mass (kg/year)           | 0.79 (0.34 to 1.81)  | 0.57                |
|                                 | Skeletal age                    | 0.89 (0.62 to 1.29)  | 0.55                |
|                                 | Δ Body mass × skeletal age      | 1.02 (0.96 to 1.08)  | 0.46                |
|                                 | Hours per week                  | 1.23 (1.02 to 1.49)  | 0.031               |
|                                 | Bone tissue<br>(49 events)      | Δ Height (cm/year)   | 0.72 (0.13 to 4.05) |
| Chronological age               |                                 | 0.63 (0.25 to 1.57)  | 0.32                |
| Δ Height × chronological age    |                                 | 1.03 (0.90 to 1.17)  | 0.69                |
| Hours per week                  |                                 | 1.17 (1.00 to 1.36)  | 0.048               |
| Δ Height (cm/year)              |                                 | 1.17 (0.45 to 3.01)  | 0.75                |
| Skeletal age                    |                                 | 0.96 (0.64 to 1.43)  | 0.83                |
| Δ Height × skeletal age         |                                 | 0.99 (0.93 to 1.06)  | 0.80                |
| Hours per week                  |                                 | 1.17 (1.01 to 1.37)  | 0.042               |
| Δ Body mass (kg/year)           |                                 | 1.50 (0.49 to 4.57)  | 0.48                |
| Chronological age               |                                 | 0.81 (0.45 to 1.46)  | 0.48                |
| Δ Body mass × chronological age |                                 | 0.97 (0.90 to 1.06)  | 0.52                |
| Hours per week                  |                                 | 1.18 (1.01 to 1.37)  | 0.040               |
| Δ Body mass (kg/year)           |                                 | 1.06 (0.56 to 2.01)  | 0.86                |
| Skeletal age                    |                                 | 0.89 (0.68 to 1.17)  | 0.41                |
| Δ Body mass × skeletal age      |                                 | 1.00 (0.95 to 1.04)  | 0.94                |
| Hours per week                  | 1.18 (1.01 to 1.37)             | 0.040                |                     |
| Physis injury<br>(27 events)    | Δ Height (cm/year)              | 1.76 (0.22 to 14.37) | 0.60                |
|                                 | Chronological age               | 0.95 (0.29 to 3.07)  | 0.93                |
|                                 | Δ Height × chronological age    | 0.97 (0.83 to 1.13)  | 0.66                |
|                                 | Hours per week                  | 1.20 (0.98 to 1.47)  | 0.08                |
|                                 | Δ Height (cm/year)              | 1.75 (0.55 to 5.59)  | 0.35                |
|                                 | Skeletal age                    | 1.12 (0.66 to 1.89)  | 0.68                |
|                                 | Δ Height × skeletal age         | 0.97 (0.89 to 1.05)  | 0.44                |
|                                 | Hours per week                  | 1.21 (0.98 to 1.48)  | 0.07                |
|                                 | Δ Body mass (kg/year)           | 1.74 (0.40 to 7.65)  | 0.46                |
|                                 | Chronological age               | 0.75 (0.32 to 1.75)  | 0.51                |
|                                 | Δ Body mass × chronological age | 0.97 (0.86 to 1.08)  | 0.53                |
|                                 | Hours per week                  | 1.23 (1.01 to 1.51)  | 0.044               |
|                                 | Δ Body mass (kg/year)           | 0.82 (0.32 to 2.11)  | 0.68                |
|                                 | Skeletal age                    | 0.78 (0.50 to 1.20)  | 0.25                |
|                                 | Δ Body mass × skeletal age      | 1.02 (0.95 to 1.09)  | 0.58                |
| Hours per week                  | 1.23 (1.00 to 1.52)             | 0.05                 |                     |

tions between changes in height and injuries have been reported in Dutch, Belgian, and English high-level football players [11–14]. Although these studies suggest a growth-injury relationship, different analytical approaches and broad injury categories make def-

inite conclusions difficult. To improve our understanding of growth as an injury risk factor, we included bone tissue and physis injuries as specific outcomes, based on the assumption that they are more closely aligned with the suggested underlying mechanisms. Impor-

tantly, these injuries are among the most common and burdensome in U13-U15 age group players [21–23], making them a priority for targeted injury reduction programs. The mean age of our sample (13.5 years) was also close to the expected age at PHV (13.6 years) in this specific academy population [30], which has been highlighted as a vulnerable phase for football players [7–10]. Still, we did not find any practically relevant effects of changes in height or body mass over an academy semester, with only one significant association suggesting an increased risk of sudden onset injuries with greater changes in body mass in the model adjusting for chronological age.

The absence of observed effects in our study may be explained by the methodological approach, using isolated and pre-determined growth intervals (i. e., Start-Mid and Mid-End of a season). The duration of growth intervals reflected recommendations for assessing growth velocity (i. e., every 3 to 4 months [33]), but they do not necessarily capture the periods of most rapid growth within an individual's growth process. They may also not be frequent enough to identify shorter bursts of growth, which could be of interest [32]. Capturing data on a more frequent basis is, however, associated with greater variance in the estimated growth velocities [40], and recommendations to focus on long-term tracking of anthropometric data therefore seem reasonable [33]. This approach was taken by Monasterio et al. [10, 15], who calculated full growth curves. They did, unfortunately, not have individual training and match data available and growth curves could only be calculated for around 10% of their initial sample, highlighting the logistical challenges associated with accurately tracking growth, injuries, and individual exposure over a sufficient duration in applied academy settings.

### Practically relevant effects of age, but not maturity, on injury risk

Studies have indicated a changing injury pattern with age, where physal or "growth-related" injuries are more common in younger age groups [21–23]. This may be attributed, in part, to maturity status [18, 19, 41], with the immature skeleton representing a relatively weak link in the muscle-tendon-bone chain prior to reaching its mature state [42]. Consequently, similar injury mechanisms would result in different pathology types (e. g., apophysitis or avulsions as opposed to tendinopathies or muscle strains) depending on a player's maturity status [42]. These patterns provide a rationale for regular maturity assessments, which can be used to better accommodate for early, on time, and late maturing players within an age group, who may be prone to different injury types at different locations.

In our sample, higher chronological age was related to a practically relevant increased risk of overall and sudden onset injuries, with point estimates suggesting moderate to large harmful effects on these injury outcomes. This is in line with our earlier study, which demonstrated increased injury incidence in older age groups in this academy [21]. Although large variations in skeletal age were observed within age groups (e. g., a six-year range in skeletal age between players in the U14 age group), we did not detect any practically relevant effects of maturity on injury risk. A significant association was found between older skeletal age and sudden onset injury risk; still, we urge caution when interpreting this, as we could not demonstrate consistent associations across models. This may

be due to a relatively small number of events for the number of variables included in our models, and studies including larger samples are needed to better understand these relationships.

### Weekly load may affect the risk of gradual onset and bone tissue injuries

Football load (operationalized as hours per week in this study) was a significant covariate in all model combinations for gradual onset and bone injuries, and in one model for physis injuries. Again, these associations could not be classified as practically relevant, and a causal relationship cannot be established based on our study design. Youth-specific consensus statements have previously highlighted training load as a risk factor for these injury types [43, 44], and studies in high-level youth football have related weekly duration, cumulative absolute training loads over three and four weeks, and greater week-to-week changes in exposure to injury risk [14, 45, 46]. While their findings are inconsistent and our results do not provide definite answers, it does appear sensible to focus on careful progression, diverse and variable movement exposure, careful scheduling of total load, and allowing for sufficient rest and recovery to ensure positive adaptations of training while minimizing the risk of injury in youth settings [3, 43, 44]. Future studies may be able to incorporate more detailed measures of load in their analyses to determine whether training load or athlete responses (e. g., physiological, perceptual) are associated with injury occurrence during specific phases of the growth and maturation process.

### Methodological considerations

We addressed several limitations from previous studies on growth, maturation, and injury risk with our models accounting for growth  $\times$  age/maturity interaction, repeated player-seasons, and daily individual football exposure, including more detailed injury outcomes recorded and verified by on-site medical staff. It is indeed rare that youth football teams have designated medical staff present at each session, and that the detailed injury data they captured can be merged with accurate individual football exposure and skeletal age assessments. Still, there are limitations with our study design that should be considered when interpreting our results.

First, our growth and maturity data were primarily collected for clinical and applied purposes, which led to a substantial number of incomplete assessments (► **Fig. 1**), presumably at random and not related to injury occurrence. Using data collected for ongoing monitoring purposes also introduces the possibility that training content could have been adjusted based on the data collected. However, we do not believe that these assessments impacted significantly on training content during the data collection period, as systematic and contextualized reports were not available to coaches at the time. The exploratory nature of our study, involving a retrospective analysis, also precluded any formal a priori sample size estimation relevant to a mixed-effects modeling framework. Future studies could improve in this area by employing strict data collection routines to avoid missing data and should attempt to include a larger and younger sample of players than we were able to in our study, to cover the entire growth process and allow for within-subject analyses.

Second, including skeletal age as our main indicator of biological maturity is arguably a strength as it is considered a precise and



reliable measure and can be used across the maturation process [4, 16]. However, we recognize its limitations. The assessment requires equipment and expertise (and is therefore costly), and importantly involves low-dose radiation (although this has been described as almost negligible [47]). It is therefore considered an invasive measure that is not feasible to include in most academy settings [3], reducing the direct applicability of our findings to practice. Furthermore, the ossification of the hand-wrist complex does not necessarily represent the maturity status of bones in other locations. Future studies may be able to use radiation-free alternatives, such as magnetic resonance imaging or ultrasound [47], to relate skeletal maturity of multiple body parts to injuries in surrounding tissues.

Finally, although we could expand on findings from previous studies by using injury outcomes that are more closely aligned with the proposed mechanisms for a growth-maturity-injury relationship, the use of a time-loss definition limited our ability to capture injuries that did not affect participation [28]. Future studies could improve in this area by using athlete-reported measures with a broader, any-complaint definition. Readers should also be cognizant that a study like ours may not be representative for other contexts (e. g., different geographical regions, training environments, or match schedules/formats) and addresses but a few of many potential risk factors for injury, which are thought to be both multifactorial, dynamic, and complex [48].

## Conclusion

Our main research question was whether changes in height or body mass between assessments at the start, middle, and end of a football season affect the risk of injury in male academy football players. After accounting for age or skeletal maturity, growth  $\times$  age/maturity interaction effects, and football exposure, no such associations were considered practically relevant based on our pre-defined thresholds. We do, however, report practically relevant harmful effects of older age on overall and sudden-onset injury risk. Additionally, significant – but not practically relevant – associations were observed between greater changes in body mass and older skeletal age on sudden onset injury risk, and higher weekly load on gradual onset, bone tissue, and physis injuries. Researchers should strive to establish robust surveillance systems that can include larger and more diverse samples (athletes of both genders from multiple settings) and capture the whole growth and maturity process (longitudinal tracking starting from pre-adolescence) alongside reliable collection of injury and exposure data (following consensus procedures and universal classification systems), to better understand the relationship between growth, maturation, and injury risk.

## Acknowledgements

The authors would like to acknowledge the efforts made by the staff members at the Aspire Academy Sports Medicine Center and Aspire Academy Football Performance and Science Department who were involved in the collection of growth, maturation, injury and football exposure data. We also thank Morten Wang Fagerland for valuable discussions regarding the study design.

## Conflict of Interest

The authors declare that they have no conflict of interest.

## References

- [1] van Mechelen W, Hlobil H, Kemper HC. Incidence, severity, aetiology and prevention of sports injuries. A review of concepts. *Sports Med* 1992; 14: 82–99. DOI: 10.2165/00007256-199214020-00002
- [2] Wik EH. Growth, maturation and injuries in high-level youth football (soccer): A mini review. *Front Sports Act Living* 2022; 4: 975900. DOI: 10.3389/fspor.2022.975900
- [3] Towson C, Salter J, Ade JD et al. Maturity-associated considerations for training load, injury risk, and physical performance in youth soccer: One size does not fit all. *J Sport Health Sci* 2021; 10: 403–412. DOI: 10.1016/j.jshs.2020.09.003
- [4] Malina RM, Bouchard C, Bar-Or O. Growth, maturation, and physical activity. 2<sup>nd</sup> ed. Champaign, IL: Human Kinetics; 2004
- [5] Largo RH, Gasser T, Prader A et al. Analysis of the adolescent growth spurt using smoothing spline functions. *Ann Hum Biol* 1978; 5: 421–434. DOI: 10.1080/03014467800003071
- [6] Preece MA, Baines MJ. A new family of mathematical models describing the human growth curve. *Ann Hum Biol* 1978; 5: 1–24. DOI: 10.1080/03014467800002601
- [7] van der Sluis A, Elferink-Gemser MT, Coelho-e-Silva MJ et al. Sport injuries aligned to peak height velocity in talented pubertal soccer players. *Int J Sports Med* 2014; 35: 351–355. DOI: 10.1055/s-0033-1349874
- [8] Bult HJ, Barendrecht M, Tak IJR. Injury risk and injury burden are related to age group and peak height velocity among talented male youth soccer players. *Orthop J Sports Med* 2018; 6: 2325967118811042. DOI: 10.1177/2325967118811042
- [9] Johnson DM, Williams S, Bradley B et al. Growing pains: Maturity associated variation in injury risk in academy football. *Eur J Sport Sci* 2020; 20: 544–552. DOI: 10.1080/17461391.2019.1633416
- [10] Monasterio X, Gil SM, Bidaurrezaga-Letona I et al. The burden of injuries according to maturity status and timing: A two-decade study with 110 growth curves in an elite football academy. *Eur J Sport Sci* 2023; 23: 1–11. DOI: 10.1080/17461391.2021.2006316
- [11] Kemper GL, van der Sluis A, Brink MS et al. Anthropometric injury risk factors in elite-standard youth soccer. *Int J Sports Med* 2015; 36: 1112–1117. DOI: 10.1055/s-0035-1555778
- [12] Rommers N, Rossler R, Goossens L et al. Risk of acute and overuse injuries in youth elite soccer players: Body size and growth matter. *J Sci Med Sport* 2020; 23: 246–251. DOI: 10.1016/j.jsams.2019.10.001
- [13] Rommers N, Rossler R, Shrier I et al. Motor performance is not related to injury risk in growing elite-level male youth football players. A causal inference approach to injury risk assessment. *J Sci Med Sport* 2021; 24: 881–885. DOI: 10.1016/j.jsams.2021.03.004
- [14] Johnson DM, Cumming SP, Bradley B et al. The influence of exposure, growth and maturation on injury risk in male academy football players. *J Sports Sci* 2022; 40: 1127–1136. DOI: 10.1080/02640414.2022.2051380
- [15] Monasterio X, Gil S, Bidaurrezaga-Letona I et al. Peak height velocity affects injury burden in circa-PHV soccer players. *Int J Sports Med* 2023; 44: 292–297. DOI: 10.1055/a-1983-6762
- [16] Beunen GP, Rogol AD, Malina RM. Indicators of biological maturation and secular changes in biological maturation. *Food Nutr Bull* 2006; 27: S244–S256. DOI: 10.1177/156482650602745508
- [17] Johnson A, Doherty PJ, Freemont A. Investigation of growth, development, and factors associated with injury in elite schoolboy footballers: prospective study. *BMJ* 2009; 338: b490. DOI: 10.1136/bmj.b490

- [18] Le Gall F, Carling C, Reilly T. Biological maturity and injury in elite youth football. *Scand J Med Sci Sports* 2007; 17: 564–572. DOI: 10.1111/j.1600-0838.2006.00594.x
- [19] Materne O, Chamari K, Farooq A et al. Association of skeletal maturity and injury risk in elite youth soccer players: A 4-season prospective study with survival analysis. *Orthop J Sports Med* 2021; 9: 2325967121999113. DOI: 10.1177/2325967121999113
- [20] Monasterio X, Bidaurrezaga-Letona I, Larruskain J et al. Relative skeletal maturity status affects injury burden in U14 elite academy football players. *Scand J Med Sci Sports* 2022; 32: 1400–1409. DOI: 10.1111/sms.14204
- [21] Wik EH, Lolli L, Chamari K et al. Injury patterns differ with age in male youth football: a four-season prospective study of 1111 time-loss injuries in an elite national academy. *Br J Sports Med* 2021; 55: 794–800. DOI: 10.1136/bjsports-2020-103430
- [22] Materne O, Chamari K, Farooq A et al. Injury incidence and burden in a youth elite football academy: a four-season prospective study of 551 players aged from under 9 to under 19 years. *Br J Sports Med* 2021; 55: 493–500. DOI: 10.1136/bjsports-2020-102859
- [23] Hall ECR, Larruskain J, Gil SM et al. An injury audit in high-level male youth soccer players from English, Spanish, Uruguayan and Brazilian academies. *Phys Ther Sport* 2020; 44: 53–60. DOI: 10.1016/j.ptsp.2020.04.033
- [24] Fuller CW, Ekstrand J, Junge A et al. Consensus statement on injury definitions and data collection procedures in studies of football (soccer) injuries. *Br J Sports Med* 2006; 40: 193–201. DOI: 10.1136/bjism.2005.025270
- [25] Wik EH, Materne O, Chamari K et al. Involving research-invested clinicians in data collection affects injury incidence in youth football. *Scand J Med Sci Sports* 2019; 29: 1031–1039. DOI: 10.1111/sms.13427
- [26] Meeuwisse WH, Wiley JP. The Sport Medicine Diagnostic Coding System. *Clin J Sport Med* 2007; 17: 205–207. DOI: 10.1097/JSM.0b013e318057518f
- [27] Orchard JW, Meeuwisse W, Derman W et al. Sport Medicine Diagnostic Coding System (SMDCS) and the Orchard Sports Injury and Illness Classification System (OSIICS): revised 2020 consensus versions. *Br J Sports Med* 2020; 54: 397–401. DOI: 10.1136/bjsports-2019-101921
- [28] Bahr R, Clarsen B, Derman W et al. International Olympic Committee consensus statement: methods for recording and reporting of epidemiological data on injury and illness in sport 2020 (including STROBE Extension for Sport Injury and Illness Surveillance (STROBE-SIIS)). *Br J Sports Med* 2020; 54: 372–389. DOI: 10.1136/bjsports-2019-101969
- [29] Stewart A, Marfell-Jones M, Olds T et al. *International Standards for Anthropometric Assessment*. 3rd ed. Lower Hutt, NZ: International Society for the Advancement of Kinanthropometry; 2011
- [30] Lolli L, Johnson A, Monaco M et al. Tanner-Whitehouse and Modified Bayley-Pinneau adult height predictions in elite youth soccer players from the Middle East. *Med Sci Sports Exerc* 2021; 53: 2683–2690. DOI: 10.1249/MSS.0000000000002740
- [31] Roche AF, Chumlea WC, Thissen D. *Assessing the skeletal maturity of the hand-wrist: Fels method*. Springfield, IL: CC Thomas; 1988
- [32] Teunissen JWA, Rommers N, Pion J et al. Accuracy of maturity prediction equations in individual elite male football players. *Ann Hum Biol* 2020; 47: 409–416. DOI: 10.1080/03014460.2020.1783360
- [33] Lloyd RS, Oliver JL, Faigenbaum AD et al. Chronological age vs. biological maturation: implications for exercise programming in youth. *J Strength Cond Res* 2014; 28: 1454–1464. DOI: 10.1519/JSC.0000000000000391
- [34] Burnham KP, Anderson DR. *Model Selection and Multimodel Inference*. 2nd ed. New York, USA: Springer-Verlag; New York; 2004. DOI: 10.1007/b97636
- [35] Sharma J, Weston M, Batterham AM et al. Gait retraining and incidence of medial tibial stress syndrome in army recruits. *Med Sci Sports Exerc* 2014; 46: 1684–1692. DOI: 10.1249/MSS.0000000000000290
- [36] Malina RM. Skeletal age and age verification in youth sport. *Sports Med* 2011; 41: 925–947. DOI: 10.2165/11590300-000000000-00000
- [37] Quatman-Yates CC, Quatman CE, Meszaros AJ et al. A systematic review of sensorimotor function during adolescence: a developmental stage of increased motor awkwardness? *Br J Sports Med* 2012; 46: 649–655. DOI: 10.1136/bjism.2010.079616
- [38] Read PJ, Oliver JL, De Ste Croix MB et al. The scientific foundations and associated injury risks of early soccer specialisation. *J Sports Sci* 2016; 34: 2295–2302. DOI: 10.1080/02640414.2016.1173221
- [39] Hawkins D, Metheny J. Overuse injuries in youth sports: biomechanical considerations. *Med Sci Sports Exerc* 2001; 33: 1701–1707. DOI: 10.1097/00005768-200110000-00014
- [40] Cole TJ. The development of growth references and growth charts. *Ann Hum Biol* 2012; 39: 382–394. DOI: 10.3109/03014460.2012.694475
- [41] Monasterio X, Gil SM, Bidaurrezaga-Letona I et al. Injuries according to the percentage of adult height in an elite soccer academy. *J Sci Med Sport* 2021; 24: 218–223. DOI: 10.1016/j.jsams.2020.08.004
- [42] Brukner P, Khan KM. *Brukner & Khan's Clinical Sports Medicine*. 4th ed. New York, NY: McGraw-Hill Medical; 2012
- [43] DiFiori JP, Benjamin HJ, Brenner JS et al. Overuse injuries and burnout in youth sports: a position statement from the American Medical Society for Sports Medicine. *Br J Sports Med* 2014; 48: 287–288. DOI: 10.1136/bjsports-2013-093299
- [44] Bergeron MF, Mountjoy M, Armstrong N et al. International Olympic Committee consensus statement on youth athletic development. *Br J Sports Med* 2015; 49: 843–851. DOI: 10.1136/bjsports-2015-094962
- [45] Delecroix B, Delaval B, Dawson B et al. Workload and injury incidence in elite football academy players. *J Sports Sci* 2019; 37: 2768–2773. DOI: 10.1080/02640414.2019.1584954
- [46] Brink MS, Visscher C, Arends S et al. Monitoring stress and recovery: new insights for the prevention of injuries and illnesses in elite youth soccer players. *Br J Sports Med* 2010; 44: 809–815. DOI: 10.1136/bjism.2009.069476
- [47] Engebretsen L, Steffen K, Bahr R et al. The International Olympic Committee Consensus statement on age determination in high-level young athletes. *Br J Sports Med* 2010; 44: 476–484. DOI: 10.1136/bjism.2010.073122
- [48] Bittencourt NFN, Meeuwisse WH, Mendonca LD et al. Complex systems approach for sports injuries: moving from risk factor identification to injury pattern recognition-narrative review and new concept. *Br J Sports Med* 2016; 50: 1309–1314. DOI: 10.1136/bjsports-2015-095850