Red Bull PET/CT Red Bull PET/CT

Introduction

Current guidelines recommend to avoid intake of coffee prior to injection of [¹⁸F]fluorodeoxyglucose ([¹⁸F]FDG), as this may hamper uptake in sites of disease and also increase physiological biodistribution, thereby limiting diagnostic accuracy [1]. Red Bull is nowadays one of the most popular soft drinks with the highest rates of brand usage and brand awareness among all generations [2]. In the present case, we report on an unintentional ingestion of this energy drink right after injection of the radiopharmaceutical, thereby emphasizing the need to inform patients about avoiding intake of these types of beverages not only prior to, but also after tracer injection.

water, sugar (12 g/100 ml), caffeine (80 mg), 2-aminoethanesulfonic acid (taurine, 1000 mg) and glucuronolactone (600 mg) [3, 4] and has been advocated to increase (an)aerobic exercise performance [5]. Given its rather low amount of sugar relative to other energy drinks [6], one may speculate whether additional ingredients may have contributed to the high uptake in skeletal muscles observed in the present case report. For instance, taurine can be involved in muscle contraction by increasing sarcoplasmic reticulum calcium aggregation and release [7]. As muscle fibers adjust their contractility by regulating intracellular levels of taurine [4, 7], elevated blood concentration of



this amino acid after Red Bull intake may also explain increased accumulation of [¹⁸F]FDG in the muscles. Nonetheless, it remains unclear how fast taurine is absorbed from the digestive tract or incorporated in muscular cells [4]. As such, beyond potential additive effects of taurine combined with caffeine, the observed high skeletal muscle uptake could also be predominantly explained by caffeine alone [4]. The latter ingredient can lead to increased exercise endurance by preservation of muscle glycogen and neuroendocrine activation [8, 9] and thus, current guidelines also recommend refraining from intake of coffee prior to [¹⁸F]FDG PET [1]. Taken together, beyond sugar,

Case Report

We herein report on a patient with Hodgkin Lymphoma who was scheduled for [¹⁸F]FDG PET/CT as part of routine followup after treatment with two cycles of chemotherapy and mediastinal external beam radiation. Following current guidelines [1], the patient was advised to stop any sugar or caffeine intake at least 4 h prior to molecular imaging. A blood glucose level of 97 mg/dl was recorded upon arrival at our PET/CT center. As shown on [18F]FDG maximum intensity projection in (A) of Fig. 1, increased uptake in large muscles along with reduced brain, increased myocardial and almost absent liver uptake was revealed, thereby questioning adherence to the fasting period. The patient then reported on ingestion of an energy drink (Red Bull) right after radiotracer administration. After having strictly followed respective dietary recommendations, the scan was repeated (> Fig. 1B), which showed normal [18F]FDG biodistribution and response to treatment.

Discussion

As one of the most popular energy drinks [2], a 250 ml can of Red Bull consists of



▶ Fig. 1 Patient with Hodgkin Lymphoma who was scheduled for [18F]FDG PET/CT. In (A), increased uptake in large muscles along with reduced brain, increased myocardial and almost absent liver uptake was revealed, thereby questioning adherence to the fasting period. The patient then reported on ingestion of an energy drink (Red Bull) right after radiotracer administration. After having strictly followed respective dietary recommendations, the scan was repeated (B), which showed normal [18F]FDG biodistribution and response to treatment.

the excitatory effects caused by taurine, its precursor glucuronolactone, and caffeine may have all contributed to the observed increased uptake in the muscles. Thus, patients should be advised to avoid such beverages not only prior to, but also right after [¹⁸F]FDG injection, even if labeled as "sugar-free", as those drinks may still include the afore-mentioned ingredients potentially causing increased muscular radiotracer accumulation.

Funding

German Research Foundation (453989101,507803309) | Okayama University ("RECTOR") | Japan Society for the Promotion of Science (22H03027) | http:// dx.doi.org/10.13039/501100001691

Conflict of Interest

The authors declare that they have no conflict of interest.

Authors

Sebastian E Serfling¹, Andreas Buck¹, Steven P Rowe², Takahiro Higuchi¹, Rudolf Werner^{1, 3}

- ¹ Department of Nuclear Medicine, University Hospital Würzburg, Würzburg, Germany
- ² Division of Molecular Imaging and Therapeutics, Department of Radiology, University of North Carolina, Chapel Hill, United States
- ³ Goethe University Frankfurt, University Hospital, Department of Nuclear Medicine, Clinic for Diagnostic and Interventional Radiology and Nuclear Medicine, Germany

Correspondence

Dr. Rudolf Werner

Department of Nuclear Medicine University Hospital Würzburg Würzburg Germany werner_r1@ukw.de

References

- Boellaard R, Delgado-Bolton R, Oyen WJ et al. FDG PET/CT: EANM procedure guidelines for tumour imaging: version 2.0. Eur J Nucl Med Mol Imaging 2015; 42 (2): 328–354. doi:10.1007/s00259-014-2961-x
- Accessed August 15, 2023 at: https://www. statista.com/insights/consumer/brand-pro files/2/5/energy-drinks/united-states/
- [3] Laquale KM. Red Bull: The other energy drink and its effect on performance. Athletic Therapy Today 2007; 12 (2): 43–45
- [4] Kim W. Debunking the effects of taurine in red bull energy drink. Nutr Bytes 2003; 9 (1)
- [5] Alford C, Cox H, Wescott R. The effects of red bull energy drink on human performance and mood. Amino Acids 2001; 21 (2): 139–150. doi:10.1007/s007260170021
- [6] Hashem KM, He FJ, MacGregor GA. Cross-sectional surveys of the amount of sugar, energy and caffeine in sugar-sweetened drinks marketed and consumed as energy drinks in the UK between 2015 and 2017: monitoring reformulation progress. BMJ Open 2017; 7 (12): e018136. doi:10.1136/bmjopen-2017-018136
- [7] Bakker AJ, Berg HM. Effect of taurine on sarcoplasmic reticulum function and force in skinned fast-twitch skeletal muscle fibres of the rat. J Physiol 2002; 538: 185–194. doi:10.1113/jphysiol.2001.012872

- [8] Laurent D, Schneider KE, Prusaczyk WK et al. Effects of caffeine on muscle glycogen utilization and the neuroendocrine axis during exercise. J Clin Endocrinol Metab 2000; 85 (6): 2170–2175
- [9] Adler GK. Exercise and fatigue–is neuroendocrinology an important factor? J Clin Endocrinol Metab 2000; 85 (6): 2167–2169. doi:10.1210/jcem.85.6.6680

Bibliography

Nuklearmedizin 2024; 63: 76–77 published online December 22, 2023 **DOI** 10.1055/a-2195-0138 **ISSN** 0029-5566 © 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https:// creativecommons.org/licenses/by/4.0/).

