


Y. UOZUMI,* Y. MATSUURA, T. ARAKAWA, Y. M. A. YAMADA (INSTITUTE FOR MOLECULAR SCIENCE, AICHI AND RIKEN, SAITAMA, JAPAN)

Asymmetric Suzuki–Miyaura Coupling in Water with a Chiral Palladium Catalyst Supported on an Amphiphilic Resin

Angew. Chem. Int. Ed. **2009**, *48*, 2708–2710.

Asymmetric Suzuki–Miyaura Coupling in Water Using a Chiral Polymeric Pd Catalyst



Significance: Asymmetric Suzuki–Miyaura coupling in water with a novel recyclable palladium complex of a polymer-supported chiral imidazoindole phosphine ligand was described. Thus, PS-PEG-supported imidazoindole dicyclohexylphosphine (PS-PEG-L*, **7**) was prepared from PS-PEG-NH₂ and compound **6**. The reaction of 2-methyl-1-iodonaphthalene with 2-methylnaphthyl boronic acid was catalyzed by the amphiphilic polymeric PS-PEG-L*-Pd complex to give 2,2'-dimethyl-1,1'-binaphthyl in 95% yield with 94% ee (S). The catalyst PS-PEG-L*-Pd was recovered readily and reused four times with little loss of its catalytic activity and stereoselectivity.

Comment: The highly stereoselective asymmetric Suzuki–Miyaura coupling affording axially chiral biaryls is an eagerly awaited reaction to stand by itself. Heterogeneous and aqueous switching of the asymmetric coupling were realized for the first time via anchoring the catalyst onto an amphiphilic polymer resin beads. Thus, this communication should become a focus of attention from both viewpoints of asymmetric catalysis and polymer-supported synthesis. Coupling of the naphthyl chloride also took place under similar conditions with excellent enantiomeric selectivity.

SYNFACTS Contributors: Yasuhiro Uozumi, Toshimasa Suzuki
Synfacts 2009, 6, 0687-0687 Published online: 25.05.2009
DOI: 10.1055/s-0029-1216762; Reg-No.: Y05509SF

2009 © THIEME STUTTGART • NEW YORK

Category

Polymer-Supported Synthesis

Key words

Suzuki–Miyaura coupling
palladium

SYNFACTS
of the month