


O. VECHORKIN, D. BARMAZ, V. PROUST (ECOLE POLYTECHNIQUE FÉDÉRALE DE LAUSANNE, SWITZERLAND)

Ni-Catalyzed Sonogashira Coupling of Nonactivated Alkyl Halides: Orthogonal Functionalization of Alkyl Iodides, Bromides, and Chlorides

J. Am. Chem. Soc. **2009**, *131*, 12078-12079.

Ni-Catalyzed Sonogashira Coupling of Nonactivated Alkyl Halides












X = Cl, Br, I

R¹ = functionalized Alk, Bn, etc.

R² = functionalized Alk

Examples:



Significance: The Sonogashira cross-coupling of nonactivated, β -H-containing alkyl halides is a highly challenging reaction. Use of nickel(II) pincer complex **1**, developed by the authors, allows for a simple and versatile coupling of functionalized alkyl halides, including chlorides, with various terminal alkynes. This is an excellent method for the preparation of internal alkynes.

Comment: The nickel(II) pincer complex **1** is also a versatile catalyst for the Kumada $\text{sp}^3\text{-sp}^2$ coupling (O. Vechorkin, V. Proust, X. L. Hu *J. Am. Chem. Soc.* **2009**, *131*, 9756). Contrary to the classical Sonogashira coupling, aryl halides are not reactive under these conditions. This method is more convenient than the classical alkyne alkylation, which requires harsh basic conditions.

SYNFACTS Contributors: Paul Knochel, Andrei Gavryushin
Synfacts 2009, 12, 1383-1383 Published online: 20.11.2009
DOI: 10.1055/s-0029-1218170; Reg-No.: P14109SF

2009 © THIEME STUTTGART • NEW YORK

Category

Metal-Mediated
Synthesis

Key words

Sonogashira
coupling

terminal alkynes

alkylation

alkyl halides

nickel

SYNFACTS
of the month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.

1383