

# New Aspects of DNA-based Authentication of Chinese Medicinal Plants by Molecular Biological Techniques

## Author

Günther Heubl

## Affiliation

Biology Department I – Systematic Botany, Ludwig-Maximilians University Munich, Munich, Germany

## Key words

- authentication
- species identification
- Chinese medicinal plants
- standardization
- molecular markers
- DNA technologies

## Abstract

DNA technology provides a powerful tool to complement chemical analyses for authentication of Chinese medicinal plants and to ensure that herbal materials are not contaminated with ineffective or potentially harmful substitutes or adulterants. In the last two decades molecular biotechnology has provided sophisticated molecular techniques for authentication of botanical materials at the DNA level. This review provides an account of the

most commonly used DNA-based technologies (RAPD, RFLP, ARMS, CAPS, AFLP, DAF, ISSR, SSR, sequencing, hybridization and microarrays) including suitable examples of Chinese medical plants. A critical evaluation of all methods is presented concerning sensitivity, reliability, reproducibility, and running costs. Recent achievements in the field of DNA barcoding and DNA chip technology that offer great potentials for screening of DNA and emerging new developments for future identification of species are briefly outlined.

## Introduction

China's plant diversity is exceptionally rich. The flora of China contains approximately 31 500 species of vascular plants representing nearly one-eighth of the world's total plant species [1, 2]. It is estimated that 10 000 species are endemic and that more than 5000 Chinese species are used for therapy in traditional Chinese medicine [3, 4]. Chinese medicinal herbs have been used for millennia and are becoming increasingly popular in the western world. Consequently, there is an enormous demand for medicinal plants.

A severe problem on the global market is that many erroneous substitutes and adulterants are traded due to their lower costs or misidentification of species with similar morphological features. There are several cases of Chinese herbs for which substitutes have been documented and reports that some of the adulterants or substitutes caused serious intoxications and even deaths [5–10].

Consequently, the authentication of Chinese medicinal plants depending on the correct identification of the species is an essential prerequisite to ensure safety, herbal drug quality, and therapeutic efficacy [11]. Identification of herbal materials, which commonly consist of dried or processed parts, is generally difficult. This is particularly true for several species which have the same

name or similar looking herbal materials that can often vary remarkably in their medicinal properties [12–15]. In practice, the identification of medicinal plants relies mainly on morphological and chemical analyses. Many pharmacopoeias [16–18] refer to macroscopic and microscopic evaluation (morphology, histology) and chemical profiling (TLC-, HPLC-, and GC-fingerprinting) for quality control and standardization of raw and processed herbs [19, 20]. However, chemical variability within the plant material often hinders the confirmation of its botanical identity as the chemical composition is affected by growth and storage conditions as well as by the harvesting process. Otherwise microscopic examination of drugs requires botanical expertise for unequivocal authentication as related species often possess similar features.

With the improvements in molecular biotechnology and plant genetics in the past decades, genetic tools are considered to provide more reliability for the authentication of herbal materials at the DNA level. Thus various DNA-based molecular marker techniques are meanwhile applied in many fields and their application is remarkably increasing for species characterization in medicinal plants [21–24]. This is especially useful in case of those taxa that are frequently substituted or adulterated with other species or varieties that

received May 12, 2010  
revised October 4, 2010  
accepted October 7, 2010

## Bibliography

DOI <http://dx.doi.org/10.1055/s-0030-1250519>  
Published online November 5, 2010  
Planta Med 2010; 76:  
1963–1974 © Georg Thieme  
Verlag KG Stuttgart · New York ·  
ISSN 0032-0943

## Correspondence

**Prof. Dr. Günther Heubl**  
Department Biologie I –  
Systematische Botanik  
LMU München  
Menzingerst. 67  
80638 München  
Germany  
Phone: +49 89 17 86 12 07  
Fax: +49 89 17 26 38  
heubl@lrz.uni-muenchen.de

are morphologically and/or phytochemically almost indistinguishable.

Benefiting in the first place from PCR techniques, DNA markers have become a powerful tool for identification and authentication of plant and animal species [25–27]. Contrary to chemical fingerprinting which is strongly influenced by age, physiological conditions, environmental factors, cultivation area, harvesting period, drying and storage conditions, DNA is an extremely stable macromolecule that is not affected by external factors and therefore can be recovered from fresh, dried and even processed biological material. Additionally, the markers are not tissue-specific and thus can be detected at any stage of organism development. Moreover, only a small amount of sample is sufficient for analysis.

Especially DNA-barcoding (an initiative by CBOL = Consortium for the Barcode of Life) uses standard DNA markers from plastid, mitochondrial, and nuclear regions to facilitate a correct taxonomic identification of species and has become a basic tool for DNA chip technology [28,29].

This review provides an account on DNA-based technologies and most commonly used assays with emphasis on those based on DNA hybridization, restriction enzymes, random PCR amplifications, species-specific PCR primers, and DNA sequencing. A critical evaluation of all methods is presented focusing on their discriminatory power, sensitivity, reproducibility, user-friendliness, and costs.

### Types of DNA Methods and Markers Used in Plant Genome Analysis

There are various types of DNA-based molecular techniques that are used to evaluate DNA polymorphism for authentication of plant taxa [23,24,26,27]. These are hybridization-based methods, polymerase chain reaction (PCR)-based methods, and sequencing-based methods (Table 1). In recent times the use of multilocus sequence analysis (MLSA), as commonly used for phylogenetic studies, has proven its discriminatory power. Additionally, DNA microarrays that contain thousands of probes are promising new developments for sensitive and high-throughput taxon identification [30,31].

#### Non-PCR-based methods

**Hybridization-based methods (RFLP):** DNA hybridization is a process in which two single stranded DNA fragments anneal into a double-stranded nucleic acid. Restriction fragment length polymorphism (RFLP) implies that a single restriction enzyme produces fragments of different lengths from the DNA marker of different strains of a species or from different related species. In a first step, genomic (or alternatively plastid) DNA is digested with one or two selected restriction enzymes, and the fragments are separated through electrophoresis on an agarose gel. The resulting differential DNA fragment profile is then transferred to a matrix (e.g., nitrocellulose or nylon membranes) and hybridized with a chemically labelled DNA probe under conditions favoring DNA-DNA hybridization. The fragments to which the probe has hybridized are fluorescently labelled or sometimes linked with enzymes that catalyze a color reaction (digoxigenin – alkaline phosphatase) [32]. Polymorphisms are detected by the presence or absence of bands. The RFLP markers are relatively polymorphic, codominantly inherited, and highly reproducible. The method also provides an opportunity to simultaneously screen numerous

samples. DNA blots can be analyzed repeatedly by stripping and reprobing with different RFLP probes. The technique is time-consuming, labor-intensive and requires a large quantity of high amounts of good quality or undegraded DNA.

RFLP combined with DNA hybridization has been mainly used for phylogenetic studies in the past, e.g., in *Lupinus* [33], *Hedysarum* [34], *Triticum* [35], *Musa* [36] and for detection of *Dendrobium* [37] and *Fritillaria* species [38].

#### PCR-based methods

PCR-based markers involve amplification of particular DNA loci, with the help of specific or arbitrary oligonucleotide primers and a thermostable DNA polymerase enzyme. The major advantages of PCR techniques are that mainly only a small amount of DNA is required, no prior sequence information is needed, and many genetic markers can be generated within a short time. Depending on the primers used for amplification, the PCR-based techniques can be grouped into 1) arbitrary or semi-arbitrary primed PCR techniques that need no prior sequence information (e.g., AP-PCR, DAF, RAPD, AFLP, ISSR); and 2) site-targeted PCR techniques that are developed from known DNA sequences (e.g., CAPS, SSR, SCAR, STS).

**RAPD (random amplified polymorphic DNA):** The RAPD technology utilizes low-stringency polymerase chain reaction (PCR) amplification with single, short, and arbitrary synthetic oligonucleotide primers (usually 10 bp length) to generate a high number of anonymous DNA fragments. DNA polymorphism can be due to mismatches at the primer site, appearance of a new primer site, and the length of the amplified region between primer sites. Amplification products are generally separated on agarose gels and stained with ethidium bromide. Decamer primer kits are commercially available from various sources. RAPDs are inherited as dominant-recessive characters which means that homozygotes and heterozygotes cannot be distinguished. Low expense, efficiency in developing large number of DNA markers in a short time, and requirement of basic molecular equipment has made the RAPD technique valuable although band reproducibility, problems of comigration, and scoring errors are a major problem.

The technique has been applied in many plant groups such as *Glycyrrhiza* [42], *Atractylodes* [43,44], *Astragalus* [45], *Amomum* [46], *Scutellaria* [47], *Panax* [48–51], *Aconitum* [52], *Ginkgo* [53], *Anectochilus* [54], *Lycium* [55], *Angelica* [56], *Bupleurum* [57], *Dendrobium* [58], *Magnolia* [59], *Asarum* [60], *Apocynum* [61], *Trollius* [62], *Phyllanthus* [63], *Indigofera* [64], *Coptis* [65], *Codonopsis* [66], *Taraxacum* [67], *Elephantopus* [68], and *Rehmannia* [69].

**AP-PCR (arbitrary polymerase chain reaction):** AP-PCR (or arbitrarily chosen primers ACP-PCR) is a special case of RAPD using single primers approximately 10–50 bp in length [39]. In AP-PCR the amplification is in three parts. In the first two cycles annealing is under nonstringent conditions. Higher primer concentrations are used in the first cycle. Often primers of variable length are used, and products are mostly analyzed on polyacrylamide gels. AP-PCR has been applied in various groups for identification of species and analysis of genetic variation [40,41].

**DAF (DNA amplification fingerprinting):** As with arbitrarily primed polymerase chain reaction (AP-PCR), DNA amplification fingerprinting (DAF) is also an independently developed methodology, which is a variant of RAPD. For PCR amplification, very short primers of 5–8 nucleotides are used that produce a very complex banding pattern [70,71]. DAF requires careful optimization of parameters and only two temperature cycles are required.

**Table 1** Quick overview of DNA based methods or technologies, investigated TCM plant taxa (genus), and reference.

DNA method	Gene	TCM plant Systematic name–genus	References	
Hybridization- RFLP		<i>Lupinus</i>	[33]	
		<i>Hedysarum</i>	[34]	
		<i>Triticum</i>	[35]	
		<i>Musa</i>	[36]	
		<i>Dendrobium</i>	[37]	
RAPD		<i>Fritillaria</i>	[38]	
		<i>Glycyrrhiza</i>	[42]	
		<i>Atractylodes</i>	[43, 44]	
		<i>Astragalus</i>	[45]	
		<i>Amomum</i>	[46]	
		<i>Scutellaria</i>	[47]	
		<i>Panax</i>	[48–51]	
		<i>Aconitum</i>	[52]	
		<i>Ginkgo</i>	[53]	
		<i>Anectochilus</i>	[54]	
		<i>Lycium</i>	[55]	
		<i>Angelica</i>	[56]	
		<i>Bupleurum</i>	[57]	
		<i>Dendrobium</i>	[58]	
		<i>Magnolia</i>	[59]	
		<i>Asarum</i>	[60]	
		<i>Apocynum</i>	[61]	
		<i>Trollius</i>	[62]	
		<i>Phyllanthus</i>	[63]	
		<i>Indigofera</i>	[64]	
	<i>Coptis</i>	[65]		
	<i>Codonopsis</i>	[66]		
	<i>Taraxacum</i>	[67]		
	<i>Elephantopus</i>	[68]		
	<i>Rehmannia</i>	[69]		
ISSR		<i>Dendrobium</i>	[75]	
		<i>Cistanche</i>	[76]	
		<i>Fritillaria</i>	[77]	
		<i>Salvia</i>	[78]	
		<i>Vitex</i>	[79]	
		<i>Cannabis</i>	[80]	
		<i>Rhodiola</i>	[81]	
		<i>Houttunya</i>	[82]	
	AFLP		<i>Panax</i>	[87, 88]
			<i>Actaea</i>	[89]
		<i>Plectranthus</i>	[90]	
		<i>Caladium</i>	[91]	
		<i>Cannabis</i>	[92]	
RAMPO		<i>Rehmannia</i>	[93]	
		<i>Phoenix</i>	[95]	
		<i>Ficus</i>	[96]	
Sequencing ITS		<i>Panax</i>	[101, 102]	
		<i>Asarum</i>	[103–105]	
		<i>Astragalus</i>	[106, 107, 171]	
		<i>Dendrobium</i>	[108–111]	
		<i>Fritillaria</i>	[112]	
		<i>Leonurus</i>	[113]	
		<i>Perilla</i>	[114]	
		<i>Rehmannia</i>	[115]	
		<i>Salvia</i>	[116]	
		<i>Swertia</i>	[117]	
		<i>Plantago</i>	[118]	
		<i>Bupleurum</i>	[119]	
		<i>Euphorbia</i>	[120, 121]	
		<i>Hedysarum</i>	[171]	

**Table 1** Quick overview of DNA based methods or technologies, investigated TCM plant taxa (genus), and reference. (continued)

DNA method	Gene	TCM plant Systematic name–genus	References						
5S rDNA		<i>Adenophora</i>	[122]						
		<i>Aconitum</i>	[123]						
		<i>Angelica</i>	[124]						
		<i>Astragalus</i>	[125, 126]						
		<i>Curcuma</i>	[127]						
		<i>Epimedium</i>	[128]						
		<i>Fritillaria</i>	[129]						
		<i>Crocus</i>	[130]						
		<i>Ligularia</i>	[131]						
		<i>Pueraria</i>	[132]						
		<i>Saussurea</i>	[133]						
	18S rDNA		<i>Dioscorea</i>	[134]					
			<i>Pinellia</i>	[135]					
			<i>Panax</i>	[136]					
			<i>Fritillaria</i>	[137]					
		<i>Phyllanthus</i>	[138]						
		<i>Panax</i>	[139]						
26S rDNA		<i>Panax</i>	[139]						
		<i>Pueraria</i>	[140]						
		<i>Rheum</i>	[141]						
		<i>Ephedra</i>	[142]						
		<i>Artemisia</i>	[163]						
		<i>Swertia</i>	[143]						
		<i>Scutellaria</i>	[144]						
		<i>Angelica</i>	[145]						
		<i>Dyosma</i>	[146]						
		<i>Actinidia</i>	[147]						
atpB-rbcL		<i>Atractylodes</i>	[148]						
		<i>Curcuma</i>	[149]						
		<i>Agastache</i>	[150]						
		<i>Panax</i>	[151]						
	trnC-trnD		<i>Dryopteris</i>	[152]					
			<i>Cnidium</i>	[153]					
			<i>Pinellia</i>	[154]					
		trnL-trnF		<i>Alisma</i>	[157]				
				<i>Angelica</i>	[158]				
				<i>Sinopodophyllum</i>	[159]				
			<i>Dyosma</i>	[159]					
			<i>Ephedra</i>	[160]					
			<i>Fritillaria</i>	[161, 162]					
			<i>Artemisia</i>	[163]					
	<i>Panax</i>		[164–167]						
	<i>Actinidia</i>		[168]						
	<i>Atractylodes</i>		[169]						
rpl16		<i>Glehnia</i>	[170]						
		<i>Astragalus</i>	[171]						
		<i>Dendrobium</i>	[172]						
		<i>Duboisia</i>	[173]						
		<i>Codonopsis</i>	[174]						
	rpl16-rpl14		<i>Stephania</i>	[176]					
			<i>Panax</i>	[177]					
		atpF-atpA		<i>Alisma</i>	[179]				
				<i>Panax</i>	[180, 181]				
				<i>Rheum</i>	[182]				
			<i>Dendrobium</i>	[183, 184]					
			<i>Curcuma</i>	[185]					
trnD-trnT				<i>Panax</i>	[190, 191]				
				<i>Artemisia</i>	[192]				
				<i>Phyllanthus</i>	[193, 194]				
			<i>Pueraria</i>	[195]					
			<i>Sinocalycanthus</i>	[196]					
	trnK		<i>Embelia</i>	[197]					
			<i>Lycium</i>	[198]					
		matK							
			rbcL						
				CAPS					
DALP									
					ARMS				
						SCAR			

continued next page

**Table 1** Quick overview of DNA based methods or technologies, investigated TCM plant taxa (genus), and reference. (continued)

DNA method	TCM plant	References
	Gene	
SSR	<i>Panax</i>	[202, 203]
	<i>Acanthopanax</i>	[204]
	<i>Dendrobium</i>	[205]
	<i>Cymbopogon</i>	[206]
	<i>Bupleurum</i>	[207]
SAMPL	<i>Schisandra</i>	[208]
	<i>Cicer</i>	[213]
	<i>Lactuca</i>	[214]
DAMD	<i>Tribulus</i>	[215]
	<i>Oryza</i>	[219]
	<i>Panax</i>	[220]
	<i>Capsicum</i>	[221]
Microarrays	<i>Salvia</i>	[222]
	<i>Morus</i>	[223]
	<i>Fritillaria</i>	[228]
	<i>Dendrobium</i>	[229, 230]
	<i>Bupleurum</i>	[231]
	<i>Panax</i>	[232]

DAF products are routinely separated by polyacrylamide gels and detected by silver staining [72].

Although DAF and AP-PCR are different with respect to the length of the random primers, amplification conditions, and visualization methods, they all differ from the standard PCR condition in that only a single oligonucleotide of the random sequence is employed and no prior knowledge of the genome is required.

**ISSR (inter-simple sequence repeat):** ISSR markers are more and more in demand, because they are known to be abundant, very reproducible, highly polymorphic, informative, and quick to use [73, 74]. Neither sequence information nor prior genetic studies are required. ISSR uses the presence throughout the genome of simple sequence repeats (SSRs) which are ubiquitous, abundant, and highly polymorphic tandem repeat motifs composed of 1 to 7 nucleotides. Inter-simple sequence repeat (ISSR) permits detection of polymorphisms in inter-microsatellite loci. The primers used can be 5' or 3' anchored by 1–3 selective nucleotides to prevent internal priming and to amplify only a subset of the targeted inter-repeat regions.

It has been used in the authentication of *Dendrobium* [75], *Cistanche* [76], *Fritillaria* [77], *Salvia* [78], *Vitex* [79], *Cannabis* [80], *Rhodiola* [81], and *Houttuynia* [82].

**AFLP (amplified fragment length polymorphism):** The AFLP technique is a powerful DNA fingerprinting technology applicable to any organism without the need for prior sequence knowledge. It is a multilocus approach combining restriction fragment analysis with selective PCR amplification [83, 84]. In a first step, total genomic DNA is digested with two or more restriction enzymes (e.g., *EcoRI*; *MseI*) producing sticky ends, and the fragments are ligated to specific adapters (oligonucleotide 10–30 bp length). The ligated DNA fragments are then amplified twice under highly stringent conditions by PCR using radioactive or fluorescence-labelled primers complementary to the adapter and restriction site sequence. Using selective primers which include additional nucleotides at their 3' end, the complexity of the mixture of fragments can be reduced. The amplicons are separated on a polyacrylamide gel, followed by visualization of the banding pattern [85]. The AFLP technique is a reliable and robust molecular

marker assay that permits the simultaneous screening of different loci randomly distributed throughout the genome [86]. It is very efficient in revealing polymorphisms even between closely related individuals. However, degraded DNA can lead to wrong banding patterns. It has been used for recognition of individuals (paternity analysis, selfing rates, identification of cultivars, clones, etc.) and studying genetic diversity of Chinese medicinal plants. AFLP-based techniques were used in *Panax* [87, 88], *Actaea* [89], *Plectranthus* [90], *Caladium* [91], *Cannabis* [92], and *Rehmannia* [93].

**RAMPO (random amplified microsatellite polymorphisms):** This method, also termed RAHM (random amplified hybridization microsatellite) or RAMS (randomly amplified microsatellites), combines arbitrarily primed PCR (RAPD) with microsatellite hybridization to produce polymorphic genetic fingerprints [94]. No prior sequence information is needed. Genomic DNA is first amplified with a single arbitrary or microsatellite-complementary PCR primer. After electrophoretic separation and staining of the PCR products, the gel is either dried or blotted onto a nylon membrane, and subsequently hybridized to a labelled, microsatellite-complementary oligonucleotide probe (e.g., [GT]8 or [GA]8). The method is mainly used for identification and discrimination of genotypes within and among populations, cultivars, germplasm, etc. [95, 96].

### PCR and DNA sequencing-based methods

DNA sequencing is a powerful tool to characterize species, analyze phylogenetic relationships, population genetics, and evolutionary processes [97]. DNA polymorphisms are revealed by determining the nucleotide sequence in a defined region of the genome and aligning the sequence with homologous regions of related organisms [98].

By choosing appropriate regions of the nuclear, plastid, or mitochondrial genome this approach provides a highly reproducible analysis at various taxonomic ranks to differentiate TCM plants from its substitutes or adulterants.

In order to provide security for a correct species identification based on DNA sequence data, it is necessary to have a herbarium specimen for verification or a reliable database that guarantees that the reference specimen was correctly identified by a taxonomic expert. Additionally, the sequence should be obtained in independent studies including related taxa. A common way to assign a particular sequence to a taxon is to perform a BLAST search (basic local alignment search tool) in the GenBank database of NCBI. However, care must be taken when assigning the questioned sequence to the species with the highest similarity, because several gaps and false sequences are known to be present in these databases.

There are many studies concerning the application of DNA sequence-based markers to differentiate medicinal taxa used in TCM from its substitutes or adulterants. The ITS region of 18S–26S rDNA has proven to be an important and useful gene locus for DNA barcoding applications [99, 100]. Especially the ITS2 locus was revealed to discriminate among 4800 species with a probability of 92.7% [99, 100].

Sequencing analyses based on this nuclear marker have been applied to *Panax* [101, 102], *Asarum* [103–105], *Astragalus* [106, 107], *Dendrobium* [108–111], *Fritillaria* [112], *Leonurus* [113], *Perilla* [114], *Phyllanthus* [115], *Rehmannia* [116], *Salvia* [117], *Swertia* [118], *Plantago* [119], *Bupleurum* [120], and *Euphorbia* [121].

Another frequently used marker is the nuclear 5S rDNA intergenic spacer used for authentication of *Adenophora* [122], *Aconitum* [123], *Angelica* [124], *Astragalus* [125, 126], *Curcuma* [127], *Epimedium* [128], *Fritillaria* [129], *Crocus* [130], *Ligularia* [131], *Pueraria* [132], and *Saussurea* [133].

From nuclear DNA also 18S rDNA has been tested in *Dioscorea* [134], *Pinellia* [135] and *Panax* [136], the 26S rDNA marker in *Fritillaria* [137].

From chloroplast DNA, a couple of markers including genes, intergenic spacers, or introns are applied. The *atpB-rbcL* region was used for differentiation of *Phyllanthus* [138], *trnC-trnD* in *Panax* [139], *trnL-F* in *Pueraria* [140], *Rheum* [141] and *Ephedra* [142], *rpl16* in *Swertia* [143], *rpl16-rpl14* spacer in *Scutellaria* [144], *atpF-atpA* in *Angelica* [145], *trnD-trnT* in *Dyosma* [146], *trnK* in *Actinidia* [147], *Atractylodes* [148] and *Curcuma* [149], *matK* in *Agastache* [150], *Panax* [151], *rbcL* in *Dryopteris* [152], *Cnidium* [153], and *Pinellia* [154].

**CAPS or PCR-RFLP (cleaved amplified polymorphic sequence):** CAPS, originally named PCR-RFLP is a combination of PCR of target DNA and subsequent digestion with a restriction enzyme [155, 156]. CAPS markers are generated in two steps. In the first step, a defined sequence is amplified using specific primers. In the second step, the PCR product is digested with a restriction enzyme usually with a four-base recognition specificity. The digested fragments are separated on agarose gels. However, the ability of CAPS to detect DNA polymorphism is not as high as SSRs or AFLPs because nucleotide changes affecting restriction sites are essential for the detection of DNA polymorphism by CAPS. Furthermore, the development of CAPS markers is only possible where mutations disrupt or create a restriction enzyme recognition site. PCR-RFLP has been used for authentication of *Alisma* [157], *Angelica* [158], *Sinopodophyllum* and *Dyosma* [159], *Ephedra* [160], *Fritillaria* [161, 162], *Artemisia* [163], *Panax* [164–167], *Actinidia* [168], *Atractylodes* [169], *Glehnia* [170], *Astragalus* [171], *Dendrobium* [172], *Duboisia* [173], and *Codonopsis* [174].

**DALP (direct amplification of length polymorphisms):** This method uses an arbitrarily primed PCR (AP-PCR) to produce genomic fingerprints and to enable sequencing of DNA polymorphisms in virtually any species [175]. For PCR higher stringency is necessary.

The uniqueness of DALP relies upon the specific design of primer pairs. It uses a selective forward primer containing a 5' core sequence of the universal M13 sequencing primer plus additional bases (usually 2–5) at the 3' end, and a common reverse M13 primer. Any of the bands generated by PCR can be excised from the gel and sequenced directly using forward or reverse primers. After sequencing the polymorphic bands among the samples, species specific primers can be designed. DALP has been used to detect polymorphisms between species [175] and to authenticate *Stephania yunnanensis*, [176], *Panax ginseng*, and *Panax quinquefolius* [177].

**ARMS and Multiplex-ARMS (amplification refractory mutation system):** ARMS, also known as allele-specific polymerase chain reaction (ASPCR) is a simple, timesaving, and effective method for detecting any mutation involving single base changes (SNPs) or small deletions. It has become a standard technique that allows the discrimination of alleles [178]. The basis of ARMS is that oligonucleotides with a mismatched 3'-residue will not function as primers in the PCR. ARMS allows amplification of test DNA only when the target allele is contained within the sample and will not amplify the nontarget allele. Following an ARMS reaction, the presence or absence of a PCR product is diagnostic for the presence or absence of the target allele. A main advantage of ARMS

is that the amplification step and the authentication step are combined, in that the presence or absence of a PCR product is diagnostic for the presence or absence of the target allele. The method provides a quick screening assay that does not require any form of labelling as the amplified products are visualized simply by agarose gel electrophoresis and ethidium bromide staining. Multiplex ARMS or MARMS is a similar approach but there are several primer combinations to be optimized simultaneously, which increases the complexity of the procedure. This technique has been applied in the authentication of *Alisma* [179], *Panax* [180, 181], *Rheum* [182], *Dendrobium* [183, 184], and *Curcuma* [185].

**SCAR (sequence characterized amplified region):** A SCAR can be used to rapidly amplify a diagnostic nucleic acid from herbal materials using a pair of specific oligonucleotide primers designed from polymorphic RAPD [186, 187] or ISSR [188] fragments.

Polymorphic regions from RAPDs or ISSR are selected among amplified fingerprints. After cloning and sequencing for the selected polymorphic regions, pairs of internal primers are designed to amplify a unique and specific sequence designed as a SCAR marker. PCR results in a positive or negative amplification in target-containing and nontarget-containing samples respectively or amplification products of different sizes in the case of closely related samples. SCARs are advantageous over RAPD markers as they detect only a single locus, their amplification is less sensitive to reaction conditions, and they can potentially be converted into codominant markers [189]. Prior sequence information (i.e., sequencing the polymorphic fragments) is required for designing the primers flanking the polymorphic region. As PCR inhibitory effects of ingredients can lead to false negative results, amplification of a control fragment using the same DNA template should be performed to ensure that the quality of sample DNA is suitable for PCR.

The SCAR technique has been used for authentication of *Panax* [190, 191] and for discrimination of species of *Artemisia* [192], *Phyllanthus* [193, 194], *Pueraria* [195], *Sinocalycanthus* [196], *Embelia* [197], and *Lycium* [198].

**Microsatellites (SSR – simple sequence repeats):** Hypervariable repetitive DNA sequences such as microsatellites, minisatellites or midisatellites, and satellites can be of great value in assessing a high level of polymorphism as they are distributed throughout the genomes.

Microsatellites are also known as simple sequence repeats (SSRs), short tandem repeats (STRs), or simple sequence length polymorphisms (SSLPs) are the smallest class of simple repetitive DNA sequences [199, 200]. They are highly polymorphic and abundant sequences that are dispersed throughout most eukaryotic genomes. These molecular markers are widely used for DNA fingerprinting, paternity testing, linkage map construction, and population genetic studies but of less importance for species identification [201]. Based on tandem repeats of short (2–6 bp) DNA sequences, these markers are highly polymorphic due to variation in the number of repeat units. The repeat length at specific SSR loci is easily assayed by PCR using primers specific to conserved regions flanking the repeat array. PCR fragments are usually separated on polyacrylamide gels in combination with fluorescent detection systems. The hypervariability and codominance of SSRs, their dispersion throughout genomes and suitability for automatization are the principle reasons for their wide utility. A major limitation of SSRs is the time and high development cost required to isolate and characterize each locus when a preexisting DNA sequence is not available. Typically, this process requires

the construction and screening of a genomic library of size-selected DNA fragments with SSR-specific probes, followed by DNA sequencing of isolated positive clones, PCR primer synthesis, and testing.

Microsatellites have been applied in *Panax* [202,203], *Acanthopanax* [204], *Dendrobium* [205], *Cymbopogon* [206], *Bupleurum* [207], and *Schisandra* [208].

**SAMPL (selective amplification of microsatellite polymorphic loci):** The SAMPL technique is an SSR-based modification of the amplified fragment length polymorphism (AFLP) procedure [209], but it differs from AFLP by using primers with compound microsatellite motifs in combination with oligonucleotides complementary to the end-ligated adapters for the selective amplification step [210]. In brief, genomic DNA is digested with restriction enzymes (commonly *EcoRI* and *MseI*), and the resulting fragments are ligated to adapters that contain sticky ends to the restriction sites for the enzymes at the genome fragments, and a preamplification reaction for all ligated DNA fragments is carried out with primers annealing to the adapters. These preamplified products, adequately diluted, are then used as templates for a selective SAMPL-polymerase chain reaction (PCR) that uses the adapter-primer (*EcoRI* oligo-1) in combination with an end-labelled microsatellite-based 15-mer oligonucleotide [211] to amplify a group of fragments from those that were restricted, ligated, and preamplified. This multiplexing genome profiling technique has not adequately been used in plant genomics, although a few reports have already documented its potential to detect polymorphisms [212]. This method was used for analysis of genetic diversity in *Cicer* [213], *Lactuca* [214] and *Tribulus* [215].

**DAMD (directed amplification of minisatellite-region DNA):** DAMD is a DNA fingerprinting method based on amplification of the regions rich in minisatellites at relatively high stringencies by using previously found VNTR core sequences as primers [216,217]. Minisatellites, also known as variable number of tandem repeats (VNTR) or hypervariable repeats (HVR), are similar to microsatellites (SSR) except that the tandem repeat DNA sequences are longer and generally consist of 10–60-bp motifs. Extreme variations in the tandem repeat copy number of minisatellite loci are responsible for the polymorphism observed. By using the VNTR core sequence as primers, the directed amplification of minisatellite-region DNA (DAMD) with PCR is capable of producing RAPD-like results for the identification of species [218]. It is also used to generate highly variable probes for DNA fingerprinting. This method is more reproducible than RAPD due to the longer primers used.

Recently, DAMD-PCR has been successfully applied for genotyping of wheat cultivars and rice species [219]. The method has been used for authentication of *Panax* [220], *Capsicum* [221], *Salvia* [222], and *Morus* [223].

### DNA-microarrays (DNA chip technology)

DNA microarrays or genechips are a high throughput technology for simultaneous analysis of multiple genes in many taxa or samples. To apply this technique for identification and authentication of herbal material it is necessary to identify a distinct DNA sequence that is unique to each species [224]. Based on the gained DNA sequences, corresponding probes are synthesized for many samples. These immobilized DNA fragments are arranged in a regular pattern on a microarray by fixation on glass slides, silicon or nylon membranes [225,226].

DNA extracted from the target sample and labelled with a specific fluorescent molecule is then hybridized to the microarray

DNA. A positive hybridization is detected and visualized with fluorescence scanning or imaging equipment. The microarray is scanned to obtain a complete hybridization pattern generated by the release of a fluorescent, chemiluminescent, or colorimetric signal associated with the binding of the probe to the target DNA. A number of terms such as DNA arrays, gene chips, and biochips are often used to describe these devices [227].

Recently this technique has been applied for the identification of various species of *Fritillaria* [228], *Dendrobium* [229,230], and *Bupleurum* [231]. In addition, microarray technology has also been used to authenticate *Panax ginseng* [232] and toxic traditional Chinese medicinal materials [233] such as *Aconitum*, *Strychnos*, and *Datura*.

The results demonstrated that DNA microarray based technology can provide a rapid, high-throughput tool for correct botanical identification, authentication of crude plant materials, standardization and quality control, testing simultaneously hundreds of samples [226].

### Future Developments

▼ New innovative automated assays and specific tools for DNA analysis are emerging and will contribute to the next generation of technologies. These are minisequencing [234,235], nanoscale DNA sequencing [236] or microsphere-based suspension arrays [237]. Further promising developments are the nanopore technology for identification of DNA bases with very high confidence, and the arrayed primer extension reaction (APEX) which is an enzymatic genotyping method to analyze hundreds to thousands of variations in the genome simultaneously in a single multiplexed reaction [238,239]. Another upcoming method for large-scale multiplex analysis of nucleic acid sequences is the multiplex oligonucleotide ligation assay (OLA), which can be applied for allelic discrimination in highly polymorphic genes [240].

These techniques have a high multiplexing capacity and great potentials for genotyping and future taxon identification [241].

### Limitation of DNA methods

Molecular authentication methods have several advantages which make them suitable for the identification of herbs used in TCM, as compared to macroscopic, microscopic, and phytochemical analyses. The DNA-based techniques are not affected by environmental factors, independent from the physical form of the plant material, and only a low amount of material is required. Although DNA analysis is currently considered to be cutting-edge technology, it has certain limitations.

- ▶ The applicability of a DNA-based method depends generally on the quality and quantity of the DNA, which might be a problem for dried or processed materials. Important drug-processing conditions, for example, temperature and pH, may lead to degradation (fragmentation) of the DNA, rendering PCR analysis impossible. However, depending on the degree of degradation of DNA some methods can still be used in processed materials. For these, it is necessary to develop very short amplicons to have a certain probability of successful application [242].
- ▶ High concentrations of secondary plant compounds (polysaccharides, tannins, essential oils, phenolics, alkaloids, etc.) may influence DNA extraction or PCR reaction. In tissues of medicinal plants, secondary compounds generally get accumulated and the problem becomes severe as the material gets older. Polysaccharide contaminations are particularly problematic as

they can inhibit the activity of many commonly used enzymes, such as polymerases, ligases, and restriction endonucleases. Polyphenol contamination of DNA makes it resistant to restriction enzymes and interacts irreversibly with proteins and nucleic acids. Choosing the most suitable DNA extraction procedure may help to eliminate the PCR inhibitors.

- ▶ Sometimes plant materials are contaminated with endophytic fungi, which might influence DNA sequencing and can be eliminated with a plant-specific primer design.
- ▶ DNA related methods can generally not be applied when the herb is processed to an extract.
- ▶ DNA markers, such as the internal transcribed spacer (ITS) region of the 18S-5.8S-26S nuclear ribosomal cistron, sometimes show intraspecific sequence variation due to nonfunctional paralogous sequences (pseudogenes). For DNA barcoding as a practical molecular method to identify species, only orthologous DNA sequences can be used. Consequently cloning of PCR products is sometimes inevitable.
- ▶ In order to establish a marker for identification of a particular species, DNA analysis of closely related species and/or varieties and common botanical contaminants and adulterants is necessary, which is a costly and time-consuming process.

## Conclusion

DNA technologies are reliable and powerful tools for identification of taxa at various taxonomic levels (e.g., species, subspecies, variety, strain) as they provide consistent results irrespective of age, tissue origin, physiological conditions, environmental factors, harvest, storage, and processing of samples. With the increasing demand of high-quality herbs, also the need for DNA authentication will accelerate for ensuring the therapeutic effectiveness, a fair trade of drugs and raising consumers' confidence. However, for the modernization of TCM it is inevitable in the future to compile a comprehensive database including DNA data for all investigated medicinal taxa with reference information on nomenclature, phylogenetic relationships, macroscopic and microscopic features, chemical constituents and profiling, toxicity, and voucher specimens in herbaria or museums.

## References

- 1 Lopez-Pujol JL, Zahng FM, Song GE. Plant biodiversity in China: richly varied, endangered, and in need of conservation. *Biodivers Conserv* 2006; 15: 3983–4026
- 2 Qian H, Ricklefs RE. A comparison of the taxonomic richness of vascular plants in China and the United States. *Am Nat* 1999; 154: 160–181
- 3 Zhong YD. *Big Chinese Herb Dictionary*. Shanghai Kexue Jishu Chu Ban Shi. Shanghai: Shanghai Science and Technology Publishing Co.; 1977
- 4 Lee HH, Itokawa H, Kozuka M. Asian herbal products: The basis for development of high-quality dietary supplements and new medicines. In: Shi J, Ho CT, Shahidi F, editors. *Asian Functional Foods*. Boca Raton: Pub CRC Press, Taylor & Francis Group; 2005
- 5 Zhao ZZ, Hu Y, Liang ZT, Yuen PSJ, Jiang ZH, Leung KSY. Authentication is fundamental for standardization of Chinese medicines. *Planta Med* 2006; 72: 865–874
- 6 Zhao ZZ, Li TYS. *Easily confused Chinese medicines in Hong Kong* (English edition). Hong Kong: Chinese Medicine Merchants Association Ltd.; 2007
- 7 Chan TY, Critchley JA. Usage and adverse effects of Chinese herbal medicines. *Hum Exp Toxicol* 1996; 15: 5–12
- 8 Gertner E, Marshall PS, Filandrinis D, Potek AS, Smith TM. Complications resulting from the use of Chinese herbal medications containing undeclared prescription drugs. *Arthritis Rheum* 1995; 38: 614–617
- 9 But PP, Tomlinson B, Cheung KO, Yong SP, Szeto ML, Lee CK. Adulterants of herbal products can cause poisoning. *Br Med J* 1996; 313: 117
- 10 Chen J, Chen L, An Z, Shi S, Zhan Y. Non-technical causes of fakes existing in Chinese medicinal material markets. *Zhongyaocai* 2002; 25: 516–519
- 11 Zhao Z, Hu Y, Liang Z, Yuen J, Jiang Z, Leung KSY. Authentication is fundamental for standardization of Chinese medicines. *Planta Med* 2006; 72: 865–874
- 12 Huang HH, Yen DHT, Wu ML, Deng JF, Huang CI, Lee CH. Acute *Erycibe henryi* Prain (“Ting Kung Teng”) poisoning. *Clin Toxicol* 2006; 44: 71–75
- 13 Sun SQ, Zhou Q, Liu J, Huang H. Study on the identification of standard and false BanXia by two-dimensional infrared correlation spectroscopy. *Spectrosc Spectr Anal* 2004; 24: 427–430
- 14 Zhang WH, Shen ZJ. Comparison on macroscopic characteristics of clinical curative effect of Banxia and Shuibanxia. *J Nanjing Univ Tradit Chin Med* 1995; 11: 32–33
- 15 Zhao ZZ, Li TYS. *Hong Kong Commonly Confused Chinese Medicines*. Hong Kong: Chinese Medicine Merchants Association Ltd.; 2004
- 16 *Society of Japanese Pharmacopoeia*. *Japanese Pharmacopoeia* (English version), 14th edition. Tokyo: Junkudo Book Shop; 2001
- 17 *Korean Food and Drug Administration*. *Korean Herbal Pharmacopoeia*, 8th edition. Seoul: Ministry of Health Family Welfare of South Korea; 2002
- 18 *State Pharmacopoeia Committee*. *Pharmacopoeia of China* (2005 edition). Beijing: Chemical Industry Publisher; 2005
- 19 Siow YL, Gong Y, Au-Yeung KK, Woo CW, Choy PC. Emerging issues in traditional Chinese medicine. *Can J Physiol Pharmacol* 2005; 83: 321–334
- 20 Chan K. Some aspects of toxic contaminants in herbal medicines. *Chemosphere* 2003; 52: 1361–1371
- 21 Shaw PC, Ngan FN, But PPH, Wang J. Molecular markers in Chinese medicinal materials. In: Shaw PC, But PPH, editors. *Authentication of Chinese medicinal material by DNA technology*. Singapore: World Scientific Publishing; 2002
- 22 Zhang YB, Shaw PC, Sze CW, Wang ZT, Tong Y. Molecular authentication of Chinese herbal materials. *J Food Drug Anal* 2007; 15: 1–9
- 23 Sucher JN, Carles MC. Genome-based approaches to the authentication of medicinal plants. *Planta Med* 2008; 74: 603–623
- 24 Shaw PC, Wong KKL, Chan AWK, Wong WC, But PPH. Patent applications for using DNA technologies to authenticate medicinal herbal material. *J Chin Med* 2009; 4: 1–11
- 25 Yip PY, Chau CF, Mak CY, Kwan HS. DNA methods for identification of Chinese medicinal materials. *J Chin Med* 2007; 2: 1–19
- 26 Kaplan J, Chavan P, Warude D, Patwardhan B. Molecular markers in herbal drug technology. *Curr Sci* 2004; 87: 159–165
- 27 Pereira F, Carneiro J, Amorim A. Identification of species with DNA-based technology: current progress and challenges. *Recent Pat DNA Gene Seq* 2008; 2: 187–200
- 28 Ratnasingham S, Hebert PDN. BOLD: the barcode of life data system ([www.barcodinglife.org](http://www.barcodinglife.org)). *Mol Ecol Notes* 2007; 7: 355–364
- 29 Chase MW, Salamin N, Wilkinson M, Dunwell JM, Kesanakurthi RP, Haidar N, Savolainen V. Land plants and DNA barcodes: short-term and long-term goals. *Philos Trans R Soc Lond Ser B Biol Sci* 2005; 360: 1889–1895
- 30 Trau D, Lee TM, Lao AI, Lenigk R, Hsing IM, Ip NY, Carles MC, Sucher NJ. Genotyping on a complementary metal oxide semiconductor silicon polymerase chain reaction chip with integrated DNA microarray. *Anal Chem* 2002; 74: 3168–3173
- 31 Schena M, Heller RA, Theriault TP, Konrad K, Lachenmeier E, Davis RW. Microarrays: biotechnology's discovery platform for functional genomics. *Trends Biotechnol* 1998; 16: 301–316
- 32 Lashermes P, Combes MC, Cros J. Use of non-radioactive digoxigenin-labelled DNA probes for RFLP analyses in coffee. *Technique et utilisations des marqueurs moleculaires* 1994; 72: 29–31
- 33 Yamazaki M, Sato A, Saito K, Murakoshi I. Molecular phylogeny based on RFLP and its relation with alkaloid patterns in *Lupinus* plants. *Biol Pharm Bull* 1993; 16: 1182–1184
- 34 Trifi-Farah N, Marrakchi M. *Hedysarum* phylogeny mediated by RFLP analysis of nuclear ribosomal DNA. *Genet Resour Crop Evol* 2001; 48: 339–345
- 35 Mori N, Moriguchi T, Nakamura C. RFLP analysis of nuclear DNA for study of phylogeny and domestication of tetraploid wheat. *Genes Genet Syst* 1997; 72: 153–161
- 36 Gawel NJ, Jarret RL, Whittemore AP. Restriction fragment length polymorphism (RFLP)-based phylogenetic analysis of *Musa*. *Theor Appl Genet* 1992; 84: 286–290

- 37 Li T, Wang J, Lu Z. Accurate identification of closely related *Dendrobium* species with multiple species-specific gDNA probes. *J Biochem Biophys Methods* 2005; 62: 111–123
- 38 Tsoi PY, Woo HS, Wong MS, Chen SL, Fong WF, Xiao PG, Yang MS. Genotyping and species identification of *Fritillaria* by DNA chips. *Yaoxue Xuebao* 2003; 38: 185–190
- 39 Welsh J, McClelland M. Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acids Res* 1990; 18: 7213–7218
- 40 Munthali M, Ford-Lloyd BV, Newbury HJ. The random amplification of polymorphic DNA for fingerprinting plants. *PCR Methods Appl* 1992; 1: 274–276
- 41 Kersten T, Daniel C, König GM, Knöß W. The potential of PCR-related methods to identify medicinal plants in herbal medicinal products. *Planta Med* 2007; 73: 256
- 42 Yamasaki M, Sato A, Shimomura K, Saito K, Murakoshi I. Genetic relationships among *Glycyrrhiza* plants determined by RAPD and RFLP analyses. *Biol Pharm Bull* 1994; 17: 1529–1531
- 43 Kohjyouma M, Nakajima S, Namera A, Shimizu R, Mizukami H, Kohda H. Random amplified polymorphic DNA analysis and variation of essential oil components of *Atractylodes* plants. *Biol Pharm Bull* 1997; 20: 502–506
- 44 Chen KT, Su YC, Lin JG, Hsin LH, Su YP, Su CH, Li SY, Cheng JH, Mao SJ. Identification of *Atractylodes* plants in Chinese herbs and formulations by random amplified polymorphic DNA. *Acta Pharmacol Sin* 2001; 22: 493–497
- 45 Cheng KT, Su B, Chen CT, Lin CC. RAPD analysis of *Astragalus* medicines marketed in Taiwan. *Am J Chin Med* 2000; 28: 273–278
- 46 Wang P, Huang F, Zhou L, Cao L, Liang S, Xu H, Liu J. Analysis of *Amomum villosum* species and some adulterants of Zingiberaceae by RAPD. *Zhong Yao Cai* 2000; 23: 71–74
- 47 Hosokawa K, Minami M, Kawahara K, Nakamura I, Shibata T. Discrimination among three species of medicinal *Scutellaria* plants using RAPD markers. *Planta Med* 2000; 66: 270–272
- 48 Cheung KS, Kwan HS, But PPH, Shaw PC. Pharmacognostical identification of American and Oriental ginseng roots by genomic fingerprinting using arbitrarily primed polymerase chain reaction (AC-PCR). *J Ethnopharmacol* 1994; 42: 67–69
- 49 Shim YH, Choi JH, Park CD, Lim CJ, Cho JH, Kim HJ. Molecular differentiation of *Panax* species by RAPD analysis. *Arch Pharm Res* 2003; 26: 601–605
- 50 Cui XM, Lo CK, Yip K L, Dong TT, Tsim K W. Authentication of *Panax notoginseng* by 5S-rRNA spacer domain and random amplified polymorphic DNA (RAPD) analysis. *Planta Med* 2003; 69: 584–586
- 51 Lim W, Mudge KW, Weston L. Utilization of RAPD markers to assess genetic diversity of wild populations of North American ginseng (*Panax quinquefolium*). *Planta Med* 2007; 73: 71–76
- 52 Cole CT, Kuchenreuther MA. Molecular markers reveal little genetic differentiation among *Aconitum noveboracense* and *A. columbianum* (Ranunculaceae) populations. *Am J Bot* 2001; 88: 337–347
- 53 Fan XX, Shen L, Zhang X, Chen XY, Fu CX. Assessing genetic diversity of *Ginkgo biloba* L. (Ginkgoaceae) populations from China by RAPD markers. *Biochem Genet* 2004; 42: 269–278
- 54 Cheng KT, Fu LC, Wang CS, Hsu FL, Tsay HS. Identification of *Anoectochilus formosanus* and *Anoectochilus koshunensis* species with RAPD markers. *Planta Med* 1998; 64: 46–49
- 55 Zhang KYB, Leung HW, Yeung HW, Wong RNS. Differentiation of *Lycium barbarum* from its related *Lycium* species using random amplified polymorphic DNA. *Planta Med* 2001; 67: 379–381
- 56 Watanabe A, Araki S, Kobari S, Sudo H, Tsuchida T, Uno T, Kosaka N, Shimomura K, Yamazaki M, Saito K. *In vitro* propagation, restriction fragment length polymorphism, and random amplified polymorphic DNA analyses of *Angelica* plants. *Plant Cell Rep* 1998; 18: 187–192
- 57 Liang ZT, Qin MJ, Wang ZT, Huang Y, Wang NH. Identification of *Bupleurum* L. plants by RAPD technology. *Zhongcaoyào* 2002; 33: 1117–1119
- 58 Zhang M, Huang HR, Liao SM, Gao JY. Cluster analysis of *Dendrobium* by RAPD and design of specific primer for *Dendrobium candidum*. *Zhongguo Zhong Yao Za Zhi* 2001; 26: 442–447
- 59 Guo BL, Wu M, Si JP, Li JS, Xiao PG. Research on DNA molecular marker of *Magnolia officinalis* Rehd. et Wils. RAPD study on certified species. *Yao Xue Xue Bao* 2001; 36: 386–389
- 60 Huang L, Wang M, Zhou C, Li N, He X, Yang B. Problems and solutions in the use of RAPD to the identification of the Chinese drugs “xi-xin” (Herba Asari) and its substitutes. *Yao Xue Xue Bao* 1998; 33: 778–784
- 61 Lu C, Zhang W, Peng X, Gu G, Chen M, Tang Z. Development of randomly amplified polymorphic DNA-sequence characterized amplified region marker for identification of *Apocynum venetum* LINN. from *A. pictum* SCHRENK. *Biol Pharm Bull* 2010; 33: 522–526
- 62 Li Y, Ding WL. Genetic diversity assessment of *Trollius* accessions in China by RAPD markers. *Biochem Genet* 2010; 48: 34–43
- 63 Dnyaneshwar W, Preeti C, Kalpana J, Bhushan P. Development and application of RAPD-SCAR marker for identification of *Phyllanthus emblica* L. *Biol Pharm Bull* 2006; 29: 2313–2316
- 64 Zhang R, Zhang B, Ye H. Identification of the Chinese herbs of Indigoferae L. by RAPD analysis. *Hongguo Zhong Yao Za Zhi* 1997; 22: 72–73
- 65 Cheng KT, Chang HC, Su CH, Hsu FL. Identification of dried rhizomes of *Coptis* species using random amplified polymorphic DNA. *Bot Bull Acad Sin* 1997; 38: 241–244
- 66 Zhang YB, Ngan FN, Wang ZT, Wang J, But PPH, Shaw PC. Differentiation of *Codonopsis pilosula* using random amplified polymorphic DNA. *Planta Med* 1999; 65: 57–60
- 67 Cao H, But PPH, Shaw PC. Identification of Herba Taraxaci and its adulterants in Hong Kong market by DNA fingerprinting with random primed PCR. *Chin J Chin Mater Med* 1997; 22: 197–200
- 68 Cao H, But PPH, Shaw PC. Authentication of the Chinese drug “ku-didan” (Herba Elephantopi) and its substitutes using random-primed polymerase chain reaction (PCR). *Acta Pharm Sin* 1996; 31: 543–553
- 69 Cheng JL, Huang LQ, Shao AJ, Lin SF. RAPD analysis on different varieties of *Rehmannia glutinosa*. *Zhongguo Zhongyao Zazhi* 2002; 27: 505–508
- 70 Caetano-Anollés G, Bassam BJ, Gresshoff PM. DNA amplification fingerprinting: a strategy for genome analysis. *Plant Mol Biol Rep* 1991; 9: 294–307
- 71 Caetano-Anollés G, Bassam BJ, Gresshoff PM. DNA amplification using very short arbitrary oligonucleotide primers. *Biotechnology* 1991; 9: 553–557
- 72 Chawla HS. Introduction to Plant Biotechnology, 2nd edition. New Hampshire: Science Publishers; 2002
- 73 Zietkiewicz E, Rafalski A, Labuda D. Genome fingerprinting by simple sequence repeat (SSR)-anchored polymerase chain reaction amplification. *Genomics* 1994; 20: 176–183
- 74 Bornet B, Branchard M. Nonanchored inter simple sequence repeat (ISSR) markers: reproducible and specific tools for genome fingerprinting. *Plant Mol Biol Rep* 2001; 19: 209–215
- 75 Shen J, Ding XY, Ding G, Liu DY, Tang F, He J. Studies on population difference of *Dendrobium officinale* II establishment and optimization of the method of ISSR fingerprinting marker. *Zhongguo Zhong Yao Za Zhi* 2006; 31: 291–294
- 76 Shi HM, Wang J, Wang MY, Tu PF, Li XB. Identification of *Cistanche* species by chemical and inter-simple sequence repeat fingerprinting. *Biol Pharm Bull* 2009; 32: 142–146
- 77 Li K, Wu W, Zheng Y, Dai Y, Xiang L, Liao K. Genetic diversity of *Fritillaria* from Sichuan province based on ISSR. *Zhongguo Zhong Yao Za Zhi* 2009; 34: 2149–2154
- 78 Song Z, Li X, Wang H, Wang J. Genetic diversity and population structure of *Salvia miltiorrhiza* Bge in China revealed by ISSR and SRAP. *Genetica* 2010; 138: 241–249
- 79 Hu Y, Zhang Q, Xin H, Qin LP, Lu BR, Rahman K, Zheng H. Association between chemical and genetic variation of *Vitex rotundifolia* populations from different locations in China: its implication for quality control of medicinal plants. *Biomed Chromatogr* 2007; 21: 967–975
- 80 Kojoma M, Iida O, Makino Y, Sekita S, Satake M. DNA fingerprinting of *Cannabis sativa* using inter-simple sequence repeat (ISSR) amplification. *Planta Med* 2002; 68: 60–63
- 81 Xia T, Chen S, Chen S, Zhang D, Zhang D, Gao Q, Ge XJ. ISSR analysis of genetic diversity of the Qinghai-Tibet plateau endemic *Rhodiola chrysanthemifolia* (Crassulaceae). *Biochem Syst Ecol* 2007; 35: 209–214
- 82 Wu W, Zheng YL, Chen L, Wei YM, Yang RW, Yan ZH. Evaluation of genetic relationships in the genus *Houttuynia* Thunb. in China based on RAPD and ISSR markers. *Biochem Syst Ecol* 2005; 33: 1141–1157
- 83 Vos P, Hogers R, Bleeker M, Reijans M, van de Lee T, Hornes M, Frijters A, Pot J, Peleman J, Kuiper M. AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Res* 1995; 23: 4407–4414
- 84 Vos P, Kuiper M. AFLP Analysis. In: Caetano-Anollés G, Gresshoff PM, editors. DNA markers: protocols, applications, and overviews. New York: Wiley-Liss; 1997: 115–132
- 85 Blears M, De Grandis S, Lee H, Trevors J. Amplified fragment length polymorphism (AFLP): a review of the procedure and its applications. *J Indian Microbiol Biotechnol* 1998; 21: 99–114



- 86 Mueller UG, Wolfenbarger L. AFLP genotyping and fingerprinting. *Trends Ecol Evol* 1999; 14: 389–394
- 87 Choi YE, Ahn CH, Kim BB, Yoon ES. Development of species specific AFLP-derived SCAR marker for authentication of *Panax japonicus* C. A. MEYER. *Biol Pharm Bull* 2008; 31: 135–138
- 88 Ha WY, Shaw PC, Liu J, Yau FC, Wang J. Authentication of *Panax ginseng* and *Panax quinquefolius* using amplified fragment length polymorphism (AFLP) and directed amplification of minisatellite region DNA (DAMD). *J Agric Food Chem* 2002; 50: 1871–1875
- 89 Zerega NJC, Mori S, Lindqvist C, Zheng Q, Motley TJ. Using amplified fragment length polymorphisms (AFLP) to identify black cohosh (*Actaea racemosa*). *Econ Bot* 2002; 56: 154–164
- 90 Passinho-Soares H, Felix D, Kaplan MA, Margis-Pinheiro M, Margis R. Authentication of medicinal plant botanical identity by amplified fragmented length polymorphism dominant DNA marker: inferences from the *Plectranthus* genus. *Planta Med* 2006; 72: 929–931
- 91 Loh JP, Kiew R, Kee A, Gan LH, Gan YY. Amplified fragment length polymorphism (AFLP) provides molecular markers for the identification of *Caladium bicolor* cultivars. *Ann Bot (Lond)* 1999; 84: 155–161
- 92 Datwyler SL, Weiblen GD. Genetic variation in hemp and marijuana (*Cannabis sativa* L.) according to amplified fragment length polymorphisms. *J Forensic Sci* 2006; 51: 371–375
- 93 Qi JJ, Li XE, Song J, Eneji AE, Ma X. Genetic Relationships among *Rehmannia glutinosa* cultivars and varieties. *Planta Med* 2008; 74: 1846–1852
- 94 Weising K, Nybom H, Wolff K, Meyer W. DNA Fingerprinting in Plants and Fungi. Boca Raton: CRC Press; 1995
- 95 Soumaya R, Dakhlaoui-Dkhil S, Salem AOM, Zehdi-Azouzi S, Rhouma A, Marrakchi M, Trifi M. Genetic diversity and phylogenetic relationships in date-palms (*Phoenix dactylifera* L.) as assessed by random amplified microsatellite polymorphism markers (RAMPOs). *Sci Hortic* 2008; 117: 53–57
- 96 Chatti K, Saddoud O, Salhi Hannachi A, Mars M, Marrakchi M, Trifi M. Analysis of genetic diversity and relationships in a Tunisian Fig (*Ficus carica*) germplasm collection by random amplified microsatellite polymorphisms. *J Integr Plant Biol* 2007; 49: 386–391
- 97 Baldwin BG, Sanderson MJ, Porter JM, Wojciechowski MF, Campbell CS, Donoghue MJ. The ITS region of nuclear ribosomal DNA: a valuable source of evidence on angiosperm phylogeny. *Ann Mo Bot Gard* 1995; 82: 247–277
- 98 Alvarez I, Wendel JF. Ribosomal ITS sequences and plant phylogenetic inference. *Mol Phylogenet Evol* 2003; 29: 417–434
- 99 Chen S, Yao H, Han J, Liu C, Song J, Shi L, Zhu Y, Ma X, Gao T, Pang X, Luo K, Li Y, Li X, Jia X, Lin Y, Leon C. Validation of the ITS2 region as a novel DNA barcode for identifying medicinal plant species. *PLoS One* 2010; 5: 1–8
- 100 Chiou SJ, Yen JH, Fang CL, Chen HL, Lin TY. Authentication of medicinal herbs using PCR-Amplified ITS2 with specific primers. *Planta Med* 2007; 73: 1421–1426
- 101 Ngan F, Shaw P, But P, Wang J. Molecular authentication of *Panax* species. *Phytochemistry* 1999; 50: 787–791
- 102 Kim OT, Bang K, In DS, Lee JW, Kim YC, Shin YS, Hyun DY, Lee SS, Cha SW, Seong NS. Molecular authentication of ginseng cultivars by comparison of internal transcribed spacer and 5.8S rDNA sequences. *Plant Biotechnol Rep* 2007; 1: 163–167
- 103 Kelly LM. Phylogenetic relationships in *Asarum* (Aristolochiaceae) based on morphology and ITS sequences. *Am J Bot* 1998; 85: 1454–1467
- 104 Liu CS, Bai GB, Yan YN. Studies on the botanical sources and DNA molecular identification of Herba Asari based on ITS sequence. *Zhongguo Zhong Yao Za Zhi* 2005; 30: 329–332
- 105 Yamaji H, Fukuda T, Yokoyama J, Pak J-H, Zhou C, Yang CS, Kondo K, Morota T, Takeda S, Sasaki H, Maki M. Reticulate evolution and phylogeography in *Asarum* sect. *Asiasarum* (Aristolochiaceae) documented in internal transcribed spacer sequences (ITS) of nuclear ribosomal DNA. *Mol Phylogenet Evol* 2007; 44: 863–884
- 106 Dong TT, Ma XQ, Clarke C, Song ZH, Ji ZN, Lo CK, Tsim KW. Phylogeny of *Astragalus* in China: molecular evidence from the DNA sequences of 5S rRNA spacer, ITS, and 18S rRNA. *J Agric Food Chem* 2003; 51: 6709–6714
- 107 Yip PY, Kwan HS. Molecular identification of *Astragalus membranaceus* at the species and locality levels. *J Ethnopharmacol* 2006; 106: 222–229
- 108 Xu H, Wang Z, Ding X, Zhou K, Xu L. Differentiation of *Dendrobium* species used as “Huangcao Shihu” by rDNA ITS sequence analysis. *Planta Med* 2006; 72: 89–92
- 109 Ding X, Xu L, Wang Z, Zhou K, Xu H, Wang Y. Authentication of stems of *Dendrobium officinale* by rDNA ITS region sequences. *Planta Med* 2002; 68: 191–192
- 110 Lau DT, Shaw PC, Wang J, But PP. Authentication of medicinal *Dendrobium* species by the internal transcribed spacer of ribosomal DNA. *Planta Med* 2001; 67: 456–460
- 111 Zhang YB, Wang J, Wang ZT, But PP, Shaw PC. DNA microarray for identification of the herb of *Dendrobium* species from Chinese medicinal formulations. *Planta Med* 2003; 69: 1172–1174
- 112 Wang CZ, Li P, Ding JY, Jin GQ, Yuan CS. Identification of *Fritillaria pallidiflora* using diagnostic PCR and PCR-RFLP based on nuclear ribosomal DNA internal transcribed spacer sequences. *Planta Med* 2005; 71: 384–386
- 113 Yang ZY, Chao Z, Huo KK, Wu BY, Pan SL. Nuclear ribosomal DNA internal transcribed spacer 1 sequences of 4 *Leonurus* species. *Nan Fang Yi Ke Da Xue Xue Bao* 2006; 26: 1593–1595
- 114 Luo YM, Zhang WM, Ding XY, Shen J, Bao SL, Chu BH, Mao SG. SNP marker and allele-specific diagnostic PCR for authenticating herbs of *Perilla*. *Acta Pharm Sin* 2006; 41: 840–845
- 115 Lee SK, Li PT, Lau DT, Yung PP, Kong RY, Fong WF. Phylogeny of medicinal *Phyllanthus* species in China based on nuclear ITS and chloroplast atpB-rbcL sequences and multiplex PCR detection assay analysis. *Planta Med* 2006; 72: 721–726
- 116 Albach DC, Li HQ, Zhao N, Jensen SR. Molecular systematics and phytochemistry of *Rehmannia* (Scrophulariaceae). *Biochem Syst Ecol* 2007; 35: 293–300
- 117 Wang H, Wang Q. Analysis of rDNA ITS sequences of Radix et Rhizoma *Salviae miltiorrhizae* and plants of *Salvia* L. *Chin Tradit Herb Drugs* 2005; 36: 1381–1385
- 118 Xue CY, Li DZ, Lu JM, Yang JB, Liu JQ. Molecular authentication of the traditional Tibetan medicinal plant *Swertia mussoitii*. *Planta Med* 2006; 72: 1223–1226
- 119 Sahin FP, Yamashita H, Guo Y, Terasaka K, Kondo T, Yamamoto Y, Shimada H, Fujita M, Kawasaki T, Sakai E, Tanaka T, Goda Y, Mizukami H. DNA authentication of *Plantago* herb based on nucleotide sequences of 18S-28S rRNA internal transcribed spacer region. *Biol Pharm Bull* 2007; 30: 1265–1270
- 120 Yang ZY, Chao Z, Huo KK, Xie H, Tian ZP, Pan SL. ITS sequence analysis used for molecular identification of the *Bupleurum* species from northwestern China. *Phytomedicine* 2007; 14: 416–422
- 121 Xue HG, Zhou SD, He XJ, Yu Y. Molecular authentication of the traditional Chinese medicinal plant *Euphorbia pekinensis*. *Planta Med* 2006; 73: 91–93
- 122 Zhao KJ, Dong TT, Cui XM, Tu PF, Tsim KW. Genetic distinction of radix adenophorae from its adulterants by the DNA sequence of 5S-rRNA spacer domains. *Am J Chin Med* 2003; 31: 919–926
- 123 Carles M, Cheung MK, Moganti S, Dong TT, Tsim KW, Ip NY, Sucher NJ. A DNA microarray for the authentication of toxic traditional Chinese medicinal plants. *Planta Med* 2005; 71: 580–584
- 124 Zhao KJ, Dong TT, Tu PF, Song ZH, Lo CK, Tsim KW. Molecular genetic and chemical assessment of radix *Angelica* (Danggui) in China. *J Agric Food Chem* 2003; 51: 2576–2583
- 125 Dong TT, Ma XQ, Clarke C, Song ZH, Ji ZN, Lo CK, Tsim KW. Phylogeny of *Astragalus* in China: molecular evidence from the DNA sequences of 5S rRNA spacer, ITS, and 18S rRNA. *J Agric Food Chem* 2003; 51: 6709–6714
- 126 Ma XQ, Duan JA, Zhu DY, Dong TT, Tsim KW. Species identification of Radix *Astragali* (Huangqi) by DNA sequence of its 5S-rRNA spacer domain. *Phytochemistry* 2000; 54: 363–368
- 127 Xia Q, Zhao KJ, Huang ZG, Zhang P, Dong TT, Li SP, Tsim KW. Molecular genetic and chemical assessment of Rhizoma *Curcumae* in China. *J Agric Food Chem* 2005; 53: 6019–6026
- 128 Sun Y, Fung KP, Leung PC, Shi D, Shaw PC. Characterization of medicinal *Epimedium* species by 5S rRNA gene spacer sequencing. *Planta Med* 2004; 70: 287–288
- 129 Cai ZH, Li P, Dong TT, Tsim KW. Molecular diversity of 5S-rRNA spacer domain in *Fritillaria* species revealed by PCR analysis. *Planta Med* 1999; 65: 360–364
- 130 Ma XQ, Zhu DY, Li SP, Dong TT, Tsim KW. Authentic identification of Stigma *Croci* (stigma of *Crocus sativus*) from its adulterants by molecular genetic analysis. *Planta Med* 2001; 67: 183–186

- 131 Zhang M, Zhang DZ, Xu XH, Zhang T, Wang ZT. 5S rRNA gene spacer sequences from *Ligularia* medicinal plants and the identification of HPAs-containing species. *Chin J Nat Med* 2005; 3: 38–40
- 132 Sun Y, Shaw PC, Fung KP. Molecular authentication of *Radix Puerariae Lobatae* and *Radix Puerariae Thomsonii* by ITS and 5S rRNA spacer sequencing. *Biol Pharm Bull* 2007; 30: 173–175
- 133 Chen F, Chan HY, Wong KL, Wang J, Yu MT, But PP, Shaw PC. Authentication of *Saussurea lappa*, an endangered medicinal material, by ITS DNA and 5S rRNA sequencing. *Planta Med* 2008; 74: 889–892
- 134 Liu YP, Cao H, Wang XT. Application of gene technology in quality control of Chinese drugs (II) – Identification of Chinese yam (*Dioscorea polystachia rhizome*) using DNA sequencing. *Chin J Tradit Herb Drugs* 2001; 32: 113–117
- 135 Liu YP, Cao H, Wang XT. Application of gene technology in quality control of Chinese drugs (I) – identification of *Pinellia ternata* species from Yuncheng, Shandong using DNA sequencing. *Chin J Pharm Anal* 2001; 21: 423–426
- 136 Zhu S, Fushimi H, Cai S, Komatsu K. Phylogenetic relationship in the genus *Panax*: inferred from chloroplast trnK gene and nuclear 18S rRNA gene sequences. *Planta Med* 2003; 69: 647–653
- 137 Wang CZ, Li P, Ding JY, Jin GQ, Yuan CS. Identification of *Fritillaria pallidiflora* using diagnostic PCR and PCR-RFLP based on nuclear ribosomal DNA internal transcribed spacer sequences. *Planta Med* 2005; 71: 384–386
- 138 Lee SK, Li PT, Lau DT, Yung PP, Kong RY, Fong WF. Phylogeny of medicinal *Phyllanthus* species in China based on nuclear ITS and chloroplast atpB-rbcL sequences and multiplex PCR detection assay analysis. *Planta Med* 2006; 72: 721–726
- 139 Lee C, Wen J. Phylogeny of *Panax* using chloroplast trnC-trnD intergenic region and the utility of trnC-trnD in interspecific studies of plants. *Mol Phylogenet Evol* 2004; 31: 894–903
- 140 Albach DC, Li HQ, Zhao N, Jensen SR. Molecular systematics and phytochemistry of *Rehmannia* (Scrophulariaceae). *Biochem Syst Ecol* 2007; 35: 293–300
- 141 Yang M, Zhang D, Liu J, Zheng J. A molecular marker that is specific to medicinal rhubarb based on chloroplast trnL/trnF sequences. *Planta Med* 2001; 67: 784–786
- 142 Long C, Kakiuchi N, Takahashi A, Komatsu K, Cai S, Mikage M. Phylogenetic analysis of the DNA sequence of the non-coding region of nuclear ribosomal DNA and chloroplast of *Ephedra* plants in China. *Planta Med* 2004; 70: 1080–1084
- 143 Xue CY, Li DZ, Lu JM, Yang JB, Liu JQ. Molecular authentication of the traditional Tibetan medicinal plant *Swertia mussotii*. *Planta Med* 2006; 72: 1223–1226
- 144 Hosokawa K, Minami M, Nakamura I, Hishida A, Shibata T. The sequences of the plastid gene rpl16 and the rpl16-rpl14 spacer region allow discrimination among six species of *Scutellaria*. *J Ethnopharmacol* 2005; 99: 105–108
- 145 Hosokawa K, Hishida A, Nakamura I, Shibata T. The sequences of the spacer region between the atpF and atpA genes in the plastid genome allows discrimination among three varieties of medicinal *Angelica*. *Planta Med* 2006; 72: 570–571
- 146 Gong W, Fu C-X, Luo Y-P, Qiu Y-X. Molecular identification of *Sinopodophyllum hexandrum* and *Dysosma* species using cpDNA sequences and PCR-RFLP markers. *Planta Med* 2006; 72: 650–652
- 147 Zhao YP, Qiu YX, Gong W, Li JH, Fu CX. Authentication of *Actinidia macrosperma* using PCR-RFLP based on trnK sequences. *Bot Stud* 2007; 48: 239–242
- 148 Mizukami H, Okabe Y, Kohda H, Hiraoka N. Identification of the crude drug *Atractylodes Rhizome* (Byaku-jutsu) and *Atractylodes lancea Rhizome* (So-jutsu) using chloroplast trnK sequence as a molecular marker. *Biol Pharm Bull* 2000; 23: 589–594
- 149 Sasaki Y, Fushimi H, Cao H, Cai S-Q, Komatsu K. Sequence analysis of Chinese and Japanese *Curcuma* drugs on the 18S rRNA gene and trnK gene and the application of amplification-refractory mutation system analysis for their authentication. *Biol Pharm Bull* 2002; 25: 1593–1599
- 150 Luo JP, Cao H, Liu YP. DNA sequencing and molecular identification of Patchouli and its substitute wrinkled giant hyssop. *Yao Xue Xue Bao* 2002; 37: 739–742
- 151 Komatsu KS, Zhu S, Fushimi H, Qui TK, Cai S, Kadota S. Phylogenetic analysis based on 18S rRNA gene and matK gene sequences of *Panax vietnamensis* and five related species. *Planta Med* 2001; 67: 461–465
- 152 Zhao ZL, Leng CH, Wang ZT. Identification of *Dryopteris crassirhizoma* and the adulterant species based on cpDNA rbcL and translated amino acid sequences. *Planta Med* 2007; 73: 1230–1233
- 153 Kondo K, Terabayashi S, Okada M, Yuan C, He S. Phylogenetic relationship of medicinally important *Cnidium officinale* and Japanese *Apiaceae* based on rbcL sequences. *J Plant Res* 1996; 109: 21–27
- 154 Lin J, Zhou X, Gao S, Wu W, Liu X, Sun X, Tang K. Authentication of *Pinellia ternata* and its adulterants based on PCR with specific primers. *Planta Med* 2006; 72: 844–847
- 155 Maeda M, Uryu N, Murayama N, Ishii H, Ota M, Tsuji K, Inoko H. A simple and rapid method for HLA-DP genotyping by digestion of PCR-amplified DNA with allele specific restriction endonucleases. *Hum Immunol* 1990; 27: 111–121
- 156 Lum MR, Potter E, Dang T, Heber D, Hardy M, Hirsch AM. Identification of botanicals and potential contaminants through RFLP and sequencing. *Planta Med* 2005; 71: 841–846
- 157 Li X, Ding X, Chu B, Ding G, Gu S, Qian L, Wang Y, Zhou Q. Molecular authentication of *Alisma orientale* by PCR-RFLP and ARMS. *Planta Med* 2007; 73: 67–70
- 158 Watanabe A, Araki S, Kobari S, Sudo H, Tsuchida T, Uno T, Kosaka N, Shimomura K, Yamazaki M, Saito K. *In vitro* propagation, restriction fragment length polymorphism, and random amplified polymorphic DNA analyses of *Angelica* plants. *Plant Cell Rep* 1998; 18: 187–192
- 159 Gong W, Fu CX, Luo YP, Qiu YX. Molecular identification of *Sinopodophyllum hexandrum* and *Dysosma* species using cpDNA sequences and PCR-RFLP markers. *Planta Med* 2006; 72: 650–652
- 160 Guo Y, Tsuruga A, Yamaguchi S, Oba K, Iwai K, Sekita S, Mizukami H. Sequence analysis of chloroplast chlB gene of medicinal *Ephedra* species and its application to authentication of *Ephedra* herb. *Biol Pharm Bull* 2006; 29: 1207–1211
- 161 Wang CZ, Li P, Ding JY, Jin GQ, Yuan CS. Identification of *Fritillaria pallidiflora* using diagnostic PCR and PCR-RFLP based on nuclear ribosomal DNA internal transcribed spacer sequences. *Planta Med* 2005; 71: 384–386
- 162 Wang CZ, Li P, Ding JY, Peng X, Yuan CS. Simultaneous identification of *Bulbus Fritillariae cirrhosae* using PCR-RFLP analysis. *Phytomedicine* 2007; 14: 628–632
- 163 Lee JH, Lee JW, Sung JS, Bang KH, Moon SG. Molecular authentication of 21 Korean *Artemisia* species (Compositae) by polymerase chain reaction-restriction fragment length polymorphism based on trnL-F region of chloroplast DNA. *Biol Pharm Bull* 2009; 32: 1912–1916
- 164 Diao Y, Lin XM, Liao CL, Tang CZ, Chen ZJ, Hu ZL. Authentication of *Panax ginseng* from its adulterants by PCR-RFLP and ARMS. *Planta Med* 2009; 75: 557–560
- 165 Do KR, Hwang WJ, Lyu YS, An NH, Kim HM. Molecular authentication of *Panax ginseng* species by RAPD analysis and PCR-RFLP. *Biol Pharm Bull* 2001; 24: 872–875
- 166 Lu KT, Lee HC, Liu FS, Lo CF, Lin JH. Identification of ginseng radix in Chinese medicine preparations by nested PCR-DNA sequencing method and nested PCR-restriction fragment length polymorphism. *J Food Drug Anal* 2010; 18: 58–63
- 167 Um JY, Chung HS, Kim MS, Na HJ, Kwon HJ, Kim JJ, Lee KM, Lee SJ, Lim JP, Do KR, Hwang WJ, Lyu YS, An NH, Kim HM. Molecular authentication of *Panax ginseng* species by RAPD analysis and PCR-RFLP. *Biol Pharm Bull* 2001; 24: 872–875
- 168 Zhao YP, Qiu YX, Gong W, Li JH, Fu CX. Authentication of *Actinidia macrosperma* using PCR-RFLP based on trnK sequences. *Bot Stud* 2007; 48: 239–242
- 169 Mizukami H, Okabe Y, Kohda H, Hiraoka N. Identification of the crude drug *Atractylodes rhizome* (Byaku-jutsu) and *Atractylodes lancea rhizome* (So-jutsu) using chloroplast TrnK sequence as a molecular marker. *Biol Pharm Bull* 2000; 23: 589–594
- 170 Mizukami H, Ohbayashi K, Umetsu K, Hiraoka N. Restriction fragment length polymorphism of medicinal plants and crude drugs. II. Analysis of *Glehnia littoralis* of different geographical origin. *Biol Pharm Bull* 1993; 16: 611–612
- 171 Lu KT, Lee HC, Liu FS, Lo CF, Lin JH. Discriminating *Astragalus Radix* from *Hedysarum Radix* in Chinese Medicine Preparations Using Nested PCR and DNA Sequencing Methods. *J Food Drug Anal* 2009; 17: 380–385
- 172 Zhang T, Xu LS, Wang ZT, Zhou KY, Zhang N, Shi YF. Molecular identification of medicinal plants: *Dendrobium chrysanthum*, *Dendrobium fimbriatum* and their morphologically allied species by PCR-RFLP analyses. *Yao Xue Xue Bao* 2005; 40: 728–733

- 173 Mizukami H, Ohbayashi K, Kitamura Y, Ikenaga T. Restriction fragment length polymorphisms (RFLPs) of medicinal plants and crude drugs. I. RFLP probes allow clear identification of *Duboisia* interspecific hybrid genotypes in both fresh and dried tissues. *Biol Pharm Bull* 1993; 16: 388–390
- 174 Fu RZ, Wang J, Zhang YB, Wang ZT, But PP, Li N, Shaw PC. Differentiation of medicinal *Codonopsis* species from adulterants by polymerase chain reaction-restriction fragment length polymorphism. *Planta Med* 1999; 65: 648–650
- 175 Desmarais E, Lanneluc I, Lagnel J. Direct amplification of length polymorphisms (DALP), or how to get and characterize new genetic markers in many species. *Nucleic Acids Res* 1998; 26: 1458–1465
- 176 Ma YS, Yu H, Li YY, Yan H, Cheng X. A study of genetic structure of *Stephania yunnanensis* (Menispermaceae) by DALP. *Biochem Genet* 2008; 46: 227–240
- 177 Ha WY, Yau FC, But PP, Wang J, Shaw PC. Direct amplification of length polymorphism analysis differentiates *Panax ginseng* from *P. quinquefolius*. *Planta Med* 2001; 67: 587–589
- 178 Newton CR, Graham A, Heptinstall LE, Powell SJ, Summers C, Kalsheker N, Smith JC, Markham AF. Analysis of any point mutation in DNA. The amplification refractory mutation system (ARMS). *Nucleic Acids Res* 1989; 17: 2503–2516
- 179 Li X, Ding X, Chu B, Ding G, Gu S, Qian L, Wang Y, Zhou Q. Molecular authentication of *Alisma orientale* by PCR-RFLP and ARMS. *Planta Med* 2007; 73: 67–70
- 180 Zhu S, Fushimi H, Cai S, Komatsu K. Species identification from Ginseng drugs by multiplex amplification refractory mutation system (MARMS). *Planta Med* 2004; 70: 189–192
- 181 Diao Y, Lin XM, Liao CL, Tang CZ, Chen ZJ, Hu ZL. Authentication of *Panax ginseng* from its Adulterants by PCR-RFLP and ARMS. *Planta Med* 2009; 75: 557–560
- 182 Yang DY, Fushimi H, Cai SQ, Komatsu K. Polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) and amplification refractory mutation system (ARMS) analyses of medicinally used *Rheum* species and their application for identification of *Rhei Rhizoma*. *Biol Pharm Bull* 2004; 27: 661–669
- 183 Ding G, Zhang D, Feng Z, Fan W, Ding X, Li X. SNP, ARMS and SSH authentication of medicinal *Dendrobium officinale* Kimura et Migo and application for identification of Fengdou drugs. *Biol Pharm Bull* 2008; 31: 553–557
- 184 Qian L, Ding G, Zhou Q, Feng ZY, Ding XY, Gu S, Wang Y, Li XX, Chu BH. Molecular authentication of *Dendrobium loddigesii* Rolfe by amplification refractory mutation system (ARMS). *Planta Med* 2008; 74: 470–473
- 185 Sasaki Y, Fushimi H, Cao H, Cai SQ, Komatsu K. Sequence analysis of Chinese and Japanese *Curcuma* drugs on the 18S rRNA gene and trnK gene and the application of amplification-refractory mutation system analysis for their authentication. *Biol Pharm Bull* 2002; 25: 1593–1599
- 186 Semagn K, Bjørnstad A, Ndjiondjop MN. An overview of molecular marker methods for plants. *African J Biotechnol* 2006; 5: 2540–2568
- 187 McDermott JM, Brandle U, Dutly F, Haemmerli UA, Keller S, Muller KE, Wolf MS. Genetic variation in powdery mildew of barley: development of RAPD, SCAR and VNTR markers. *Phytopathology* 1994; 4: 1316–1321
- 188 Albani MC, Battey NH, Wilkinson MJ. The development of ISSR derived SCAR markers around the seasonal flowering locus (SFL) in *Fragaria vesca*. *Theor Appl Genet* 2004; 109: 571–579
- 189 Paran I, Kesseli R, Michelmore R. Identification of restriction fragment length polymorphism and random amplified polymorphic DNA markers linked to downy mildew resistance genes in lettuce, using near isogenic lines. *Genome* 1991; 34: 1021–1027
- 190 Wang J, Ha WY, Ngan FN, But PP, Shaw PC. Application of sequence characterized amplified region (SCAR) analysis to authenticate *Panax* species and their adulterants. *Planta Med* 2001; 67: 781–783
- 191 Choi YE, Ahn CH, Kim BB, Yoon ES. Development of species specific AFLP-derived SCAR marker for authentication of *Panax japonicus* C. A. MEYER. *Biol Pharm Bull* 2008; 31: 135–138
- 192 Lee MY, Doh EJ, Park CH, Kim YH, Kim ES, Ko BS, Oh SE. Development of SCAR marker for discrimination of *Artemisia princeps* and *A. argyi* from other *Artemisia* herbs. *Biol Pharm Bull* 2006; 29: 629–633
- 193 Dnyaneshwar W, Preeti C, Kalpana J, Bhushan P. Development and application of RAPD-SCAR marker for identification of *Phyllanthus emblica* LINN. *Biol Pharm Bull* 2006; 29: 2313–2316
- 194 Theerakulpisut P, Kanawapee N, Maensiri D, Bunnag S, Chantaranonthai P. Development of species-specific SCAR markers for identification of three medicinal species of *Phyllanthus*. *J Syst Evol* 2008; 46: 614–621
- 195 Devaiah KM, Venkatasubramanian P. Development of SCAR marker for authentication of *Pueraria tuberosa* (Roxb. ex. Willd.) DC. *Curr Sci* 2008; 94: 1306–1308
- 196 Ye Q, Qiu YX, Quo YQ, Chen JX, Yang SZ, Zhao MS, Fu CX. Species-specific SCAR markers for authentication of *Sinocalycanthus chinensis*. *J Zhejiang Univ Sci B* 2006; 7: 868–872
- 197 Devaiah KM, Venkatasubramanian P. Genetic characterization and authentication of *Embelia ribes* using RAPD-PCR and SCAR marker. *Planta Med* 2008; 74: 194–196
- 198 Sze SCW, Song JX, Wong RNS, Feng YB, Ng TB, Tong Y, Zhang KYB. Application of SCAR (sequence characterized amplified region) analysis to authenticate *Lycium barbarum* (wolfberry) and its adulterants. *Biotechnol Appl Biochem* 2008; 51: 15–21
- 199 Litt M, Luty JA. A hypervariable microsatellite revealed by in vitro amplification of a dinucleotide repeat within the cardiac muscle actin gene. *Am J Hum Genet* 1989; 44: 397–401
- 200 Tautz D. Hypervariability of simple sequences as a general source of polymorphic DNA markers. *Nucleic Acids Res* 1989; 17: 6463–6471
- 201 Gupta PK, Balyan HS, Sharma PC, Ramesh B. Microsatellites in plants: a new class of molecular markers. *Curr Sci* 1996; 70: 45–54
- 202 Kim J, Jo BH, Lee KL, Yoon ES, Ryu GH, Chung KW. Identification of new microsatellite markers in *Panax ginseng*. *Mol Cells* 2007; 24: 60–68
- 203 Jo BH, Suh DS, Cho EM, Kim J, Ryu GH, Chung KW. Characterization of polymorphic microsatellite loci in cultivated and wild *Panax ginseng*. *Genes Genomics* 2009; 31: 119–127
- 204 Kim J, Chung KW. Isolation of new microsatellite-containing sequences in *Acanthopanax senticosus*. *J Plant Biol* 2007; 50: 557–561
- 205 Fan WJ, Luo YM, Li XX, Gu S, Xie ML, He J, Cai WT, Ding XY. Development of microsatellite markers in *Dendrobium fimbriatum* Hook, an endangered Chinese endemic herb. *Mol Ecol Res* 2009; 9: 373–375
- 206 Kumar J, Verma V, Shahi AK, Qazi GN, Balyan HS. Development of simple sequence repeat markers in *Cymbopogon* species. *Planta Med* 2007; 73: 262–266
- 207 Chun S, Jian-He W, Shi-Lin C, Huai-Qiong C, Cheng-Min Y. Development of genomic SSR and potential EST-SSR markers in *Bupleurum chinense* DC. *African J Biotechnol* 2009; 8: 6233–6240
- 208 Boqian Y, Jing W, Guopei C, Ting W. Isolation and characterization of polymorphic microsatellite loci in a traditional Chinese medicinal plant, *Schisandra sphenanthera*. *Conserv Genet* 2009; 10: 615–617
- 209 Vos P, Hogers R, Bleeker M, Reijers M, Van de Lee T, Hornes M, Frijters A, Pot J, Peleman J, Kuiper M, Zabeau M. AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Res* 1995; 23: 4407–4441
- 210 Paglia GE, Olivieri AM, Morgante M. Towards second-generation STS (sequence-tagged sites) linkage maps in conifers, a genetic map of Norway spruce (*Picea abies* K.). *Mol Gen Genet* 1998; 258: 466–478
- 211 Karp A, Edwards KJ. DNA markers, a global overview. In: Caetano-Anollds G, Gresshoff PM, editors. *DNA Markers, Protocols, Applications, and Overviews*. New York: Wiley-Liss; 1997
- 212 Molina C, Kahl G. Genomics of two banana pathogens, genetic diversity, diagnostics, and phylogeny of *Mycosphaerella fijiensis* and *M. musicola*. In: Jain SM, editor. *Banana Improvement, Cellular and Molecular Biology, and Induced Mutations*. Vienna: FAO/IAEA; 2002
- 213 Winter E, Pfaff T, Udupa SM, Hiittel B, Sharma PC, Sahi S, Arreguin-Espinosa R, Weigand F, Muehlbauer FJ, Kah G. Characterization and mapping of sequence-tagged microsatellite sites in the chickpea (*Cicer arietinum* L.) genome. *Mol Gen Genet* 1999; 262: 90–101
- 214 Witsenboer H, Vogel J, Michelmore RW. Identification, genetic localization, and allelic diversity of selectively amplified polymorphic loci in lettuce and wild relatives (*Lactuca* spp.). *Genome* 1997; 40: 923–936
- 215 Sarwat M, Das S, Srivastava PS. Analysis of genetic diversity through AFLP, SAMPL, ISSR and RAPD markers in *Tribulus terrestris*, a medicinal herb. *Plant Cell Rep* 2008; 27: 519–528
- 216 Heath DD, Iwama GK, Devlin RH. PCR primed with VNTR core sequences yields species specific patterns and hypervariable probes. *Nucleic Acids Res* 1993; 21: 5782–5785
- 217 Somers DJ, Demmon G. Identification of repetitive, genome-specific probes in crucifer oilseed species. *Genome* 2002; 45: 485–492
- 218 Silva LM, Montes de Oca H, Diniz CR, Fortes-Dias CL. Fingerprinting of cell lines by directed amplification of minisatellite-region DNA (DAMD). *Braz J Med Biol Res* 2001; 34: 1405–1410

- 219 Zhou Z, Bebeli PJ, Somers DJ, Gustafson JP. Direct amplification of minisatellite-region DNA with VNTR core sequences in the genus *Oryza*. *Theor Appl Genet* 1997; 95: 942–949
- 220 Ha WY, Shaw PC, Liu J, Yau F, Wang J. Authentication of *Panax ginseng* and *Panax quinquefolius* using amplified fragment length polymorphism (AFLP) and directed amplification of minisatellite region DNA (DAMD). *J Agric Food Chem* 2002; 50: 1871–1875
- 221 Ince AG, Karaca M, Onus AN. Development and utilization of diagnostic DAMD-PCR markers for *Capsicum* accessions. *Genet Resour Crop Evol* 2009; 56: 211–221
- 222 Karaca M, Ince AG, Tugrul S, Turgut K, Onus AN. PCR-RFLP and DAMD-PCR genotyping for *Salvia* species. *J Sci Food Agric* 2008; 88: 2508–2516
- 223 Bhattacharya E, Dandin SB, Ranade SA. Single primer amplification reaction methods reveal exotic and indigenous mulberry varieties are similarly diverse. *J Biosci* 2005; 30: 669–677
- 224 Chavan P, Joshi K, Patwardhan B. DNA microarrays in herbal drug research. *Evid Based Complement Alternat Med* 2006; 3: 447–457
- 225 Gebauer M. Microarray applications: emerging technologies and perspectives. *Drug Discov Today* 2004; 9: 915–917
- 226 Debouck C, Goodfellow PN. DNA microarrays in drug discovery and development. *Nat Genet* 1999; 21: 48–50
- 227 Trau D, Lee TM, Lao AI, Lenigk R, Hsing IM, Ip NY. Genotyping on a complementary metal oxide semiconductor silicon polymerase chain reaction chip with integrated DNA microarray. *Anal Chem* 2002; 74: 3168–3173
- 228 Tsoi PY, Wu HS, Wong MS, Chen SL, Fong WF, Xiao PG, Yang MS. Genotyping and species identification of *Fritillaria* by DNA chip technology. *Acta Pharm Sin* 2003; 4: 185–190
- 229 Li T, Wang J, Lu Z. Accurate identification of closely related *Dendrobium* species with multiple species-specific gDNA probes. *J Biochem Biophys Methods* 2005; 62: 111–123
- 230 Zhang YB, Wang J, Wang ZT, But PP, Shaw PC. DNA microarray for identification of the herb of *Dendrobium* species from Chinese medicinal formulations. *Planta Med* 2003; 69: 1172–1174
- 231 Lin WY, Chen LR, Lin TY. Rapid authentication of *Bupleurum* species using an array of immobilized sequence-specific oligonucleotide probes. *Planta Med* 2008; 74: 464–469
- 232 Qin J, Leung FC, Fung Y, Zhu D, Lin B. Rapid authentication of ginseng species using microchip electrophoresis with laser-induced fluorescence detection. *Anal Bioanal Chem* 2005; 381: 812–819
- 233 Carles M, Cheung MK, Moganti S, Dong TT, Tsim KW, Ip NY, Sucher NJ. A DNA microarray for the authentication of toxic traditional Chinese medicinal plants. *Planta Med* 2005; 71: 580–584
- 234 Pastinen T, Partanen J, Syvanen AC. Multiplex, fluorescent, solid-phase minisequencing for efficient screening of DNA sequence variation. *Clin Chem* 1996; 42: 1391–1397
- 235 Cai H, White PS, Torney D, Deshpande A, Wang Z, Keller RA, Marrone B, Nolan JP. Flow cytometry-based minisequencing: a new platform for high-throughput single-nucleotide polymorphism scoring. *Genomics* 2000; 66: 135–143
- 236 Pastinen T, Raitio M, Lindroos K, Tainola P, Peltonen L, Syvänen AC. A system for specific, high-throughput genotyping by allele-specific primer extension on microarrays. *Genome Res* 2000; 10: 1031–1042
- 237 Lowe CR. Nanobiotechnology: the fabrication and applications of chemical and biological nanostructures. *Curr Opin Struct Biol* 2000; 10: 428–434
- 238 Fortina P, Kricka LJ, Surrey S, Grodzinski P. Nanobiotechnology: the promise and reality of new approaches to molecular recognition. *Trends Biotechnol* 2005; 23: 168–173
- 239 Pirrung MC, Connors RV, Odenbaugh AL, Montague-Smith MP, Walcott NG, Tollett JJ. The arrayed primer extension method for DNA microchip analysis. *Molecular computation of satisfaction problems*. *J Am Chem Soc* 2000; 122: 1873–1882
- 240 Kurg A, Tönisson N, Georgiou I, Shumaker J, Tollett J, Metspalu A. Arrayed primer extension: solid-phase four-color DNA resequencing and mutation detection technology. *Genet Test* 2000; 4: 1–7
- 241 Grossman PD, Bloch W, Brinson E, Chang CC, Eggerding FA, Fung S, Iovannisci DM, Woo S, Winn-Deen ES, Iovannisci DA. High-density multiplex detection of nucleic acid sequences: oligonucleotide ligation assay and sequence-coded separation. *Nucleic Acids Res* 1994; 22: 4527–4534
- 242 Cimino MT. Successful isolation and PCR amplification of DNA from National Institute of Standards and Technology herbal dietary supplement standard reference material powders and extracts. *Planta Med* 2010; 76: 495–497