

Giant atypical lymphoid hyperplasia of the colon



Fig. 1 Double-contrast barium study disclosed a mass of approximately 2 × 3 cm (arrow) in the ascending colon.




Fig. 2 Abdominal computed tomography (CT) showed one mass (arrow) in the ascending colon.




Fig. 3 Colonoscopy revealed one polypoid mass (arrow) in the ascending colon.




Fig. 4 Histopathological examination showed proliferation and aggregation of many atypical lymphoid cells, consistent with atypical lymphoid hyperplasia (hematoxylin and eosin [H&E]; × 400).

A 57-year-old woman with a history of acute myelogenous leukemia had received chemotherapy. Thereafter, she was relatively stable with complete remission. After 8 years, the woman visited hospital for evaluation of intermittent abdominal fullness. Both physical examinations and laboratory studies were unremarkable. Double-contrast barium study and abdominal computed tomography disclosed a single mass of about 2 × 3 cm in the ascending colon (► **Figs. 1** and **2**). Colonoscopy confirmed a single polypoid mass in the ascending colon (► **Fig. 3**).

Endoscopic mucosal resection was accordingly performed. Histopathological examination showed proliferation and aggregation of many atypical lymphoid cells, consistent with atypical lymphoid hyperplasia (► **Fig. 4**).

The woman underwent colonoscopy 3 months later, which revealed that the mucosa of the lesion site had healed well. Atypical lymphoid hyperplasia is a condition usually related to an underlying immune dysregulation, or a reactive change to various inciting antigens or irritating stimuli [1–4]. To our knowledge, this is

the first reported case of polyp-like atypical lymphoid hyperplasia in the colon; this may be considered a pre-lymphomatous state, and close follow-up is recommended [4,5].

Competing interests: None

Endoscopy_UCTN_Code_CCL_1AD_2AC

W.-C. Huang¹, T.-Y. Hsieh¹, H.-C. Chu¹,
T.-K. Chao², T.-Y. Huang¹

¹ Division of Gastroenterology, Department of Internal Medicine, Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan

² Department of Pathology, Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan

References

- 1 Crowson AN, Magro CM. Antidepressant therapy. A possible cause of atypical cutaneous lymphoid hyperplasia. *Arch Dermatol* 1995; 131: 925–929
- 2 Tanahashi T, Tatsumi Y, Sawai N et al. Regression of atypical lymphoid hyperplasia after eradication of *Helicobacter pylori*. *J Gastroenterol* 1997; 32: 543–547
- 3 Viraben R, Lamant L, Brousset P. Losartan-associated atypical cutaneous lymphoid hyperplasia. *Lancet* 1997; 350: 1366
- 4 Krauss E, Konturek P, Maiss J et al. Clinical significance of lymphoid hyperplasia of the lower gastrointestinal tract. *Endoscopy* 2010; 42: 334–337
- 5 Takano Y, Kato Y, Sugano H. Histopathological and immunohistochemical study of atypical lymphoid hyperplasia and benign lymphoid hyperplasia of the stomach. *Jpn J Cancer Res* 1992; 83: 288–293

Bibliography

DOI 10.1055/s-0030-1255691

Endoscopy 2010; 42: E195–E196

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

T.-Y. Huang, MD, PhD

Division of Gastroenterology
Department of Internal Medicine
Tri-Service General Hospital
National Defense Medical Center
325 Cheng-Kung Road, Section 2
Neihu
Taipei 114
Taiwan
Fax: +866-2-8792-7138
teinyu.chun@msa.hinet.net