


M. NEUMANN, S. FÜLDNER, B. KÖNIG, K. ZEITLER* (UNIVERSITÄT REGENSBURG, GERMANY)

Metal-Free, Cooperative Asymmetric Organophotoredox Catalysis with Visible Light

Angew. Chem. Int. Ed. **2010**, DOI: 10.1002/anie.201002992.

Organophotoredox Catalysis



Significance: Classic xanthenes were demonstrated to be efficient photoredox catalysts for the organocatalytic asymmetric intermolecular α -alkylation of aldehydes. Under visible light irradiation eosin Y (**2**) gave comparable results to precious metal catalysts.

Comment: In 2008, Nicewicz and MacMillan reported an asymmetric intermolecular α -alkylation of aldehydes via a combination of photoredox catalysis and organocatalysis (*Science* **2008**, 322, 77). Now, Zeitler and co-workers demonstrated that precious metal complexes can be replaced by xanthenes as photoredox catalysts rendering the process purely organic. The extension of highly versatile photoredox catalysis to classic organic dyes is likely to find utility in a variety of other transformations.

SYNFACTS Contributors: Benjamin List, Ilija Čorić
Synfacts 2010, 12, 1419-1419 Published online: 22.11.2010
DOI: 10.1055/s-0030-1258925; Reg-No.: B12010SF

2010 © THIEME STUTTGART • NEW YORK

Category

Organo- and Biocatalysis

Key words

photoredox catalysis

α -alkylation

xanthenes dyes

SYNFACTS
of the month