


K. TANINO,* M. TAKAHASHI, Y. TOMATA, H. TOKURA, T. UEHARA, T. NARABU, M. MIYASHITA* (HOKKAIDO UNIVERSITY, SAPPORO; NATIONAL AGRICULTURE RESEARCH CENTER FOR HOKKAIDO REGION, SAPPORO AND KOGAKUIN UNIVERSITY, HACHIOJI, JAPAN)

Total Synthesis of Solanoeclepin A

Nat. Chem. 2011, 3, 484-488.

Synthesis of Solanoeclepin A



Significance: Solanoeclepin A, a complex triterpenoid, was isolated in 1986 by Mulder and co-workers and is a hatch stimulus for the potato cyst nematode, which inflicts considerable damage to crop plants. Cyst nematodes are notoriously hard to exterminate and as such attention has been drawn to study of such stimuli in order to eradicate this problem.

Comment: Key steps in the synthesis include several rearrangements and a Charette-modified Simmons–Smith cyclopropanation to complete the key DEFG fragment **L**. Further elaboration via an intramolecular Diels–Alder reaction completed the ABCDEFG scaffold (**N**). The first total synthesis of solanoeclepin A was completed in further 16 steps.

SYNFACTS Contributors: Steven V. Ley, James R. Frost
Synfacts 2011, 8, 0809-0809 Published online: 20.07.2011
DOI: 10.1055/s-0030-1260753; Reg-No.: N03911SF

2011 © THIEME STUTTGART • NEW YORK

Category

Synthesis of Natural Products and Potential Drugs

Key words

solanoeclepin A

Meinwald rearrangement

stereoselective epoxidation

Charette-modified Simmons–Smith cyclopropanation

intramolecular Diels–Alder reaction

SYNFACTS
of the month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.

809