

Cat scratch esophagus

Cat scratch lesion of the gastrointestinal tract is a rare entity. It was first reported in the colon, in 2007 [1]. A few case reports of cat scratch colon were published in the English literature following the initial description of this entity [2–4]. Recently, a case was described of linear esophageal tears in a man, similar to the colonic cat scratch lesion [5]. We report the second case of cat scratch esophagus and describe its natural clinical course.




Fig. 1 First endoscopy showing cat scratch lesion on the esophageal wall.

A 45-year-old woman was admitted to our gastroenterology clinic with dyspepsia. She had undergone an upper endoscopy in another hospital 1 week before admission to our clinic. We performed an upper endoscopy in July 2010, and it revealed tears along the long axis of the esophagus. These superficial longitudinal lesions were evenly distributed around the circumference, resembling a cartwheel (● Fig. 1). Mucosal biopsies from the esophageal mucosa showed detachment of the squamous epithelial layer with mild lymphoplasmacytic infiltration (● Fig. 2). A control upper endoscopy of the patient in August 2010 demonstrated the complete healing of the esophageal lesions without any treatment. Cat scratch colon is characterized by bright, superficial linear erythematous marks and parallel corkscrew lesions without significant hemorrhage. Colonic cat scratch lesions are usually localized in the right colon. The lesion has been observed mainly in normal patients with normal histology [1]. The single reported case of cat scratch lesion involving the esophagus was similar to the colonic counterparts [5]. The etiopathogenesis of this lesion is unclear. However, it has been suggested that the lesion is caused by barotrauma from air insufflation rather than by direct colonoscopy trauma [4]. In our case, the patient's previous upper endoscopy probably gave rise to this morphology. We believe that the ileocecal valve in the right colon and the lower esophageal sphincter

in the esophagus may act as barriers and increase the intraluminal pressure if they do not open properly at the right time during air insufflation.

Cat scratch morphology is accepted as an innocent finding due to its normal histology [2]. However, there were insufficient data regarding the natural clinical course of the lesion. Our case demonstrates that cat scratch esophagus is a self-healing lesion and confirms that it is an innocent lesion.

Endoscopy_UCTN_Code_CCL_1AB_2AC_3AH

Competing interests: None

K. Deniz¹, M. Başkol², Ş. Gürsoy², K. Güven²

¹ Department of Pathology, Erciyes University, Medical School, Kayseri, Turkey

² Department of Gastroenterology, Erciyes University, Medical School, Kayseri, Turkey

References

- 1 McDonnell WM, Loura F, Pointon MJ et al. Cat scratch colon. *Endoscopy* 2007; 39: 459–461
- 2 Katsinelos P, Kountouras J, Chatzimavroudis G et al. "Cat scratch" colon: an impressive but innocent endoscopic finding. *Endoscopy* 2007; 39: 1026
- 3 Tominaga K, Shigiyama F, Ito S et al. Emergence of "cat scratch colon" during a colonoscopy. *Endoscopy* 2008; 40: 353
- 4 Baudet JS, Diaz-Bethencourt D, Arguinarena X et al. Cat scratch colon is caused by barotrauma secondary to insufflation during colonoscopy. *Endoscopy* 2008; 40: 878
- 5 Velayos B, Fernandez L, del-Olmo L et al. Cat scratch esophagus: a new entity to be described. *Rev Esp Enferm Dig* 2011; 103: 427–428

Bibliography

DOI <http://dx.doi.org/10.1055/s-0031-1291522>
Endoscopy 2012; 44: E44
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

K. Deniz, MD
 Erciyes University
 Faculty of Medicine
 Department of Pathology
 38039 Kayseri
 Turkey
 Fax: +90-352-4375273
 drkdeniz@yahoo.com




Fig. 2 Histology of the esophageal biopsy, showing traumatic breaks within the squamous epithelial layers. Mild lymphocytic inflammation was observed (Hematoxylin and eosin, ×100.)