

ERRATA

Immune-Mediated Encephalopathies with an Emphasis on Paraneoplastic Encephalopathies

Amy A. Pruitt, M.D.¹

The publisher regrets an error in the dosing units of measure in the above article in *Seminars in Neurology*, Volume 31, Number 2, 2011, p. 165. The correct dosing is indicated below.

Rituximab dosage should be 375 mg/m² (meter squared, not millimeter squared)

Cytosan dosage should be 750 mg/m² (meter squared, not millimeter squared)

¹Department of Neurology, University of Pennsylvania, Philadelphia Pennsylvania.

Address for correspondence and reprint requests: Amy A. Pruitt, M.D., Department of Neurology, University of Pennsylvania, 3400 Spruce Street, Philadelphia, PA 19104 (e-mail: pruitt@mail.med.upenn.edu).

Emerging and Controversial Issues in Neurology: Infectious

Diseases; Guest Editor, Tracey A. Cho, M.D.

Semin Neurol 2011;31:356. Copyright © 2011 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA. Tel: +1(212) 584-4662.

DOI: <http://dx.doi.org/10.1055/s-0031-1291959>.

ISSN 0271-8235.