

Endoscopic snare polypectomy of a pedunculated adenocarcinoma of the duodenal bulb arising from a hyperplastic polyp

Primary non-ampullary adenocarcinoma of the duodenum is a rare occurrence, arising mainly from adenomatous polyps [1]. We report the first case of a pedunculated adenocarcinoma of the duodenal bulb, arising from a hyperplastic polyp, treated with endoscopic snare polypectomy.

A 76-year-old man underwent esophago-gastroduodenoscopy (OGD) for asymptomatic iron-deficiency anemia. A 2-cm pedunculated polyp in the duodenal bulb (Fig. 1) and a small antral polyp were noted. Snare polypectomy of the pedunculated polyp was performed. Histopathology revealed a polyp sharing microscopic similarities to hyperplastic gastric polyps with hyperplastic irregular crypts, basal cystic change, and a focally retained villiform surface. Reactive eosinophilic syncytial cells with large vesicular nuclei

were present along with foci of severe dysplasia and possible early invasion (Fig. 2 and Fig. 3). Antral biopsies revealed *Helicobacter pylori* gastritis and a hyperplastic polyp with low grade dysplasia. A colonoscopy, staging computed tomography (CT) scan, and small-bowel barium studies were normal.

Following treatment of the patient with triple therapy and proton pump inhibitors, a repeat OGD at 3 months confirmed *H. pylori* eradication but showed high grade dysplasia of the stomach body. Subsequent OGDs at 6 months and 1 year showed downstaging to low grade dysplasia and intestinal metaplasia respectively. The patient is due for a repeat OGD in 1 year.

Incidental non-ampullary duodenal polyps are frequently encountered at endoscopy, and are mostly of an adeno-

matous nature, especially if ≥ 10 mm in size [2]. While adenomas are premalignant for duodenal adenocarcinoma [1], hyperplastic polyps in the stomach and colon have also rarely been shown to possess malignant potential [3]. Progression of duodenal hyperplastic polyps though remains uncertain [4]. This case is the first to report malignant transformation from a hyperplastic polyp occurring in the duodenum, emphasizing the importance of complete excision of such polyps. Endoscopic snare polypectomy is safe when polyps are pedunculated [5], as in our case.

Endoscopy_UCTN_Code_CCL_1AB_2AZ_3AB

Competing interests: None

J. Gerada¹, D. Babic², J. Degaetano², J. Pocock¹

¹ Division of Gastroenterology, Mater Dei Hospital, Msida, Malta

² Department of Histopathology, Mater Dei Hospital, Msida, Malta



Fig. 1 Pedunculated polyp in the duodenal bulb before endoscopic snare polypectomy.




Fig. 3 Scattered abnormal mitoses (hematoxylin and eosin, $\times 650$).




Fig. 2 Basal cystic change in crypts, and crypt epithelial atypia (hematoxylin and eosin, $\times 50$).

References

- 1 *Culver EL, McIntyre AS.* Sporadic duodenal polyps: classification, investigation, and management. *Endoscopy* 2011; 43: 144–155
- 2 *Abbass R, Rigaux J, Al-Kawas F.* Nonampullary duodenal polyps: characteristics and endoscopic management. *Gastrointes Endosc* 2010; 71: 754–759
- 3 *Okano A, Takakuwa H, Matsubayashi Y.* Gastric intramucosal carcinoma in a small hyperplastic foveolar polyp. *Endoscopy* 2004; 36: 1134
- 4 *Rosty C, Buchanan D, Walters R et al.* Hyperplastic polyp of the duodenum: a report of 9 cases with immunohistochemical and molecular findings. *Hum Pathol* 2011; 42: 1953–1959
- 5 *Kawaratani H, Tsujimoto T, Nishimura N et al.* A case of lobulated and pedunculated duodenal hyperplastic polyp treated with snare polypectomy. *Case Rep Gastroenterol* 2011; 5: 404–410

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1309395>
Endoscopy 2012; 44: E244–E245
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

J. Gerada, MD
Mater Dei Hospital
Msida
Malta
Fax: +356-25457582
jurgen.gerada@gmail.com