

Pharmacokinetics and Bioequivalence Study of Two Indapamide Formulations after Single-dose Administration in Healthy Chinese Male Volunteers

Authors

G. Li^{1,2}, X. Zhang^{1,2}, Y. Tian^{1,2}, Z. Zhang^{1,2}, J. Rui³, X. Chu³

Affiliations

¹ Key Laboratory of Drug Quality Control and Pharmacovigilance, China Pharmaceutical University, Ministry of Education, Nanjing, P. R. China

² Center for Instrumental Analysis, China Pharmaceutical University, Nanjing, P. R. China

³ Nanjing Command General Hospital, Nanjing, P. R. China

Key words

- indapamide
- pharmacokinetics
- bioequivalence
- healthy volunteers
- HPLC-MS
- human plasma

Abstract

The pharmacokinetics and relative bioavailability/bioequivalence of 2 formulations of indapamide were assessed in this paper. The study was conducted in 20 healthy Chinese male volunteers according to an open, randomized, single-blind, 2-way crossover study design with a wash-out phase of 7 days. Blood samples for pharmacokinetic profiling were taken up to 84h post-dose, and indapamide concentrations in plasma were determined by a validated liquid chromatography-electrospray ionization-mass spectrometry (LC-ESI-MS) method. Based on the plasma concentration-time data of each individual in each period, pharmacokinetic parameters, C_{max} , AUC_{0-t} , $AUC_{0-\infty}$ and $t_{1/2}$, were

calculated by non-compartmental analysis. Pharmacokinetic parameters for test and reference formulations were analyzed statistically to assess bioequivalence of the 2 formulations. The values of C_{max} , T_{max} , $t_{1/2}$, AUC_{0-t} , $AUC_{0-\infty}$ for test and reference formulations were 49.53 ± 5.53 and 47.79 ± 4.68 ng/mL, 1.9 ± 0.6 and 2.0 ± 0.5 h, 22.49 ± 5.93 and 23.23 ± 4.48 h, 859.51 ± 160.92 and 840.90 ± 170.62 ng·h/mL, 934.35 ± 190.60 and 919.52 ± 179.74 ng·h/mL, respectively. The 90% confidence intervals of the T/R-ratios of logarithmically transformed data were within the accepted range of 80–125%. It showed that the 2 formulations of indapamide were bioequivalent. Both preparations were well tolerated and no adverse reactions were found throughout the study.

received 08.06.2012

accepted 07.11.2012

Bibliography

DOI <http://dx.doi.org/10.1055/s-0032-1331181>

Published online:

December 13, 2012

Drug Res 2013;

63: 13–18

© Georg Thieme Verlag KG

Stuttgart · New York

ISSN 2194-9379

Correspondence

Dr. Z.-j. Zhang

Center for Instrumental

Analysis

China Pharmaceutical

University

24 Tongjia Lane

Nanjing 210009

(P. R. China)

Tel.: +86/25/8327 1454

Fax: +86/25/8327 1454

zunjianzhangcpu@hotmail.com

Introduction

Indapamide, 4-chloro-N-(2-methyl-1-indolyl)-3-sulphamoyl benzamide, is a non-thiazide indole derivative of O-chloro benzene sulfonamide, which has diuretic and natriuretic effects and thus exhibits an anti-hypertensive action causing a decrease in systolic, diastolic and mean blood pressure [1,2]. However, Specific side chains endows the drug some characteristic properties. Indapamide presents an indolyl ring and uniquely possesses free-radical scavenging activity as well as a direct vasodilator action [2–4], besides it has no adverse lipid effects in contrast to thiazide diuretics [5–7]. Accordingly, indapamide was defined by the 1999 WHO/ISH Hypertension Guidelines and JNC VII as a first-line drug for treating mild-to-moderate hypertension [8].

Indapamide was first developed as 2.5 mg immediate release tablets and subsequently 1.5 mg sustained release tablets. Indapamide is quickly absorbed from the gastrointestinal tract and peak plasma concentrations are seen 1–2 h after dosing immediate release tablets. Indapamide is

widely distributed throughout the body, with extensive binding to some specific sites, which leads its relatively highly combination to plasma proteins (79%) in plasma [9]. It has a long half-life (about 20h) so it is effective in once daily dosing [10]. Indapamide is extensively metabolised, with only about 7% of the total dose administered recovered in the urine as unchanged drug during the first 48 h after administration [10,11]. Although Li Ding et al. [12], Wei-Dong Chen et al. [13] and Tai-Jun Hang et al. [9] have researched the pharmacokinetics and bioavailability of indapamid in the Chinese population, they focus on the method for the quantitation of indapamide in vivo, which is simple, sensitive and rapid. The information concerning the bioavailability and pharmacokinetics of indapamide in the Chinese population has not been reported in detail up to now. In this article, pharmacokinetic parameters of indapamid in Chinese after dosing 5 mg indapamide are presented. Meanwhile, this study also focused on a comparison of the bioavailability of generic drug (test) and original drug (reference) formulations of indapamid of the same

strength (5 mg) produced by 2 different manufacturers. The generic drug product is therapeutically equivalent to the original product and can be used alternatively. Besides, as a way of curbing healthcare expenditure, many countries including China have taken measures to boost their generics markets. Therefore, it is of great significance to evaluate bioequivalence of the generic drug reasonably.

Materials and Methods

Materials

Indapamide test tablets (2.5 mg/tablet; Batch No. 20110302) were supplied by Jiangsu Fubang Pharmaceutical Co. Ltd., (Jiangsu, P. R. China); Indapamide reference tablets (2.5 mg/tablet; Batch No. 869000) were obtained from Servier Pharmaceutical Co. Ltd., France. Indapamide standard and prednisone standard (IS) were purchased from the Chinese National Institute for the Control of Pharmaceutical and Biological Products (Beijing, P. R. China).

Methanol was HPLC grade and purchased from Merck (Merck Company, Darmstadt, Germany). Other chemicals were all of analytical grade. Deionized water was distilled twice before use, and the mobile phase was filtered with a 0.22 µm film before use. Other reagents were prepared as required.

Analytical method

Preparation of stock solutions, calibration standards and quality control samples

The stock solution of indapamide and IS were prepared in methanol at concentration of 1.0 mg/ml. Working solutions of indapamide (0.05, 0.1, 0.25, 0.5, 1, 2.5, 4 and 5 µg/mL) and IS (30 µg/mL) were prepared daily by appropriate dilution.

Calibration standards (1, 2, 5, 10, 20, 50 and 100 ng/mL) were prepared by spiking 500 µL blank plasma with 10 µL working solutions (0.05, 0.1, 0.25, 0.5, 1, 2.5 and 5 µg/mL) of indapamide. In each run, a blank plasma sample (processed without IS) was analyzed to confirm absence of interference but not used to construct the calibration function. Quality control (QC) samples were prepared by spiking 500 µL blank plasma with 10 µL working solutions (0.1, 1 and 4 µg/mL) to produce a final concentration equivalent to 2 ng/mL (low level), 20 ng/mL (middle level) and 80 ng/mL (high level) of indapamide.

Sample preparation and instrumental conditions

A simple, sensitive and specific liquid chromatography-electrospray ionization-mass spectrometry (LC-ESI-MS) method has been developed and validated for the determination of indapamide in human plasma.

The plasma samples were prepared by liquid-liquid extraction. After the addition of prednisone as the internal standard (IS, 10 µL × 30 µg/mL) and 100 µL saturated Na₂CO₃, 500 µL plasma samples were extracted with 4.0 mL diethyl ether. Then the upper organic phase was transferred and evaporated to dryness. The residue was reconstituted in 100 µL of mobile phase, and an aliquot of 10 µL was injected to LC-ESI-MS (SHIMADZU LCMS-2010 A) for analysis.

The analytical procedure was performed on an Agilent Zorbax Eclipse XDB-phenyl (150 mm × 2.1 mm I.D., 5 µm) at 40 °C with a mobile phase of methanol- 10 mM ammonium acetate buffer solution (pH=5.0) (50:50, v/v). Analytes were determined in a single quadrupole mass spectrometer with negative ion and

selected ion monitoring (SIM) mode to monitor target ions at *m/z* of 364.0 for indapamide and 357.0 for prednisone. The MS operating conditions were optimized as follows: drying gas 1.5 L/min, CDL temperature 250 °C, block temperature 200 °C, probe voltage -4.5 kV.

Method validation

The method was validated in our analytical laboratory according to the United States Food and Drug Administration (FDA) bioanalytical method validation guidance [14] on specificity, matrix effect, linearity, accuracy, precision, extraction recovery and stability.

Specificity was assessed by testing 6 different drug-free plasma batches. The matrix effect caused by ionization competition between analytes (indapamide and IS) and co-eluted compounds was evaluated at 3 QC concentrations. 5 calibration curves (1, 2, 5, 10, 20, 50, 100 ng/mL) were generated in 5 consecutive days to test the linearity. The limit of detection (LOD) and the lower limit of quantification (LLOQ) were determined as the concentrations with a signal-to-noise ratio (S/N) no less than 3 and 10, respectively. The accuracy and precision were evaluated by analyzing QC standards 5 times daily and over 3 consecutive days. Extraction recovery was evaluated by comparing the indapamide-IS peak area ratios obtained from extracted samples to those from the samples containing the same amount of indapamide which was added after the extraction step (IS was added after the extraction step in both batches). Stability was evaluated at each QC concentration under different conditions that occurred during sample analysis. The short-term stability and long-term stability was evaluated by keeping QC samples at room temperature for 8 h and low temperature (-20 °C) for 15 days respectively. The post-preparative stability was measured by placing prepared QC samples under the autosampler conditions (15 °C) for 48 h. The freeze and thaw stability was tested by analyzing QC samples undergoing 3 freeze (-20 °C)-thaw (room temperature) cycles on consecutive days. The working solutions and stock solutions of indapamide and the IS were also evaluated at room temperature for 24 h and at 4 °C for 15 days, respectively.

Experimental design

This open, randomized, single-dose, 2-sequence, 2-period crossover study was performed in strict accordance with the Declaration of Helsinki, Good Clinical Practice (GCP) Guidelines and Good Laboratory Practice (GLP) Guidelines. The protocol and the consent form of the study were approved by the Research Ethics Committee of Nanjing Command General Hospital, Nanjing (P. R. China).

Enrolled in this study were 20 healthy, male, adult Chinese volunteers provided written informed consent prior to study participation. All of them completed the study and were evaluable for this pharmacokinetics and bioequivalence study. Their mean age ± SD was 23.60 ± 1.39 years. Their mean body weight was 66.15 ± 7.01 kg and their mean height was 172.85 ± 3.18 cm, giving a mean Body Mass Index (BMI) of 22.11 ± 1.94 kg/m². Screening at study start included medical history, physical examination, vital signs, ECG, laboratory safety tests (routine hematology; glucose, creatinine, sodium, potassium, chloride, CO₂, ALT, ALP, and total bilirubin level; and urinalysis). Subjects were excluded if they smoked, used alcohol, had a history or evidence of hepatic, renal, gastrointestinal, or hematological deviations, had acute or chronic disease, allergies to any medications, and/or had any clinically significant abnormality. Subjects were instructed to abstain from using any

medications for at least 2 weeks prior to and during the study; and from smoking and taking alcohol, caffeine, related xanthenes-containing beverages or food for at least 48 h before administering the study drug and throughout the study.

This was a single-dose, single-blind, randomized, 2-way crossover trial in which the subjects were randomly assigned to the treatment sequences R/T or T/R. The subjects received a single dose of 5 mg indapamide as 2 tablets of formulation R (2 indapamide reference tablets) or a single dose of 5 mg indapamide as 2 tablets of formulation T (2 indapamide test tablets) under supervision by study investigators. The reference formulation and test formulation were administered alternately in the second period. The wash-out period between the 2 dosing periods was 1 week. In each treatment period, volunteers were required to fast for approximately 10 h prior to the dosing. The formulations were administered with 250 mL water, then water was not allowed until 2 h after administration. Standardized lunch and dinner were served after a 6- and 12-h blood sampling, respectively. For the measurement of pharmacokinetic parameters, venous blood samples were collected by study nurses via an indwelling cannula or direct venipuncture into the heparinized tubes from all subjects prior to the dosing (T_0) and then after 0.33, 0.67, 1, 1.5, 2, 3, 4, 6, 8, 12, 24, 48, 72 and 84 h from administration during each treatment period. Samples were centrifuged at 3000g for 5 min immediately then the plasma was transferred and stored at -20°C until assay.

Pharmacokinetic and statistical analysis

Pharmacokinetic parameters

Pharmacokinetic assessment of indapamide was estimated from the concentration-time data using non-compartmental methods. Maximum plasma concentrations (C_{\max}) and time to reach this value (T_{\max}) were considered as the observed highest concentration and time of its occurrence, respectively. The elimination half-life ($t_{1/2}$) was calculated as $t_{1/2} = (\ln 2)/K_{el}$, where the apparent elimination rate constant, K_{el} was calculated as the slope of the linear regression line of natural log-transformed plasma concentration against time. $AUC_{0-\tau}$ was calculated using the linear trapezoidal rule from zero to the last sampling time-point (T_{last}) at which a concentration was measurable (C_T). $AUC_{0-\infty}$ was calculated according to the following formula: $AUC_{0-\infty} = AUC_{0-\tau} + C_T/K_{el}$. The rate of absorption was evaluated by means of the ratio: $C_{\max}/AUC_{0-\infty}$. The ratio $AUC_{0-\tau}/AUC_{0-\infty}$ was determined as an indicator of the adequacy of the blood sampling period.

Statistical analysis

The pharmacokinetic primary variables for bioequivalence testing in this study were $AUC_{0-\infty}$, $AUC_{0-\tau}$ and C_{\max} . An analysis of variance (ANOVA) was performed on these parameters, after natural logarithmic transformation using general linear model procedure for randomized crossover design to evaluate the effects of formulation, period, sequence, and subject (within sequence) at the significance level (α) of 0.05. ANOVA was performed with the statistical software package, SAS (Statistical Analysis System, SAS Institute Inc., Cary, NC, USA), using the General Linear Model (GLM) Procedure. The Wilcoxon signed rank test was used for the non-parametric analysis to determine differences in T_{\max} . The sequence effect was tested against the mean squares term of subjects nested in sequence. All other effects were tested against the mean residual error. Effects were considered statistically significant if the probability obtaining the calculated F test was ≤ 0.05 . According to the guidance from the FDA, product bioequivalence

was based on the 90% confidence intervals (CIs) for the ratios of least squares means of the ln-transformed values for $AUC_{0-\infty}$, $AUC_{0-\tau}$ and C_{\max} . In addition, the bioequivalence between the 2 formulations was assessed by Schurimann's 2 one-sided t-tests.

Results



Analytical method

The representative chromatograms of indapamide are depicted in **Fig. 1a-c**. As shown, the retention times of indapamide and IS were 7.8 and 11.4 min, respectively. The total LC-MS analysis time was 13 min per sample.

The sensitivity and specificity of the method were proved by testing 6 different plasma batches (**Fig. 1a**). No significant matrix effect for indapamide and the IS was implied in the method. The calibration curve was linear over the concentration range of 1–100 ng/mL, with $r^2 > 0.99$. LLOQ for indapamide proved to be 1 ng/mL (**Fig. 1b**), and LOD was 0.2 ng/mL. The accuracy and precision at 3 QC levels (2, 20 and 80 ng/mL) are $100.8\% \pm 10.8\%$, $102.2\% \pm 8.2\%$ and $100.9\% \pm 5.1\%$ for intra-batch, and $100.6\% \pm 9.1\%$, $101.4\% \pm 9.2\%$ and $97.6\% \pm 6.8\%$ for intra-batch. The accuracy and precision at LLOQ are within $(100 \pm 20)\%$ and 20%, respectively. The data of extraction efficiency measured for indapamide in human plasma was consistent, precise and reproducible. The results (**Table 1**) showed that both indapamide and the IS were stable under the studied conditions. **Table 1** showed the data for intra-batch and inter-batch precision and accuracy of the method, extraction recovery and sample stability test.

Clinical observations

20 subjects completed the study so that 20 cases were available for each treatment. No clinically significant adverse effects were reported in the present study population.

All subjects were discharged in good health after repeating physical examination and laboratory analysis after period 2. A clinically relevant difference in both tolerability and safety of 2 treatments was not detected.

Pharmacokinetic and statistical results

The plasma concentration-time profiles of indapamide following single-dose oral administration of the test and reference formulations are presented in **Fig. 2**. A summary of the pharmacokinetic parameters, $AUC_{0-\tau}$, $AUC_{0-\infty}$, C_{\max} , T_{\max} , $t_{1/2}$, MRT, $AUC_{0-\tau}/AUC_{0-\infty}$, $F_{0-\tau}$ and $F_{0-\infty}$ derived from the 2 formulations is shown in **Table 2**.

The calculated C_{\max} values for the test and reference formulations were 49.53 ± 5.53 ng/mL and 47.79 ± 4.68 ng/mL, $AUC_{0-\infty}$ of 934.35 ± 190.60 ng·h/mL (test) and 919.52 ± 179.74 ng·h/mL (reference) were achieved. Elimination half-lives and elimination rate constants were not significantly different between the preparations. In addition, the mean T_{\max} values in this study were 1.9 ± 0.6 h for test formulation and 2.0 ± 0.5 h for reference formulation.

The results of ANOVA for assessment of treatment, period and subject (within sequence) effects and 90% CIs for the ratio of C_{\max} , $AUC_{0-\tau}$ and $AUC_{0-\infty}$ values for test and reference formulations, using logarithmic transformed data, are shown in **Table 3**.




Fig. 1 The representative chromatograms of indapamide. **a** Drug-free sample, **b** LOQ (1 ng/mL), **c** plasma sample of a healthy volunteer with determined indapamide concentration of 21.74 ng/mL. The retention time of indapamide and IS were 7.8 and 11.4 min, respectively.

Discussion and Conclusion


The proposed method for determination of indapamide in human plasma was selective, accurate, and precise. And it was proved to

be reliable and suitable for the pharmacokinetics and bioequivalence study of indapamide in healthy Chinese male volunteers. In our method, liquid-liquid extraction was necessary and important, because this technique can purify and concentrate

Table 1 Accuracy, precision, extraction recovery and stability data for method validation.

Plasma samples	2 ng/mL		20 ng/mL		80 ng/mL	
	Accuracy (%)	Precision (%)	Accuracy (%)	Precision (%)	Accuracy (%)	Precision (%)
Intra-batch (n=5)	100.8	10.8	102.2	8.2	100.9	5.1
Inter-batch (n=15)	100.6	9.1	101.4	9.2	97.6	6.8
Extraction recovery ¹	83.2	5.3	82.7	8.1	87.1	4.7
Long-term stability ²	97.3	7.3	102.2	6.9	96.7	7.5
Short-term stability ³	100.0	3.7	107.4	3.0	93.1	4.6
Freeze and thaw stability ⁴	97.9	6.2	101.3	7.7	96.3	6.4
Post-preparative stability ⁵	101.0	6.2	104.7	8.2	96.4	7.6
Standard solutions	Indapamide (25ng/mL)		IS (15µg/mL)		-	
	Accuracy (%)	Precision (%)	Accuracy (%)	Precision (%)	-	
working solution Stability ⁶	105.5	4.1	104.8	5.8	-	
stock solution Stability ⁷	105.9	2.5	105.8	7.7	-	

¹n=5; ²15 days, -20°C, n=6; ³8 h, room temperature, n=6; ⁴3 cycles, -20°C-room temperature, n=6; ⁵48 h, 15°C, n=9; ⁶24 h, room temperature, n=3; ⁷15 days, 4°C, n=3

**Fig. 2** Mean plasma concentration-time profiles of indapamide in 20 Chinese male volunteers after oral administration of 5 mg indapamide.**Table 2** Pharmacokinetic parameters of indapamide in 20 male volunteers after oral administration of 5 mg indapamide.

Parameters	Indapamide	
	T	R
C _{max} (ng/mL)	49.53 ± 5.53	47.79 ± 4.68
T _{max} (h)	1.9 ± 0.6	2.0 ± 0.5
t _{1/2} (h)	22.49 ± 5.93	23.23 ± 4.48
MRT (h)	29.17 ± 7.52	30.19 ± 4.94
AUC ₀₋₈₄ (ng·h/mL)	859.51 ± 160.92	840.90 ± 170.62
AUC _{0-∞} (ng·h/mL)	934.35 ± 190.60	919.52 ± 179.74
AUC _{0-∞} /AUC ₀₋₈₄ (%)	92.41 ± 4.35	91.42 ± 3.72
C _{max} /dose (ng/mL)	9.91 ± 1.11	9.56 ± 0.94
AUC _{0-∞} /dose (ng·h/mL)	186.87 ± 38.12	183.90 ± 35.95
F ₀₋₈₄ (%)	102.71 ± 8.57	
F _{0-∞} (%)	102.03 ± 12.64	

R: reference tablets; T: test tablets; mean ± SD, n=20

the sample. A preliminary evaluation of different extraction solvents such as diethyl ether, ethyl acetate or dichloromethane under alkalizing and neutral conditions showed that extraction with diethyl ether in alkalizing condition exhibited higher extraction efficiency. Then, 1M NaOH, saturated Na₂CO₃ solution and saturated NaHCO₃ solution were attempted. Finally, 100 µL of saturated Na₂CO₃ solution was adopted for sample preparation in terms of precise, consistent and reproducible recoveries and Specific.

To exclude the interference of unknown peak to indapamide peak (showed in **Fig. 1c**), a mobile phase consisting of 50%

methanol and 50% 10mM ammonium acetate buffer solution (pH=5.0) was adopted, achieved good resolution, symmetric peak shape and suitable retention time without matrix effects.

Several articles [15,16] have reported the pharmacokinetics of indapamide in healthy human. Pierre Schiavi [15] researched the pharmacokinetics of sustained and immediate release formulations (1.5 mg and 2.5 mg respectively) of indapamide, indicating much higher values of C_{max}/dose and AUC_{0-∞}/dose (39 ± 11 ng/mL and 564 ± 146 ng·h/mL after dosing one 2.5 mg immediate release tablet, determined and calculated by use of concentrations in whole blood) than this study. The main reason should be that indapamide is preferentially and reversibly taken up by the erythrocytes in the blood [17]. Moreover, Dragica Zendelovska [18] developed a new method for determining indapamide in biological fluids to confirm that the C_{max} in blood was much higher than in serum. Consequently, it should be remarked that blood samples should avoid hemolysis until to obtain the plasma. Moreover, bioequivalence study of indapamide in plasma requires a more sensitive analytical method than in whole blood. LLOQ of 1 ng/mL in this study is enough for plasma concentration determination. In contrast, HPLC-UV methods [9,18], giving a higher LLOQ of 10 ng/mL, are not suitable for pharmacokinetics study of indapamide in plasma. Florin Albu [16] and Deepak S. Jain [19] reported HPLC-MS/MS methods with LLOQ of 1 ng/mL and 0.5 ng/mL respectively, but they adopted a superior instrument (HPLC-MS/MS), which is more expensive than HPLC-MS. Consequently, HPLC-MS is economic and suitable for pharmacokinetics study of indapamide.

The results from the ANOVA found that formulation, period, and sequence had no statistically significant effect on AUC_{0-t}, AUC_{0-∞} and C_{max} at the significance level of 0.05. However, as might be expected, the subject variability of those pharmacokinetic parameters was significant (P<0.0001). This was further explored by calculating the inter-subject and intra-subject coefficient of variations. The mean ratio of AUC_{0-t}/AUC_{0-∞} for test and reference formulations were both above 90%, indicated that the sampling time was adequate. The 90% CIs for AUC_{0-t}, AUC_{0-∞} and C_{max} were within the FDA acceptable bioequivalence range of 80–125% and the results of the Schuirmann's test indicated that the lower and upper limits of the calculated t-test were greater than the critical t-value.

Therefore, it can be concluded that the test and reference formulations are bioequivalent according to both of the rate and extent of absorption and consequently can be used interchangeably in

Table 3 Statistical analysis of log-transformed data.

	Parameter	ANOVA (p-Value)				t-Test	
		Treatment	Period	Subject in sequence	90% CI (%)	Lower	Upper
Mosapride citrate	C _{max}	0.0591	0.6133	0.0001	100.48–106.55	19.11	23.08
	AUC ₀₋₈₄	0.2325	0.5981	<0.0001	99.06–105.79	10.53	13.01
	AUC _{0-∞}	0.6217	0.1091	<0.0001	96.85–105.97	8.1	9.1

medical practice. The results of this study, obtained from samples of healthy Chinese volunteers, might serve as a reference for future controlled studies of indapamide in the Chinese population.

Acknowledgement

We thank all co-workers in Nanjing Command General Hospital and China Pharmaceutical University. The study was funded by Jiangsu Fubang Pharmaceutical Co. Ltd., (Jiangsu, P. R. China).

Conflict of Interest Statement

The authors declare that they have no conflict of interest with respect to this paper.

References

- Ganado P, Ruiz E, Del Rio M *et al.* Growth inhibitory activity of indapamide on vascular smooth muscle cells. *Eur J Pharmacol* 2001; 428: 19–27
- Campbell DB, Taylor AR, Moore RA. Observations on the pharmacokinetics of indapamide in man. *Br J Clin Pharmacol* 1976; 3: 971
- Chillon JM, Baumbach GL. Effects of indapamide, a thiazide-like diuretic, on structure of cerebral. *Hypertension* 2004; 43: 1092–1097
- Inaba M, Noguchi Y, Yamamoto T *et al.* Effects of a low dose of indapamide, a diuretic, given daily or every-other-day on blood pressure and metabolic parameters. *Hypertens Res* 2004; 27: 141–145
- Kyncl J, Oheim K, Seki T *et al.* Antihypertensive action of indapamide. Comparative studies in several experimental models. *Arzneimittelforschung* Oct 1975; 25 (10): 1491–1495
- Ames RP. A comparison of blood lipid and blood pressure responses during the treatment of systemic hypertension with indapamide and with thiazides. *Am J Cardiol* 1996; 77: 12B–16B
- Carey PA, Sheridan DJ, Cordoue A *et al.* Effect of indapamide on left ventricular hypertrophy in hypertension. *Am J Cardiol* 1996; 77: 17B–19B
- Ren C, Fang L, Ling L *et al.* Design and in vivo evaluation of an indapamide transdermal patch. *International Journal of Pharmaceutics* 2009; 370: 129–135
- Hang TJ, Zhao W, Liu J *et al.* A selective HPLC method for the determination of indapamide in human whole blood: Application to a bioequivalence study in Chinese volunteers. *Journal of Pharmaceutical and Biomedical Analysis* 2006; 40: 202–205
- Livingstone K, McKay G, Fisher M. Indapamide. *Pract Diab Int Sep* 2008; 25 (7): 290–291
- Caruso FS, Szabadi RR, Vukovich RA. Pharmacokinetics and clinical pharmacology of indapamide. *Am heart J* 1983 Jul; 106 (1 Pt 2): 212–220
- Ding L, Yang L, Liu F *et al.* A sensitive LC–ESI–MS method for the determination of indapamide in human plasma: Method and clinical applications. *Journal of Pharmaceutical and Biomedical Analysis* 2006; 42: 213–217
- Chen WD, Liang Y, Zhang H *et al.* Simple, sensitive and rapid LC–MS method for the quantitation of indapamide in human plasma—application to pharmacokinetic studies. *J Chromatogr B* 2006; 842: 58–63
- US Department of Health and Human Services. Food and Drug Administration, Center for Drug Evaluation and Research (CDER), Guidance for Industry. *Bioanalytical Method Validation*. 2001
- Schiavi P, Jochemsen R, Guez D. Pharmacokinetics of sustained and immediate release formulations of indapamide after single and repeated oral administration in healthy volunteers. *Fundamental & Clinical Pharmacology* 2000; 14: 139–146
- Florin A, Cristina G, Victor D *et al.* Liquid chromatography–electrospray tandem mass spectrometry method for determination of indapamide in serum for single/multiple dose bioequivalence studies of sustained release formulations. *J Chromatogr B* 2005; 816: 35–40
- Campbell DB, Taylor AR, Hopkins YW *et al.* Pharmacokinetics and metabolism of indapamide: A review. *Current Medical Research and Opinion* 1977; 5 (Suppl 1): 13–24
- Zendelovska D, Stafilov T, Stefova M. Optimization of a solid-phase extraction method for determination of indapamide in biological fluids using high-performance liquid chromatography. *J Chromatogr B* 2003; 788: 199–206
- Jain DS, Subbaiah G, Sanyal M *et al.* Liquid chromatography–tandem mass spectrometry validated method for the estimation of indapamide in human whole blood. *J Chromatogr B* 2006; 834: 149–154