

# Cost-Effectiveness of Trial of Labor after Previous Cesarean in a Minimally Biased Cohort

Sharon A. Gilbert, MS, MBA<sup>1</sup> William A. Grobman, MD, MBA<sup>2</sup> Mark B. Landon, MD<sup>3</sup>  
 Catherine Y. Spong, MD<sup>4</sup> Dwight J. Rouse, MD<sup>5</sup> Kenneth J. Leveno, MD<sup>6</sup> Michael W. Varner, MD<sup>7</sup>  
 Ronald J. Wapner, MD<sup>8</sup> Yoram Sorokin, MD<sup>9</sup> Mary J. O'Sullivan, MD<sup>10</sup> Baha M. Sibai, MD<sup>11</sup>  
 John M. Thorp, MD<sup>12</sup> Susan M. Ramin, MD<sup>13</sup> Brian M. Mercer, MD<sup>14</sup> for the *Eunice Kennedy Shriver* National  
 Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network

<sup>1</sup>The George Washington University Biostatistics Center,  
 Washington, District of Columbia

<sup>2</sup>The Department of Obstetrics and Gynecology, Northwestern  
 University, Chicago, Illinois

<sup>3</sup>The Ohio State University, Columbus, Ohio

<sup>4</sup>The *Eunice Kennedy Shriver* National Institute of Child Health and  
 Human Development, Bethesda, Maryland

<sup>5</sup>University of Alabama at Birmingham, Alabama

<sup>6</sup>University of Texas Southwestern Medical Center, Dallas, Texas

<sup>7</sup>University of Utah, Salt Lake City, Utah

<sup>8</sup>Thomas Jefferson University, Philadelphia, Pennsylvania

<sup>9</sup>Department of Obstetrics and Gynecology, Wayne State University,  
 Detroit, Michigan

<sup>10</sup>University of Miami, Miami, Florida

<sup>11</sup>University of Tennessee, Memphis, Tennessee

<sup>12</sup>University of North Carolina at Chapel Hill, Chapel Hill, North  
 Carolina

<sup>13</sup>The University of Texas Health Science Center at Houston, Houston,  
 Texas

<sup>14</sup>Case Western Reserve University-MetroHealth Medical Center,  
 Cleveland, Ohio

\* Other members of the *Eunice Kennedy Shriver* National Institute of  
 Child Health and Human Development Maternal-Fetal Medicine  
 Units Network are listed in the Acknowledgments

Address for correspondence and reprint requests Sharon A. Gilbert,  
 MS, MBA, The George Washington University, 6110 Executive Blvd,  
 Suite 750, Rockville, MD 20852 (e-mail: sgilbert@bsc.gwu.edu).

Am J Perinatol 2013;30:11–20.

## Abstract

### Keywords

- ▶ cost-effectiveness
- ▶ elective repeat cesarean
- ▶ trial of labor
- ▶ propensity scores

**Objective** To estimate the cost-effectiveness of a trial of labor after one previous cesarean delivery (TOLAC).

**Study Design** A model comparing TOLAC with elective repeat cesarean delivery (ERCD) was developed for a hypothetical cohort with no contraindication to a TOLAC. Probabilistic estimates were obtained from women matched on their baseline characteristics using propensity scores. Cost data, quality-adjusted life-years (QALYs), and data on cerebral palsy were incorporated from the literature.

**Results** The TOLAC strategy dominated the ERCD strategy at baseline, with \$138.6 million saved and 1703 QALYs gained per 100,000 women. The model was sensitive to five variables: the probability of uterine rupture, the probability of successful TOLAC, the

received  
 November 29, 2012  
 accepted after revision  
 December 6, 2012  
 published online  
 January 2, 2013

Copyright © 2013 by Thieme Medical  
 Publishers, Inc., 333 Seventh Avenue,  
 New York, NY 10001, USA.  
 Tel: +1(212) 584-4662.

DOI <http://dx.doi.org/10.1055/s-0032-1333206>.  
 ISSN 0735-1631.

QALY of failed TOLAC, the cost of ERCD, and the cost of successful TOLAC without complications. When the probability of TOLAC success was at the base value, 68.5%, TOLAC was preferred if the probability of uterine rupture was 4.2% or less. When the probability of uterine rupture was at the base value, 0.8%, the TOLAC strategy was preferred as long as the probability of success was 42.6% or more.

**Conclusion** A TOLAC is less expensive and more effective than an ERCD in a group of women with balanced baseline characteristics.

In 2003 the Agency for Healthcare Research and Quality (AHRQ) published the results of an evidence report and technology assessment comparing the harms and benefits of delivery options for women after a prior cesarean delivery.<sup>1</sup> Incorporated in this report was a review of the economic literature on this topic. Of 13 publications reviewed, only two had methodologies that were highly rated.<sup>2,3</sup> These two studies compared the cost-effectiveness of a trial of labor after a previous cesarean (TOLAC) with an elective repeat cesarean delivery (ERCD). Both concluded that TOLAC could be cost-effective when the probability of a successful TOLAC was sufficiently high.

Inherent to these analyses was the assumption that the probabilistic data used in the decision analytical models were derived from two groups (i.e., those women who had TOLAC and those who had ERCD) with similar baseline characteristics. However, this assumption was not accurate, as the studies from which the probabilistic data were derived actually were composed of two groups of women—those who underwent TOLAC and those who underwent ERCD—with different baseline characteristics.<sup>4–7</sup> Consequently, it is uncertain whether the data used in the decision analytic models were biased and led to a bias in the ultimate results.

Ideally, a trial in which women were randomized to one of the delivery approaches would balance observed as well as unobserved baseline covariates and would produce the true treatment effect, but such a trial is unlikely to be undertaken. In the absence of a randomized trial, propensity score analysis may be used in an observational study to derive two groups with similar baseline characteristics. The data from these groups can then be used in decision analytic models with less concern for biased outcomes. Thus, this analysis was undertaken to determine whether TOLAC or ERCD is the more cost-effective strategy after one prior cesarean based on data derived from groups of women with similar baseline characteristics.

## Methods

We developed a decision analysis model comparing a TOLAC with an ERCD for a hypothetical cohort of 100,000 women with no contraindication to a TOLAC. The analysis was based on the societal perspective, incorporating all health outcomes and economic costs regardless of who experienced the outcome or paid the costs.<sup>8</sup> The primary outcome was cost-effectiveness, measured as the marginal cost per quality-

adjusted life-year (QALY) gained, with a marginal cost per QALY ratio of less than \$50,000 used to indicate a strategy is cost-effective.

The decision tree was developed using TreeAge Pro 2009 (TreeAge Software, Inc. Williamstown, MA). The initial decision represented a woman's approach to delivery, either a TOLAC or an ERCD. Women in the TOLAC arm experienced either a successful vaginal delivery, required a repeat cesarean during labor, or had a uterine rupture in association with a successful or failed TOLAC. Additional maternal and neonatal morbidity that occurred depended upon these outcomes or upon the alternate choice of an ERCD.

The probabilities for the decision tree primarily were obtained from data collected from 1999 through 2002 in a registry (the Cesarean Registry) by the Eunice Kennedy Shriver National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network. Nineteen academic centers comprising 32 hospitals throughout the United States participated in this observational study, in which data were collected on all women with a prior cesarean delivery. Institutional review board approval at each participating center was obtained. Study personnel at the medical centers abstracted data from patient charts under a waiver of informed consent. Further detail on specific methods of the study can be obtained from previously published articles.<sup>5,9</sup>

Women who were eligible to have either an ERCD or a TOLAC and who had a singleton, term, vertex gestation and one prior low transverse incision ( $n = 22,068$ ) comprised the population that was analyzed to obtain probabilities requisite for the model. A gestation was considered "term" if delivery occurred at or beyond 37 weeks' gestation. An ERCD was defined as a cesarean delivery without any indication other than the prior cesarean. Thus, women who had a repeat cesarean for indications such as placenta previa or active herpes were excluded from this analysis ( $n = 759$ ). To ensure that women who underwent ERCD truly had no indication for the cesarean other than their choice, those who were reported to have a cesarean that was elective but who had an additional reported indication implying this was not the case (i.e., cephalopelvic disproportion, failure to progress, cord prolapse, nonreassuring tracing or abruption) were excluded ( $n = 262$ ). Also, women were ineligible for the cohort if they had an ERCD prior to 39 weeks without spontaneous labor or premature rupture of membranes given that elective delivery prior to 39 weeks is associated with known adverse outcomes unrelated to mode of delivery ( $n = 3188$ ).<sup>10</sup> Women who

underwent ERCD, after experiencing spontaneous labor or rupture of membranes prior to 39 weeks, were included because they were eligible for, and would need to choose between, either ERCD or labor. Women who underwent a TOLAC but whose labors were induced also were excluded ( $n = 3235$ ) because this intervention has been associated with a lower probability of success and a higher probability of uterine rupture and is not a probabilistic possibility but a choice that a woman and her provider make.<sup>11,12</sup> Women with fetuses with major congenital malformations were not included because these conditions, unrelated to mode of delivery, would influence the newborn outcome ( $n = 120$ ). Consequently, 14,504 women were available for analysis, of whom 8297 had a TOLAC and 6207 had an ERCD.

The propensity score methodology (R MatchIt library, <http://www.r-project.org/>) of one-to-one matching without replacement using the nearest Mahalanobis distance<sup>13</sup> was used to derive 3981 matched pairs of women who underwent either ERCD or TOLAC and who were balanced according to 43 baseline characteristics. This final cohort, approximately 80% of whom had no previous vaginal delivery, had a TOLAC success rate of 68.1%. All 31 cases of uterine rupture occurred in the TOLAC group, 27 in association with TOLAC failure. Further detailed information on the development of this cohort and the propensity score methodology can be found in Gilbert et al.<sup>14</sup>

The maternal outcomes recorded through delivery included endometritis (clinical diagnosis of puerperal uterine infec-

tion in the absence of findings suggesting another source), wound complication (seroma, hematoma, or infection), operative injury (broad ligament hematoma, cystotomy, or bowel or ureteral injury), peripartum hysterectomy, thromboembolism (deep vein thrombosis or pulmonary embolus), and maternal death. Cases of uterine rupture, modeled as a separate branch of the decision tree, were defined as a disruption or tear of the uterine muscle and visceral peritoneum or a uterine muscle separation with extension to adjacent structures. Neonatal outcomes recorded up to 120 days after delivery or hospital discharge (whichever occurred first) were acidemia (arterial cord pH less than 7.0), transient tachypnea of the newborn, respiratory distress syndrome (RDS), proven or confirmed sepsis, hypoxic ischemic encephalopathy (HIE), and infant death. A separate maternal and infant mutually exclusive hierarchy incorporating these outcomes was developed following the reverse order above, with death first. Because the Cesarean Registry was a short-term observational study and cerebral palsy (CP) could occur as a long-term consequence of an event (HIE) at the time of delivery, the probability of CP also was incorporated in the model by estimating that 12% of infants with HIE would ultimately be diagnosed with CP.<sup>15</sup>

The probabilities used in the model are shown in ► **Table 1**. The ranges were obtained from the 95% Blyth-Still-Casella binomial confidence intervals (from Stat-Xact, Cytel Software) based on the proportion of events in the matched data set.<sup>16</sup> Because the probability of successful TOLAC and

**Table 1** Probability Estimates in the Model

Outcomes	ERCD ( $n = 3981$ )		Uterine rupture ( $n = 31$ )		Failed TOLAC ( $n = 1244$ )		Successful TOLAC ( $n = 2706$ )	
	Baseline	Range	Baseline	Range	Baseline	Range	Baseline	Range
<b>Maternal</b>								
Death	0.025	0.001–0.138	0	0–9.733	0	0–0.278	0	0–0.128
Thromboembolism	0.025	0.001–0.138	0	0–9.733	0	0–0.278	0	0–0.128
Hysterectomy	0.277	0.138–0.489	3.226	0.165–16.060	0.161	0.028–0.558	0	0–0.128
Operative injury	0.126	0.050–0.288	12.900	4.530–28.600	1.206	0.676–1.967	0	0–0.128
Wound complication	0.855	0.601–1.175	0	0–9.733	1.206	0.676–1.967	0.074	0.013–0.256
Endometritis	2.111	1.705–2.599	12.900	4.530–28.600	7.556	6.149–9.103	1.516	1.090–2.023
<b>Infant</b>								
Death	0.025	0.001–0.138	3.226	0.165–16.060	0.080	0.004–0.443	0	0–0.128
HIE	0	0–0.087	3.226	0.165–16.060	0.080	0.041–0.443	0	0–0.128
Sepsis	3.115	2.597–3.687	29.030	15.760–46.650	6.758	5.426–8.292	4.250	3.521–5.054
RDS	0.528	0.327–0.804	0	0–9.733	1.046	0.558–1.767	0.517	0.300–0.841
TTN	1.181	0.879–1.552	0	0–9.733	2.172	1.436–3.140	0.628	0.366–0.996
Acidemia	0.226	0.112–0.427	6.452	1.159–20.030	0.724	0.360–1.340	0.037	0.002–0.203

Data presented as percent. Abbreviations: ERCD, elective repeat cesarean delivery; HIE, hypoxic ischemic encephalopathy; RDS, respiratory distress syndrome; TOLAC, trial of labor after a previous cesarean; TTN, transient tachypnea of the newborn.

uterine rupture have previously been shown to be variables to which the results are sensitive, these two variables were varied across a range wider than that which would have been derived from the dataset alone—34.0 to 100% and 0 to 5.0%, respectively.

With an exception for CP, the following costs were incorporated into the model and based on mode of delivery: hospital, obstetrician, pediatrician, anesthesiologist, and maternal and caregiver opportunity costs. A summary of these costs is provided in **Table 2** with further detail regarding the basis for these costs provided in the appendix. Hospital costs were obtained from the 2009 AHRQ's Healthcare Cost and Utilization Project Nationwide Inpatient Sample (HCUPnet), a nationwide database of hospital inpatient stays containing about 95% of all hospital discharges in the United States.<sup>17</sup> Based on the *International Classification of Diseases, Ninth Revision* codes, these costs represent direct and indirect costs. Obstetrician and pediatrician costs were obtained from the 2010 Current Procedural Terminology from the American Medical Association (AMA).<sup>18</sup> Because the Cesarean Registry did not contain data that would allow estimation of anesthesia costs, these costs were derived from the literature.<sup>3</sup> Maternal and caregiver postpartum opportunity costs were derived from the Bureau of Labor Statistics using the 2009 median hourly wage and salary averages for women 25 to 34 years old and for all individuals 16 years and older,

respectively.<sup>19</sup> Because the costs associated with maternal and infant death are hard to quantify, as these events occur in such a large variety of circumstances, a range of 0 to \$1 million was used, with baseline estimates of \$20,000 and \$50,000, respectively. For CP, hospital costs after delivery were estimated as twice the base cost of HIE, with the addition of approximately \$9,000 for pediatrician fees and \$23,800 per year for the next 49 years.<sup>20–22</sup>

In sensitivity analysis, costs were ranged from 50 to 400% of the base-case estimate with the exception of those associated with maternal and well infant discharge, in which cases a range of 50 to 150% was used. Although such ranges included values that appeared beyond plausible in some cases, such a wide range ensured that the plausible range was contained within the interval and that threshold analyses could be judiciously performed.<sup>23</sup>

With the exception of CP, all outcomes (e.g., wound infection) occurred and were resolved during the initial hospitalization. Correspondingly, for the costs associated with these variables, no discounting was performed. Conversely, CP continued to affect a child and incur health care costs through the child's life, and thus these costs were discounted at 3% annually in the base-case. All costs are presented in 2009 dollars, with adjustments used, when needed, according to the medical care component of the Consumer Price Index.<sup>24</sup>

**Table 2** Cost Estimates in the Model

Outcomes	ERCD		Uterine rupture		Failed TOLAC		Successful TOLAC	
	Baseline	Range	Baseline	Range	Baseline	Range	Baseline	Range
Maternal								
Death	27.41	0–1,000.0	27.9	0–1,000.0	27.9	0–1,000.0	24.1	0–1,000.0
Thromboembolism	16.61	8.3–66.4	19.1	9.6–76.5	18.4	9.2–73.5	13.3	6.7–53.3
Hysterectomy	17.9	8.9–71.5	20.4	10.2–81.5	19.6	9.8–78.6	14.6	7.3–58.4
Operative injury	13.9	6.9–55.6	16.4	8.2–65.6	15.7	7.8–62.6	10.6	5.3–42.5
Wound complication	15.6	7.8–62.3	18.1	9.0–72.3	17.3	8.7–69.4	12.3	6.2–49.2
Endometritis	15.7	7.9–63.0	18.2	9.1–73.0	17.5	8.8–70.0	12.5	6.2–49.9
Well (no adverse outcome)	11.4	5.7–17.0	13.9	6.9–20.8	13.1	6.6–19.7	8.1	4.0–12.1
Infant								
Death	52.2	0–1,000.0	52.2	0–1,000.0	52.2	0–1,000.0	52.2	0–1,000.0
Cerebral palsy	n/a	n/a	688.4	344.2–2,753.6	688.4	344.2–2,753.6	688.4	344.2–2,753.6
HIE	40.9	20.4–163.5	40.9	20.4–163.5	40.9	20.4–163.5	40.9	20.4–163.5
Sepsis	8.6	4.3–34.2	8.6	4.3–34.2	8.6	4.3–34.2	8.6	4.3–34.2
RDS	25.5	12.7–101.9	25.9	12.9–103.5	25.5	12.7–101.9	25.5	12.7–101.9
TTN	8.7	4.3–34.6	9.1	4.5–36.3	8.7	4.3–34.6	8.7	4.3–34.6
Acidemia	7.3	3.7–29.3	7.7	3.9–30.9	7.3	3.7–29.3	7.3	3.7–29.3
Well (no adverse outcome)	0.9	0.5–1.3	0.9	0.5–1.4	0.9	0.4–1.3	0.9	0.4–1.3

Currency in dollars (\$thousands). Abbreviations: ERCD, elective repeat cesarean delivery; HIE, hypoxic ischemic encephalopathy; n/a, not applicable; RDS, respiratory distress syndrome; TOLAC, trial of labor after a previous cesarean; TTN, transient tachypnea of the newborn.

**Table 3** Utility and QALY Estimates by Mode of Delivery or Outcome

	Disutility	Disutility days		QALY		
Mode of delivery/outcome	Baseline	Baseline	Range	Baseline	Range	Reference
ERCD	0.45	21	14–180	27.140	26.944–27.149	<sup>3</sup>
Uterine rupture	0.49	21	14–180	27.138	26.925–27.147	<sup>3</sup>
Failed TOLAC <sup>a</sup>	0.47	21	14–180	27.139	26.934–27.148	<sup>3</sup>
Successful TOLAC	0.35	7	2–42	27.160	27.126–27.164	<sup>3</sup>
Hysterectomy <sup>b</sup>	0.49	21	14–180	24.355	22.724–25.894	<sup>3,25</sup>
Cerebral palsy <sup>c</sup>	0.44	All	All	14.840	10.336–19.611	<sup>26</sup>
HIE	0.75	42	14–180	30.824	30.787–30.901	Assumed
Infant <sup>d</sup>	n/a	n/a	n/a	30.910	7.728–23.183	Assumed

Abbreviations: ERCD, elective repeat cesarean delivery; HIE, hypoxic ischemic encephalopathy; n/a, not applicable; QALY, quality-adjusted life-years; TOLAC, trial of labor after a previous cesarean.

<sup>a</sup>Extrapolated from Chung et al, midway between ERCD and rupture.<sup>3</sup>

<sup>b</sup>Blend of Harris et al and Chung et al at 55% and 45%, respectively, to represent the proportion of women with a hysterectomy who would and would not have desired another pregnancy.<sup>3,25</sup> For Chung et al, assumed the disutility and disutility days in the table, and from Harris et al, disutilities of 0.31 (0.14–0.48) until age 50.

<sup>c</sup>Disutility range for cerebral palsy of 0.26–0.61.

<sup>d</sup>Baseline utility of 1 for all infants without cerebral palsy or HIE, with a range of (0.25–1) tested in sensitivity analysis.

Disutilities or utility decrements were assigned based on the literature (►Table 3).<sup>3,25,26</sup> Because information was limited, infants were assigned full utility (1) except in the case of infant death, CP, and HIE where disutilities of 0, 0.44, and 0.75, respectively, were assigned. QALYs were determined based on the disutilities and life expectancy, discounted at 3% in the base case. It was assumed mode of delivery per se did not alter maternal or neonatal life expectancy. Maternal and infant life expectancy was estimated to be 78 years except for in the case of CP where 50 years was assumed.<sup>21,27</sup>

To test the robustness of the results obtained from the base-case model, sensitivity analyses were performed. One-way sensitivity analysis was conducted on all probabilities, costs, and QALYs by varying one variable at a time from the low to high value in its range while holding other variables fixed. Multivariable sensitivity analysis also was conducted by varying more than one probability at a time. This included bivariable as well as probabilistic sensitivity analysis using Monte Carlo simulation with 10,000 iterations to determine how often the base-case strategy was preferred. Simulation was conducted using the  $\beta$  or uniform distribution for the probabilities where appropriate and the gamma distribution for costs. Sensitivity analysis was also conducted on the discount rate, using 0%, 5%, and 7%.

## Results

The base-case analysis revealed that, for a hypothetical cohort of 100,000 women, the choice of TOLAC resulted in 68,077 fewer cesarean deliveries, 201 fewer hysterectomies, and 25 fewer maternal deaths (►Table 4). Conversely, TOLAC was associated with 779 additional uterine ruptures, as well as adverse neonatal outcomes of sepsis, RDS, and acidemia.

Additionally, among those undergoing TOLAC, CP was estimated to occur in an additional six offspring. The TOLAC strategy, therefore, was dominant and resulted in \$138.6 million saved and 1703 QALYs gained per 100,000 women.


One-way sensitivity analysis was performed across the full range for all the variables. The results were robust to all changes except for five variables (►Fig. 1). These variables and their thresholds, or where the preferred strategy changed, were the probability of uterine rupture (4.2%), the

**Table 4** Maternal and Infant Outcomes per 100,000 Women

	TOLAC	ERCD
Deliveries	100,000	100,000
Cesarean deliveries	31,923	100,000
Maternal		
Uterine rupture	779	0
Maternal death	0	25
Hysterectomy	76	277
Endometritis	3492	2111
Infant		
Infant death	51	25
Cerebral palsy	6	0
HIE	51	0
Sepsis	5227	3115
RDS	679	528
Acidemia	302	226

Abbreviations: ERCD, elective repeat cesarean delivery; HIE, hypoxic ischemic encephalopathy; RDS, respiratory distress syndrome; TOLAC, trial of labor after a previous cesarean.





**Fig. 1** Tornado diagram of five variables with thresholds. Abbreviations: ERCD, elective repeat cesarean delivery; QALY, quality-adjusted life-years; TOLAC, trial of labor after a previous cesarean.

probability of successful TOLAC (42.2%), and without any complications the cost of ERCD (\$9040.00) and the cost of successful TOLAC (\$11,428.00) as well as the QALY of failed TOLAC (27.00), which represented a disutility of 0.47 for 132 days or more. Bivariable analysis on the probability of uterine rupture and successful TOLAC indicated that when the probability of uterine rupture was at 0%, the TOLAC strategy was preferred if the probability of success was 36.6% or more. When the probability of uterine rupture was at the base value, 0.8%, the TOLAC strategy was preferred as long as the probability of success was 42.6% or more. With the uterine rupture rate set at 1.5% and 3.0%, the probability of success had to be 47.2% and 58.4% or less, respectively, for the preferred strategy to change to an ERCD. When the probability of success was at the base value, 68.5%, TOLAC was preferred when the probability of rupture was 4.2% or less. When the probability of TOLAC success was 36.0% or less, ERCD was preferred over the entire range of the probability of uterine rupture (0 to 5.0%).

Monte Carlo simulation of the five sensitive variables at cost-effectiveness thresholds of \$25, \$50, and \$100 thousand found TOLAC to be preferred 91.2%, 91.9%, and 91.1% of the time, respectively.

## Discussion

Under base-case assumptions, after one cesarean with a low transverse incision, TOLAC was the most cost-effective strategy and would save approximately \$138.6 million per 100,000 women when compared with ERCD. This analysis improves upon prior analyses in several ways. First, we utilized an observational study specifically conducted to answer questions related to modes of delivery after a previous cesarean to obtain maternal and perinatal outcome probabilities. Second, the probabilities used for the decision analytic model were derived using propensity scores. This allowed us to uniquely develop TOLAC and ERCD groups with well-balanced baseline covariates with minimal bias. Moreover, whereas previous studies relied on cost data that were derived from a single institution, the cost data for this analysis were obtained from U.S. national sources, the AHRQ and the AMA.

Of the two economic reports noted to have the most methodological rigor in the 2003 AHRQ technology report,

the present analysis is most comparable to the study by Chung et al, because both consider outcomes of the current pregnancy.<sup>3</sup> A main criticism of the Chung et al work was that minimal sensitivity analysis was used.<sup>1</sup> The present analysis demonstrates that although TOLAC is cost-effective under many circumstances, this conclusion is highly dependent upon several key variables which, if altered sufficiently, result in the alternate strategy of ERCD being preferred.

Indeed, this finding reveals that the cost-effectiveness of TOLAC depends upon the characteristics of women who choose to attempt a vaginal birth. For example, TOLAC will no longer be cost-effective when the chance of VBAC success is low. One-way sensitivity analyses found TOLAC not cost-effective when the probability of successful TOLAC was below 42%.

Limitations of this analysis should be noted. For feasibility and clarity, the maternal and infant outcome probabilities were based on a hierarchy and therefore no more than one complication could be experienced by an individual. However, the effect this would have on the cost-effectiveness results would be *de minimus* at best because less than 0.03% and 2.5% of the mothers and neonates, respectively, experienced more than one outcome. In addition, this study did not include the potential long-term maternal outcomes of fecal and urinary incontinence due to the fact that the marginal increase in these outcomes due to TOLAC is not well known.<sup>1,28,29</sup> The 2003 AHRQ Evidence report specifically criticized Chung et al for including incontinence because no conclusive evidence linking the probability of incontinence outcomes to delivery approach could be found.<sup>1</sup> Unfortunately, conclusive evidence is still lacking and consequently they were not incorporated in the present model.

This analysis also excluded the important long-term outcomes of placenta accreta and previa in subsequent pregnancies. The determination of the exact cost-effectiveness over the life course would require a separate, more complex model taking into account uncertainties of future reproduction. Nevertheless, including these data inputs would only make the cost effectiveness of TOLAC greater. Additionally, the base case of this analysis did not include women undergoing labor induction and instead was predicated on women in spontaneous labor and included probabilities of success and rupture consistent with this type of labor. The sensitivity analysis, however, allows insight into whether induction would be cost-effective as well. The 2010 AHRQ evidence report estimated that the frequency of rupture for those induced at any gestational age was approximately 1.5%.<sup>30</sup> Even at this frequency, the preferred strategy changes to an ERCD only when the probability of success was approximately less than 47%. Most women induced, and particularly those with a favorable cervix, would be expected to have a chance of success greater than that threshold.

In conclusion, using an analytic framework designed to minimize bias, we found that a trial of labor after one previous low transverse cesarean is more cost-effective than an ERCD under a wide range of circumstances. This conclusion is strengthened in particular for women who undergo a spontaneous TOLAC with a high chance of success.

## Acknowledgments

The authors thank the following core committee network members who participated in protocol development and coordination between clinical research centers (Francee Johnson, BSN, Julia Gold BSN/APN), data management (Sandra Meadows), protocol/data management and statistical analysis (Elizabeth Thom, PhD, and John C. Hauth, MD), and manuscript review (Yasser El-Sayed, MD).

In addition to the authors, other members of the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network are as follows:

*The George Washington University Biostatistics Center*—E. Thom, H. Juliussen-Stevenson, M. Fischer, L. Leuchtenburg

*Northwestern University*—A. Peaceman, M. Socol, D. Gradishar, G. Mallett

*The Ohio State University*—J. Iams, F. Johnson, S. Meadows, H. Walker

*University of Alabama at Birmingham*—J. Hauth, A. Northen, S. Tate

*University of Texas Southwestern Medical Center*—S. Bloom, J. Gold, D. Bradford

*University of Utah*—M. Belfort (Utah Valley Regional Medical Center), F. Porter (Intermountain Healthcare), B. Oshiro (McKay-Dee Hospital Center), K. Anderson (University of Utah Health Sciences Center), A. Guzman (McKay-Dee Hospital Center)

*University of Pittsburgh*—S. Caritis, K. Lain, M. Cotroneo, D. Fischer, M. Luce

*Wake Forest University Health Sciences*—P. Meis, M. Harper, M. Swain, C. Moorefield, K. Lanier, L. Steele

*Thomas Jefferson University*—A. Sciscione, M. DiVito, M. Talucci, M. Pollock

*Wayne State University*—M. Dombrowski, G. Norman, A. Millinder, C. Sudz, B. Steffy

*University of Cincinnati*—M. Miodovnik, T. Siddiqi, H. How, N. Elder

*Columbia University*—M. Miodovnik, F. Malone, M. D'Alton, V. Pemberton, V. Carmona, H. Husami

*Brown University*—M. Carpenter, H. Silver, J. Tillinghast, D. Catlow, D. Allard

*University of Miami*—G. Burkett, J. Gilles, J. Potter, F. Doyle, S. Chandler

*University of Tennessee*—W. Mabie, R. Ramsey

*University of Texas at San Antonio*—O. Langer, D. Conway, S. Barker, M. Rodriguez

*University of North Carolina*—K. Moise, K. Dorman, S. Brody, J. Mitchell

*The University of Texas Health Science Center at Houston*—L. Gilstrap, M. Day, M. Kerr, E. Gildersleeve

*Case Western Reserve University-MetroHealth Medical Center*—P. Catalano, C. Milluzzi, B. Slivers, C. Santori

*University of Chicago*—A. Moawad, J. Hibbard, P. Jones, M. Ramos-Brinson, M. Moran, D. Scott

*Eunice Kennedy Shriver National Institute of Child Health and Human Development*—D. McNellis, K. Howell, S. Tolivaisa

MFMU Steering Committee Chair (*Vanderbilt University Medical Center*)—S. Gabbe

## Disclosure

The project described was supported by grants from the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development (NICHD) of the National Institutes of Health (NIH) [HD21410, HD21414, HD27860, HD27861, HD27869, HD27905, HD27915, HD27917, HD34116, HD34122, HD34136, HD34208, HD34210, HD40500, HD40485, HD40544, HD40545, HD40560, HD40512, and HD36801] and its contents are solely the responsibility of the authors and do not necessarily represent the official view of NICHD or the NIH.

Presented at the 32nd Annual Meeting of the Society for Maternal-Fetal Medicine. February 6 to 11, 2012. Dallas, Texas.

## References

- 1 Guise J-M, McDonagh M, Hashima J, et al. Vaginal birth after cesarean (VBAC). Evidence report/technology assessment no. 71 (prepared by the Oregon Health & Science University Evidence-based Practice Center under Contract no. 290-97-0018). AHRQ Publication No. 03-E018. Rockville, MD: Agency for Healthcare Research and Quality; 2003
- 2 Grobman WA, Peaceman AM, Socol ML. Cost-effectiveness of elective cesarean delivery after one prior low transverse cesarean. *Obstet Gynecol* 2000;95:745-751
- 3 Chung A, Macario A, El-Sayed YY, Riley ET, Duncan B, Druzin ML. Cost-effectiveness of a trial of labor after previous cesarean. *Obstet Gynecol* 2001;97:932-941
- 4 Macones GA, Peipert J, Nelson DB, et al. Maternal complications with vaginal birth after cesarean delivery: a multicenter study. *Am J Obstet Gynecol* 2005;193:1656-1662
- 5 Landon MB, Hauth JC, Leveno KJ, et al; National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network. Maternal and perinatal outcomes associated with a trial of labor after prior cesarean delivery. *N Engl J Med* 2004;351:2581-2589
- 6 McMahon MJ, Luther ER, Bowes WA Jr, Olshan AF. Comparison of a trial of labor with an elective second cesarean section. *N Engl J Med* 1996;335:689-695
- 7 Flamm BL, Goings JR, Liu Y, Wolde-Tsadiq G. Elective repeat cesarean delivery versus trial of labor: a prospective multicenter study. *Obstet Gynecol* 1994;83:927-932
- 8 Gold MR, Siegel JE, Russell LB, Weinstein MC; for the Panel on Cost-Effectiveness in Health and Medicine. Cost-Effectiveness in Health and Medicine. New York, NY: Oxford University Press; 1996:6
- 9 Spong CY, Landon MB, Gilbert S, et al; National Institute of Child Health and Human Development (NICHD) Maternal-Fetal Medicine Units (MFMU) Network. Risk of uterine rupture and adverse perinatal outcome at term after cesarean delivery. *Obstet Gynecol* 2007;110:801-807
- 10 American College of Obstetricians and Gynecologists. ACOG Committee Opinion No. 394, December 2007. Cesarean delivery on maternal request. *Obstet Gynecol* 2007;110:1501
- 11 ACOG Committee on Practice Bulletins—Obstetrics. ACOG Practice Bulletin No. 107: Induction of labor. *Obstet Gynecol* 2009;114(2 Pt 1):386-397

- 12 Grobman WA, Gilbert S, Landon MB, et al. Outcomes of induction of labor after one prior cesarean. *Obstet Gynecol* 2007;109(2 Pt 1): 262–269
- 13 Rosenbaum PR, Rubin DB. The central role of the propensity score in observational studies for causal effects. *Biometrika* 1983;70:41–55
- 14 Gilbert SA, Grobman WA, Landon MB, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network. Elective repeat cesarean delivery compared with spontaneous trial of labor after a prior cesarean delivery: a propensity score analysis. *Am J Obstet Gynecol* 2012;206:e1–e9
- 15 Badawi N, Felix JF, Kurinczuk JJ, et al. Cerebral palsy following term newborn encephalopathy: a population-based study. *Dev Med Child Neurol* 2005;47:293–298
- 16 Casella G. Refining binomial confidence intervals. *Can J Stat* 1986;14:113–129
- 17 U.S. Department of Health and Human Services. HCUPnet, Healthcare Cost and Utilization Project. Agency for Healthcare Research and Quality, Rockville, MD. Available at: <http://hcupnet.ahrq.gov/>. Accessed July 7, 2011
- 18 American Medical Association. Available at: [http://catalog.ama-assn.org/Catalog/cpt/cpt\\_search.jsp](http://catalog.ama-assn.org/Catalog/cpt/cpt_search.jsp). Accessed September 28, 2010
- 19 United States Department of Labor, Bureau of Labor Statistics. Available at: <http://www.bls.gov/cps/cpswktabs.htm>. Accessed November 6, 2010
- 20 Centers for Disease Control and Prevention (CDC). Economic costs associated with mental retardation, cerebral palsy, hearing loss, and vision impairment—United States, 2003. *MMWR Morb Mortal Wkly Rep* 2004;53:57–59
- 21 Katz RT. Life expectancy for children with cerebral palsy and mental retardation: implications for life care planning. *Neuro-Rehabilitation* 2003;18:261–270
- 22 Cerebral Palsy Source. Available at: [http://www.cerebralpalsy-source.com/About\\_CP/life\\_cp/index.html](http://www.cerebralpalsy-source.com/About_CP/life_cp/index.html). Accessed October 25, 2009
- 23 Haddox AC, Teutsch SM, Corso PS. *Prevention Effectiveness a Guide to Decision Analysis and Economic Evaluation*. New York, NY: Oxford University Press; 2003:115
- 24 United States Department of Labor, Bureau of Labor Statistics. Consumer Price Index. Available at: <http://www.bls.gov/cpi/>. Accessed July 7, 2011
- 25 Harris RA, Washington AE, Nease RF Jr, Kuppermann M. Cost utility of prenatal diagnosis and the risk-based threshold. *Lancet* 2004;363:276–282
- 26 Carroll AE, Downs SM. Comprehensive cost-utility analysis of newborn screening strategies. *Pediatrics* 2006;117(5 Pt 2): S287–S295
- 27 Arias E. *United States Life Tables, 2006*. National Vital Statistics Reports. Vol. 58, No. 21. Hyattsville, MD: National Center for Health Statistics; 2010
- 28 NIH State-of-the-Science Conference Statement on cesarean delivery on maternal request. *NIH Consens State Sci Statements* 2006;23:1–29
- 29 Vintzileos AM, Beazoglou T. Design, execution, interpretation, and reporting of economic evaluation studies in obstetrics. *Am J Obstet Gynecol* 2004;191:1070–1076
- 30 Guise JM, Eden K, Emeis C, et al. Vaginal birth after cesarean: new insights. Evidence/report technology assessment no. 191. (Prepared by the Oregon Health and Science University Evidence-based Practice Center under contract no. 290–2007–10057-I. AHRQ publication no. 10–E003. Rockville, MD: Agency for Healthcare Research and Quality). 2010

## Appendix

**Table A1** Additional Detail on Maternal Cost Estimates

Maternal outcomes	Hospital	Obstetrician	Anesthesia	Subtotal	Maternal	Caregiver	Total
Uterine rupture		2,325	728		3,840	1,008	
Maternal death	20,000			23,053			27,901
Thromboembolism	11,218			14,271			19,119
Hysterectomy	12,486			15,539			20,387
Operative injury	8,496			11,549			16,397
Wound infection	10,184			13,237			18,085
Endometritis	10,348			13,401			18,249
Maternal no other morbidity	5,964			9,017			13,865
Failed TOLAC		2,325	728		3,840	1,008	
Maternal death	20,000			23,053			27,901
Thromboembolism	10,475			13,528			18,376
Hysterectomy	11,743			14,796			19,644
Operative injury	7,753			10,806			15,654
Wound infection	9,441			12,494			17,342
Endometritis	9,605			12,658			17,506
Maternal well discharge	5,221			8,274			13,122



**Table A1** (Continued)

Maternal outcomes	Hospital	Obstetrician	Anesthesia	Subtotal	Maternal	Caregiver	Total
Successful TOLAC		2,068	297		1,280	432	
Maternal death	20,000			22,365			24,077
Thromboembolism	9,260			11,625			13,337
Hysterectomy	10,528			12,893			14,605
Operative injury	6,538			8,903			10,615
Wound infection	8,226			10,591			12,303
Endometritis	8,390			10,755			12,467
Maternal well discharge	4,006			6,371			8,083
ERCD		2,222	342		3,840	1,008	
Maternal death	20,000			22,564			27,412
Thromboembolism	9,198			11,762			16,610
Hysterectomy	10,466			13,030			17,878
Operative injury	6,476			9,040			13,888
Wound infection	8,164			10,728			15,576
Endometritis	8,328			10,892			15,740
Maternal well discharge	3,944			6,508			11,356

Currency in dollars (\$). Hospital costs were obtained from HCUPnet,<sup>17</sup> *International Classification of Diseases, Ninth Revision* codes: ERCD, 669.71; rupture, 665.11; failed TOLAC, 660.61; successful TOLAC, 654.21; thromboembolism, 671.44; hysterectomy, 68.49 (procedure code); operative injury, 665.54; wound infection, 674.34; endometritis 670.02. Maternal well discharge represents the cost of the mode of delivery, whereas maternal outcomes represent the cost for the mode of delivery plus the cost of the outcome. For example, the cost of an ERCD was \$3,944 and the cost of thromboembolism was \$5,254, resulting in an ERCD with thromboembolism cost of \$9,198. Obstetrician costs were obtained from the 2010 American Medical Association Current Procedural Terminology codes adjusted to 2009 dollars.<sup>18</sup> ERCD Current Procedural Terminology code 59510; Rupture 59618; Failed TOLAC 59618; Successful TOLAC 59610. Anesthesia costs were obtained from Chung et al by mode of delivery and translated into 2009 dollars.<sup>3</sup> Maternal and caregiver postpartum opportunity costs were derived from the Bureau of Labor Statistics using the 2009 median hourly wage and salary averages for women 25 to 34 years old and for all sexes 16 years and older, respectively.<sup>19</sup> For ERCD, rupture, failed TOLAC maternal opportunity costs were included for 6 weeks at 40 hours per week at \$16 per hour whereas for caregiver costs were included for 7 days, 8 hours per day at \$18 per hour. For successful TOLAC, maternal opportunity costs were included for 2 weeks, 40 hours per week at \$16 per hour and caregiver costs were included for 3 days, 8 hours per day at \$18 per hour. Abbreviations: ERCD, elective repeat cesarean delivery; TOLAC, trial of labor after a previous cesarean.

**Table A2** Additional Detail on Infant Cost Estimates

Infant outcome	Hospital	Pediatrician	Total
Failed TOLAC/ERCD			
Neonatal death	50,000	2,151	52,151
CP (failed TOLAC only)	73,438	8,976	82,414
HIE	36,719	4,158	40,877
Sepsis	5,597	2,954	8,551
RDS	23,317	2,151	25,468
TTN	6,509	2,151	8,660
Acidemia	5,183	2,151	7,334
Infant well discharge	771	124	895
Uterine rupture			
Neonatal death	50,000	2,151	52,151

(Continued)

**Table A2** (Continued)

Infant outcome	Hospital	Pediatrician	Total
CP	73,438	8,976	82,414
HIE	36,719	4,158	40,877
Sepsis	5,597	2,954	8,551
RDS	23,317	2,552	25,869
TTN	6,509	2,552	9,061
Acidemia	5,183	2,552	7,735
Infant well discharge	771	156	927
Successful TOLAC			
Neonatal death	50,000	2,151	52,151
CP	73,438	8,976	82,414
HIE	36,719	4,158	40,877
Sepsis	5,597	2,954	8,551
RDS	23,317	2,151	25,468
TTN	6,509	2,151	8,660
Acidemia	5,183	2,151	7,334
Infant well discharge	771	91	862

Currency in dollars (\$). Hospital costs were obtained from HCUPnet,<sup>17</sup> *International Classification of Diseases, Ninth Revision* codes: HIE, 768.7; infection, 771.81; RDS, 769; TTN, 770.6; infant well 795 (DRG). Pediatrician costs based on length of stay from the Cesarean Registry and 2010 Current Procedural Terminology codes<sup>18</sup> adjusted to 2009 dollars; 99460, 99466, 99468, 99469; for no other infant morbidity, pediatrician coverage was assigned in the base-case as 4 days for rupture, 3 days for ERCD and failed TOLAC, 2 days for successful TOLAC. For ERCD, rupture, failed TOLAC, an additional day was added for the outcomes RDS, TTN, acidemia, and an additional 2 days for successful TOLAC. For all modes of delivery, infection, HIE, and CP were assumed to have a pediatrician coverage of 6, 9, and 21 days, respectively. CP costs through hospital discharge. Abbreviations: ERCD, elective repeat cesarean delivery; CP, cerebral palsy; HIE, hypoxic ischemic encephalopathy; RDS, respiratory distress syndrome; TOLAC, trial of labor after a previous cesarean; TTN, transient tachypnea of the newborn.