Category

Polymer-Supported Synthesis

Key words

phenols aryl halides O-arylation

diaryl ethers

Y.-P. ZHANG,* Y.-C. JIAO, Y.-S. YANG,* C.-L. LI (LANZHOU UNIVERSITY OF TECHNOLOGY, P. R. OF CHINA)

 $\label{ligand-Free Catalytic System for the Synthesis of Diarylethers over Cu_2O/Cu-CNTs as Heterogeneous Reusable Catalyst$

Tetrahedron Lett. 2013, 54, 6494-6497.

Cu₂O/Cu-CNTs Catalyzed the O-Arylation of Phenols with Aryl Halides

$$R^{1} \stackrel{\text{Cu}_{2}\text{O/Cu-CNTs}}{||||} + X = I, Br, CI$$

$$\frac{\text{Cu}_{2}\text{O/Cu-CNTs}}{\text{Cs}_{2}\text{CO}_{3} (1.0 \text{ equiv})} \\ \text{DMF}, 140 °C, 24 h}$$

$$\text{up to 97\% yield} \\ (20 \text{ examples})$$

Selected examples:

Significance: Cu₂O/Cu-Coated carbon nanotubes (Cu₂O/Cu-CNTs) catalyzed the O-arylation of phenols with aryl halides under ligand-free conditions to give the corresponding diaryl ethers in up to 97% yield (20 examples).

Comment: Cu₂O/Cu-CNTs were recovered by filtration and reused three times without significant loss of catalytic activity. Lee and co-workers have previously reported the preparation and characterization of Cu₂O/Cu-CNTs (*Scr. Mater.* **2008**, *58*, 1010).

 SYNFACTS Contributors: Yasuhiro Uozumi, Shiguang Pan

 Synfacts 2014, 10(1), 0106
 Published online: 13.12.2013

 DOI: 10.1055/s-0033-1340398; Reg-No.: Y13913SF