

Endoloop application for the removal of a self-expandable metallic stent (SEMS) in an esophagocolonic anastomotic stricture

Anastomotic strictures occur in 3%–46.2% of patients after colonic reconstruction of the esophagus [1]. Self-expanding metal stents (SEMS) are increasingly considered for refractory or complex benign strictures of the esophagus [2]. Migration is a common complication and the stent should be removed to avoid gastrointestinal complications.

A 54-year-old man was referred to our department for dysphagia following esophagectomy with colonic interposition to treat an esophageal adenocarcinoma. On esophagoscopy, a 5-mm wide and 3-cm long stricture, corresponding to the esophagocolonic anastomosis at 25 cm from the incisors, could not be traversed. After five dilation sessions at 2-week intervals the patient was still dysphagic and a fully covered stent (HanaroStent, 80 mm in length, 18 mm in diameter; MI Tech, Seoul, Korea) was positioned. The patient reported clinical improvement for 2 weeks but then the dysphagia recurred. Radiographic examination disclosed stent migration. Endoscopy confirmed its location at the distal part of the colonic segment, proximal to the cologastric anastomosis, and the persistence of the proximal anas-

tomotic stenosis. We decided to re-dilate the stenosis up to 15 mm (● **Video 1**), and mobilize the stent proximally and try to remove it using endoloops to reduce the stent diameter. With the stent positioned in the colonic segment, four detachable ligating devices (MAJ 254; Olympus, Tokyo, Japan) were applied. Because of the eversion of its distal edge and the risk of the stent getting caught in the tissue, it was rotated and then retrieved by utilizing the “lasso” stitch at the stent edge and pulling it against the endoscope. The whole assembly was subsequently removed under fluoroscopic control through the stricture without complications. The removal of migrated SEMS is technically challenging and different methods have been reported, including the use of endoloops [3–5]. To our knowledge this is the first video report of a stent retrieved from an esophagocolonic anastomosis.

Endoscopy_UCTN_Code_TTT_1AO_2AZ

Competing interests: None

**B. M. Gonçalves¹, A. Ferreira^{1,2,3},
A. C. Caetano^{1,2,3}, P. Bastos¹,
C. Rolanda^{1,2,3}**

¹ Department of Gastroenterology, Hospital Braga, Portugal

² Life and Health Sciences Research Institute (ICVS), School of Health Sciences, University of Minho, Braga, Portugal

³ ICVS/3Bs – PT Government Associate Laboratory, Braga/Guimarães, Portugal

References

- 1 Yasuda T, Shiozaki H. Esophageal reconstruction with colon tissue. *Surg Today* 2011; 41: 745–753
- 2 Siersema PD. Stenting for benign esophageal strictures. *Endoscopy* 2009; 41: 363–373
- 3 Seitz U, Thonke F, Bohnacker S et al. Endoscopic extraction of a covered esophageal Z-stent with the aid of endoloops. *Endoscopy* 1998; 30: S91
- 4 Molina-Infante J, Fernandez-Bermejo M, Perez-Gallardo B. Removal of a migrated covered metallic stent through an esophageal stricture, with multiple endoloops. *Endoscopy* 2010; 42: E268–E269
- 5 An HJ, Lee HY, Kim BW et al. Endoscopic removal of a migrated esophageal self-expandable metal stent after compression with detachable snares through an intact esophageal stent. *Gastrointest Endosc* 2010; 71: 205–207

Bibliography

DOI <http://dx.doi.org/10.1055/s-0033-1344132>
Endoscopy 2013; 45: E209
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

B. M. Gonçalves
Serviço de Gastroenterologia
Hospital de Braga
Sete Fontes – São Vitcor
4710-243 Braga
Portugal
Fax: +351 253027999
brunommgoncalves@gmail.com

Video 1

Removal of a migrated self-expandable metallic stent (SEMS) with endoloop application. The use of endoloops reduced the stent diameter and permitted its retrieval through an esophagocolonic anastomotic stricture.