

## Biliary stent migration presenting as a recurrent pilonidal abscess with underlying rectocutaneous fistula


A 53-year-old patient was admitted to the gastroenterology department for the management of a benign biliary stricture secondary to chronic alcoholic pancreatitis. He underwent sphincterotomy and placement of a 10Fr×10cm plastic biliary stent. Endoscopic follow-up 2 months later revealed spontaneous migration of the stent and no further intervention was performed.

The patient was admitted to the general surgery outpatient clinic 7 years later for the management of a recurrent pilonidal abscess. Excision of the abscess revealed a long fistulous track towards the coccyx with a plastic stent inside it (► Fig. 1 a, b); the findings indicated a rectocutaneous fistula secondary to biliary stent migration. In addition, computed tomography scan showed a 1-cm defect of the sacrococcygeal junction in contact with the rectum (► Fig. 2). Postoperatively, the patient was treated with antibiotics and vacuum-assisted closure therapy and went on to make an excellent recovery. His follow-up over 4 years was unremarkable.


Complications of stent placement are well known and include migration with fistula formation between a variety of organs, such as: duodenocolic fistula [1], duodenoscrotal fistula [2], enterosplenic fistula [3], enterocutaneous fistula [4], and colovaginal fistula [5]. However, to the best of our knowledge recurrent abscess secondary to a rectocutaneous fistula has not been described previously.

Endoscopy\_UCTN\_Code\_CPL\_1AK\_2AI

**Competing interests:** None



**Fig. 1** The patient was admitted for management of a recurrent pilonidal abscess. **a** Excision of the pilonidal abscess revealed a fistulous track (arrowhead). **b** A 10Fr×10cm plastic biliary stent was retrieved from the fistula.



**Fig. 2** Postoperative computed tomography scan disclosed a 1-cm bone defect of the sacrococcygeal junction (arrowhead) in contact with the posterior wall of the rectum. The asterisk indicates the site of excision of the abscess.

**G. Mavrogenis<sup>1</sup>, M. Lalot<sup>2</sup>, Y. Hoebeke<sup>3</sup>,  
P. Warzée<sup>1</sup>, P. Van Ende<sup>2</sup>, A. Sibille<sup>1</sup>**

<sup>1</sup> Department of Gastroenterology, Site Notre Dame, Grand Hôpital de Charleroi, Charleroi, Belgium

<sup>2</sup> Department of Anesthesiology, Site Notre Dame, Grand Hôpital de Charleroi, Charleroi, Belgium

<sup>3</sup> Department of General Surgery, Site Notre Dame, Grand Hôpital de Charleroi, Charleroi, Belgium

## References

- 1 *Ang BK, Wee SB, Kaushik SP* et al. Duodenal-colic fistula resulting from migration of a biliary stent: a case report. *Gastrointest Endosc* 1998; 48: 80–83
- 2 *Basile A, Macri' A, Lamberto S* et al. Duodenoscolic fistula secondary to retroperitoneal migration of an endoscopically placed plastic biliary stent. *Gastrointest Endosc* 2003; 57: 136–138
- 3 *Baccarani U, Risaliti A, Sainz-Barriga M* et al. Ileosplenic fistula and splenic abscesses caused by migration of biliary stents in a liver transplant recipient. *Gastrointest Endosc* 2003; 58: 811–813
- 4 *Karim A, Orbell JH, Bhatti K* et al. Biliary stent migration presenting as a recurrent abdominal wall abscess with underlying enterocutaneous fistula. *Gastrointest Endosc* 2006; 63: 874–876
- 5 *Blake AM, Monga N, Dunn EM*. Biliary stent causing colovaginal fistula: case report. *JLS* 2004; 8: 73–75

## Bibliography

**DOI** <http://dx.doi.org/10.1055/s-0033-1344413>  
*Endoscopy* 2013; 45: E301–E302  
© Georg Thieme Verlag KG  
Stuttgart · New York  
ISSN 0013-726X

## Corresponding author

**G. Mavrogenis, MD**  
Grand Hôpital de Charleroi  
Site Notre Dame  
3 Grand Rue  
Charleroi 6000  
Belgium  
Fax: +32-71-102779  
[mavrogenis@gmail.com](mailto:mavrogenis@gmail.com)