

Cellular mechanisms of ALS mutations – a loss or a gain of function?

C. Haass^{1,2}

¹ Adolf-Butenandt-Institute, Biochemistry,
Ludwig-Maximilians-University Munich

² German Center for Neurodegenerative Diseases (DZNE),
Munich

TDP-43 and FUS are genetically and pathologically associated with amyotrophic lateral sclerosis (ALS) and frontotemporal lobar degeneration (FTLD). I will summarize our recent advances in understanding the cellular mechanisms of disease causing mutations in FUS and TDP-43 and show evidence, which may suggest a loss of function component in ALS/FTLD. A loss of function appears to be the case for the FUS associated ALS causing mutations. Consistent evidence from several laboratories demonstrates that these mutations reduce nuclear transport by disturbing a PY-nuclear localization signal, which leads to a cytoplasmic accumulation of the mutant proteins. Additional stressors are then required to initiate aggregation probably via stress granules.

Much less is known about the mechanism of TDP-43 mutations. We have recently generated zebrafish TDP-43 loss of

function mutants. Homozygous loss of function mutations in zebrafish *tardbp* show no morphological phenotype due to compensation by a splice variant of *tardbpl* (Tar DNA binding protein of 43 kDa like), a second zebrafish orthologue of human *TARDBP*. *tardbp* and *tardbpl* double homozygous mutants show muscle degeneration, strongly reduced blood circulation and a dramatic mispatterning of intersomitic vessels, impaired spinal motor axon outgrowth, and early death. A quantitative proteomic approach identified a muscle specific protein to be upregulated in *tardbp* and *tardbpl* double homozygous mutants. Strikingly, the same protein is similarly increased in the frontal cortex of FTLD-TDP patients suggesting aberrant expression in vascular smooth muscle cells. Thus, these findings reveal an unexpected role of TDP-43 in vascular patterning and muscle maintenance. Evidence will be presented that TDP-43 mutations only partially rescue the vascular phenotype.

Conflict of Interest: The author declares no conflict of interest.

Bibliography

DOI <http://dx.doi.org/10.1055/s-0033-1346714>
Drug Res 2013; 63, Suppl. 1: S17–S17

© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 2194-9379