

Peroral endoscopic myotomy (POEM) for achalasia after Roux-en-Y gastric bypass




Fig. 1 Timed barium esophagram showing dilated distal esophagus with delayed emptying of esophageal contents, and tapering at the lower esophageal sphincter (LES).




Fig. 2 High-resolution esophageal manometry demonstrating esophageal aperistalsis, and impaired lower esophageal sphincter (LES) relaxation.

Achalasia is a relatively rare esophageal motility disorder which has been reported to have an incidence of 1% in the morbidly obese population [1,2]. In line with the increase in number of severely obese individuals and concomitant rise in weight-loss procedures, post-bariatric surgery patients with symptomatic achalasia pose a surgical challenge. We present a patient with achalasia and Roux-en-Y gastric bypass (RYGB) successfully treated with peroral endoscopic myotomy (POEM).

A 64-year-old woman with RYGB presented with a 2-year history of progressive dysphagia, regurgitation, and a baseline Eckardt score of 4. Esophagogastroduodenoscopy (EGD) revealed a dilated esophagus, a gastric pouch without lesions, and a wide gastrojejunostomy. Esophagography showed delayed esophageal emptying and narrowing at the lower esophageal sphincter (LES) (Fig. 1). High resolution manometry demonstrated esophageal aperistalsis and impaired LES relaxation (Fig. 2). We performed POEM as previously described [3,4], with selective



Fig. 3 Peroral endoscopic myotomy (POEM).
a Mucosal incision site in the mid-esophagus.
b Endoscopic view of submucosal tunnel.
c Selective myotomy of circular muscle fibers.

myotomy of the circular muscle fibers and advancing 4 cm into the gastric pouch (total myotomy length of 14 cm) (Fig. 3). Esophagography on postoperative day 1 revealed no leak of oral contrast with adequate passage into the gastric pouch and beyond the gastrojejunostomy (Fig. 4). The patient's dysphagia and regurgitation had resolved at follow-up 5 weeks postoperatively, with an Eckardt score of 0. Laparoscopic Heller's myotomy (LHM) and pneumatic balloon dilation are historically considered the standard of care for achalasia [5]. However, management of post-bariatric surgery patients with achalasia is often complicated by exten-




Fig. 4 Barium esophagram on postoperative day 1 revealing satisfactory emptying of oral contrast into the gastric pouch and beyond the gastrojejunostomy.

sive adhesions and altered anatomy [6,7]. The potential advantages of POEM include minimal invasiveness, allowance of a longer esophageal myotomy if required, and that it can be more easily revisited without requiring open surgical access. In this study, we demonstrate that POEM was feasible, safe, and effective for symptomatic relief in an achalasia patient with RYGB. POEM should be included as a potential treatment option for achalasia patients with previous bariatric surgeries.

Endoscopy_UCTN_Code_TTT_1AO_2AN

Competing interests: None of the authors have relevant disclosures or conflict of interest to declare related to this study.

Dennis Yang, Peter V. Draganov

Division of Gastroenterology, University of Florida, Gainesville, Florida

References

- 1 Koppman JS, Poggi L, Szomstein S et al. Esophageal motility disorders in the morbidly obese population. *Surg Endosc* 2007; 21: 761–764
- 2 Almog G, Anthone GJ, Crookes PF. Achalasia in the context of morbid obesity: a rare but important association. *Obes Surg* 2003; 13: 896–900
- 3 Inoue H, Minami H, Kobayashi Y et al. Peroral endoscopic myotomy (POEM) for esophageal achalasia. *Endoscopy* 2010; 42: 265–271
- 4 Yang D, Wagh MS. Peroral endoscopic myotomy (POEM) for the treatment of achalasia: an analysis. *Diagn Therapeut Endosc* 2013. doi: 10.1155/2013/389596 [Epub ahead of print]
- 5 Boeckstaens GE, Annese V, des Varannes SB et al. Pneumatic dilation versus laparoscopic Heller's myotomy for idiopathic achalasia. *N Engl J Med* 2011; 364: 1807–1816
- 6 Ramos AC, Murakami A, Lanzarini EG et al. Achalasia and laparoscopic gastric bypass. *Surg Obes Relat Dis* 2009; 5: 132–134
- 7 Chapman R, Rotundo A, Carter N et al. Laparoscopic Heller's myotomy for achalasia after gastric bypass: A case report. *Int J Surg Case Rep* 2013; 4: 396–398

Bibliography

DOI <http://dx.doi.org/10.1055/s-0033-1359140>
Endoscopy 2014; 46: E11–E12
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Peter V. Draganov, MD
 Division of Gastroenterology
 University of Florida
 1600 SW Archer Road, Room HD 602
 Gainesville, FL 32610
 USA
 Fax: +1-352-392-3618
peter.draganov@medicine.ufl.edu