



## A case of delayed bleeding 9 days after endoscopic ultrasound-guided pancreatic pseudocyst drainage



**Fig. 1** Computed tomography scan showed a 68-mm multilocular pancreatic pseudocyst (arrow) in contact with the stomach and pancreas.



**Fig. 2** Puncture of the pancreatic pseudocyst. **a** The color Doppler function confirmed the absence of intervening blood vessels in the puncture line. **b** The pseudocyst was punctured with a 19-G needle under endoscopic ultrasound guidance.




Endoscopic ultrasound-guided pancreatic pseudocyst drainage (EUS-PPD) has become a safe and effective procedure for pancreatic pseudocysts. However, post-procedural complication rates of 8.5%–20% have been reported [1–4]. For delayed complications, only one case of bleeding, which occurred 23 days after EUS-PPD from the transjejunal approach, has been reported [5]. We report a case of bleeding 9 days after EUS-PPD from the transgastric approach.

A 79-year-old man was admitted to our hospital for the treatment of biliary pancreatitis. Heparin was administered because of a history of myocardial infarction. A bile duct stone was successfully removed, but pseudocyst formation occurred (● Fig. 1). After 36 days, pseudocyst infection was suspected and EUS-PPD was performed. Heparin was stopped only on the day of the EUS-PPD procedure. Before puncturing the pseudocyst, the absence of intervening blood vessels was confirmed using the color Doppler function (● Fig. 2a). The pseudocyst was punctured with a 19-G needle (Sonotip; Medi-Globe GmbH, Rosenheim, Germany) under EUS guidance (● Fig. 2b). Then, an electrical cautery needle (Cyst-Gastro set; ENDO-FLEX GmbH, Voerde, Germany) was used to dilate the gastric and cystic wall. Finally, a 5-Fr nasocatheter was inserted into the pseudocyst. There were no procedure-related adverse events and the symptoms disappeared. Although the serum hemoglobin level did not drop markedly (Day 1: 12.0 mg/dL; Day 8: 11.7 mg/dL), hematemesis was observed 9 days after EUS-PPD and the hemoglobin level dropped to 6.9 mg/dL. Emergency endoscopy showed abundant clots around the pseudocyst fistula (● Fig. 3a, b). Angiography showed no aneurysms or contrast medium extravasation (● Fig. 4).


The late bleeding was possibly caused by the physical stimulus from the inserted stent which caused intracystic mural vessel destruction; the antithrombotic treatment might have been a contributing factor. To our knowledge, this is the first case of delayed bleeding 9 days after EUS-PPD. Sufficient long-term care must be provided after EUS-PPD, particularly in patients receiving antithrombotic treatment.

Endoscopy\_UCTN\_Code\_CPL\_1AK\_2AG

**Competing interests:** None



**Fig. 3** Emergency endoscopy findings. **a, b** Abundant clots (arrow) were found around the pseudocyst fistula.



**Fig. 4** Angiography showed no aneurysms or contrast medium extravasation.

**Tomoaki Matsumori, Akio Katanuma, Hiroyuki Maguchi, Kuniyuki Takahashi, Manabu Osanai, Kei Yane, Toshifumi Kin, Ryo Takaki, Kazuyuki Matsumoto, Katsushige Gon, Akiko Tomonari**

Center for Gastroenterology,  
Teine-Keijinkai Hospital, Hokkaido, Japan

#### References

- 1 *Varadarajulu S, Bang JY, Phadnis MA et al.* Endoscopic transmural drainage of peripancreatic fluid collections: outcomes and predictors of treatment in 2011 consecutive patients. *J Gastrointest Surg* 2011; 15: 2080–2088
- 2 *Park DH, Lee SS, Moon SH et al.* Endoscopic ultrasound-guided versus conventional
- 3 *Künzli HT, Timmer R, Schwartz MP et al.* Endoscopic ultrasonography-guided drainage is an effective and relatively safe treatment for peripancreatic fluid collections in a cohort of 108 symptomatic patients. *Eur J Gastroenterol Hepatol* 2013; 25: 958–963
- 4 *Puri R, Mishra SR, Thandassery RB et al.* Outcome and complications of endoscopic ultrasound guided pancreatic pseudocyst drainage using combined endoprosthesis and naso-cystic drain. *J Gastroenterol Hepatol* 2012; 27: 722–727
- 5 *Itoi T, Itokawa F, Sofuni A et al.* Late bleeding after EUS-guided transjejunal drainage of a pancreatic pseudocyst in a Roux-en-Y patient. *Dig Endosc* 2011; 23 (Suppl. 01): 51–53

transmural drainage for pancreatic pseudocysts: a prospective randomized trial. *Endoscopy* 2009; 41: 842–848

#### Bibliography

**DOI** <http://dx.doi.org/10.1055/s-0034-1364881>  
*Endoscopy* 2014; 46: E245–E246  
 © Georg Thieme Verlag KG  
 Stuttgart · New York  
 ISSN 0013-726X

#### Corresponding author

**Tomoaki Matsumori, MD**  
 Center for Gastroenterology  
 Teine-Keijinkai Hospital  
 1-12-1-40 Maeda  
 Teine-ku, Sapporo 006-8555  
 Japan  
 Fax: +81-11-6852967  
 tom.matsu1979@gmail.com