

Esophageal duplication cyst causing megaesophagus in a young woman presenting with dysphagia



Fig. 1 Barium esophagram suggesting megaesophagus.




Fig. 2 Endoscopy image showing a bulge in the distal esophagus.




Fig. 3 Computed tomography showing a hypodense lesion located in the posterior mediastinum, adjacent to the left atrium and lower esophagus.



Fig. 4 Endoscopic ultrasound showing the submucosa and muscularis propria of the esophageal duplication cyst.

A 35-year-old woman had dysphagia to solids for 6 months. Barium esophagram suggested megaesophagus (▶ Fig. 1), and endoscopy (▶ Fig. 2) showed bulging of the distal esophagus, suggesting extrinsic compression. The patient had no epidemiologic background for Chagas disease. Computed tomography (CT) showed a hypodense lesion in the posterior mediastinum, adjacent to the left atrium and lower esophagus (▶ Fig. 3).

The patient was referred for endoscopic ultrasound (EUS), which showed an anechoic, homogeneous lesion measuring 49 mm in the distal esophagus, inside the esophageal wall (▶ Fig. 4). EUS-guided fine-needle aspiration (EUS-FNA) of the cyst (▶ Fig. 5) obtained a mucoid fluid and small tissue fragments of the cystic wall. Histopathological examination showed cuboidal cells. The patient underwent thoracotomy with cyst excision from the wall of the lower esophagus with no complications. Histopathological examination showed evidence of two muscle layers and cuboidal cells, confirming the diagnosis of esophageal duplication cyst. After 1 year's follow-up, the patient is asymptomatic and the appearance on upper endoscopy is normal.

Esophageal duplication cysts are rare congenital anomalies, most frequently located in the right posterior mediastinum at the level of the distal third of the esophagus.




Fig. 5 Endoscopic ultrasound-guided fine-needle aspiration of the esophageal duplication cyst.

gus [1,2]. The cysts are lined by squamous, columnar, cuboidal, pseudostratified, or ciliated epithelium. They are covered by two thick muscle layers which are in contiguity with the muscularis propria. Because this condition is rare, it is sometimes not diagnosed or is confused with other diseases [3]. EUS can be helpful for preoperative diagnosis by demonstrating a cystic lesion with an intimate attachment to the esophageal wall and the presence of a smooth muscle coat.

In the present case we were able to demonstrate an esophageal duplication cyst causing megaesophagus clinically and radiologically.

Endoscopy_UCTN_Code_CCL_1AB_2AC_3AH

Competing interests: None

Augusto Carbonari, Marcelo Frota, Rogerio Colaiacovo, Lucio Rossini, Romeu Nakamura

Irmadade da Santa Casa de Misericordia de São Paulo – Department of Endoscopy and French-Brazilian Centre of Endoscopic Ultrasound (CFBEUS), São Paulo, São Paulo, Brazil

References

- 1 Cioffi U, Bonavina L, Simone M et al. Presentation and surgical management of bronchogenic and esophageal duplication cysts in adults. *Chest* 1998; 113: 1492–1496. doi: 10.1378/chest.113.6.1492
- 2 Procházka V, Kala Z, Novotný I et al. Laparoscopic management of an infected oesophageal cyst. *Rozhl Chir* 2005; 84: 526–528
- 3 Ringley C, Bochkarev V, Oleynikov D. Esophageal duplication cyst – a guest case in robotic and computer-assisted surgery from the University of Nebraska Medical Center. *MedGenMed* 2006; 8: 25

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1365288>
 Endoscopy 2014; 46: E201–E202
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

Corresponding author

Augusto Carbonari, MD

Irmadade da Santa Casa de Misericordia de São Paulo
 Department of Endoscopy and French-Brazilian Centre of Endoscopic Ultrasound (CFBEUS)
 Rua Manuel Figueiredo Landim 600 ap 52A
 São Paulo
 São Paulo 04693-130
 Brazil
 Fax: +55-19-996040645
augustocarbonari@gmail.com