

## Retroperitoneal schwannoma diagnosed by endoscopic ultrasound-guided fine-needle aspiration


A 59-year-old white man was referred for evaluation of epigastric pain that was radiating to his back and a 5-kg weight loss over 6 months. A computed tomography (CT) scan revealed a heterogeneous low-attenuation tumor, measuring 5 cm in size, adjacent to the celiac artery and the pancreatic body (► **Fig. 1**). Sectorial endoscopic ultrasound (GF-UCT140-AL5; Olympus America Inc., New York, USA), coupled to an ultrasound unit (Prosound alfa-5 SX; Aloka), detected a retroperitoneal well-circumscribed, hypoechoic mass, which measured 5×4 cm and was located immediately above the celiac tripod (► **Fig. 2**).

Endoscopic ultrasound-guided fine needle aspiration (EUS-FNA) was performed by three passes of a 22-gauge needle (EchoTip Ultra Echo-22; Cook Medical, Winston-Salem, North Carolina, USA). Histology revealed a spindle cell tumor that was strongly immunoreactive for S-100, thereby defining it as a schwannoma (► **Fig. 3**). The tumor was successfully removed despite its adherence to the celiac tripod arteries (► **Fig. 4**).


A schwannoma is a tumor arising from neural crest-derived Schwann cells in the sheaths of peripheral nerves. Retroperitoneal schwannomas are very rare, comprising less than 6% of all retroperitoneal tumors [1]. They are clinically asymptomatic until the tumors reach a large size and cause compressive symptoms. Diagnosis is difficult because of their rarity, asymptomatic course, and the lack of any specific diagnostic blood test or features on imaging studies [2]. Tumor size is related to its malignant potential and to the formation of cysts [3]. Surgery is the treatment of choice and is usually curative [4]. The findings on EUS generally reveal a well-circumscribed hypoechoic mass. EUS-FNA of retroperitoneal tumors is a valuable method for the preoperative diagnosis of schwannomas [2].

Endoscopy\_UCTN\_Code\_CCL\_1AF\_2AZ\_3AB


**Competing interests:** None




**Fig. 1** A computed tomography (CT) scan showing a heterogeneous low-attenuation tumor measuring 5 cm in size, adjacent to the celiac artery and the pancreatic body, that is displacing the splenic vessels anteriorly.



**Fig. 2** Linear-array endoscopic ultrasound (EUS) showing: **a** a retroperitoneal hypoechoic heterogeneous mass with well-defined borders measuring 5×4 cm; **b** the absence of a hyperechoic interface between the tumor and the celiac artery, and no evidence of thrombi or collateral circulation.



**Fig. 3** Histopathological findings of the cell block specimen showing: **a** large cohesive groups of spindle cells with nuclear palisading on hematoxylin and eosin (H&E) staining (original magnification  $\times 100$ ); **b** diffuse nuclear and cytoplasmic on immunostaining with polyclonal S-100 (original magnification  $\times 50$ ).



**Fig. 4** Macroscopic appearance of the cut surface of the resected tumor showing a firm, yellowish, 4.5-cm well-circumscribed, encapsulated solid tumor with focal areas of hemorrhage.

**César Vivian Lopes<sup>1</sup>, Manuela Zereu<sup>2</sup>, Roque Domingos Furian<sup>3</sup>, Bianca Canela Furian<sup>3</sup>, Tiago Auatt Paes Remonti<sup>4</sup>**

<sup>1</sup> Department of Gastroenterology and Digestive Endoscopy, Santa Casa Hospital, Porto Alegre, Brazil

<sup>2</sup> Department of Clinical Oncology, Santa Casa Hospital, Porto Alegre, Brazil

<sup>3</sup> Department of Pathology, Santa Casa Hospital, Porto Alegre, Brazil

<sup>4</sup> Department of Surgical Oncology, Santa Casa Hospital, Porto Alegre, Brazil

## References

- 1 Facciorusso D, Federici T, Giacobbe A et al. Retroperitoneal neurilemoma diagnosed by endosonographically guided fine needle aspiration. *J Clin Ultrasound* 2006; 34: 241–243
- 2 Kudo T, Kawakami H, Kuwatani M et al. Three cases of retroperitoneal schwannoma diagnosed EUS-FNA. *World J Gastroenterol* 2011; 17: 3459–3464
- 3 Moriya T, Kimura W, Hirai I et al. Pancreatic schwannoma. Case report and an updated 30-year review of the literature yielding 47 cases. *World J Gastroenterol* 2012; 18: 1538–1544
- 4 Li Q, Gao C, Juzi JT et al. Analysis of 82 cases of retroperitoneal Schwannoma. *ANZ J Surg* 2007; 77: 237–240

## Bibliography

**DOI** <http://dx.doi.org/10.1055/s-0034-1365816>  
Endoscopy 2014; 46: E287–E288  
© Georg Thieme Verlag KG  
Stuttgart · New York  
ISSN 0013-726X

## Corresponding author

**César Vivian Lopes, PhD**  
Rua Prof. Cristiano Fischer 668/1001  
Porto Alegre-RS  
91410-000  
Brazil  
Fax: +55-51-33388054  
[drcvlopes@gmail.com](mailto:drcvlopes@gmail.com)