




## Sequential endoscopic drainage and clip closure of an intrathoracic esophagogastric anastomotic dehiscence



**Fig. 1** Endoscopic ultrasound (EUS) view of the hypoechoic lesion in the submucosa of the distal esophagus.



**Fig. 2** Thoracic computed tomography (CT) scan showing a posterior mediastinal collection (arrow).



**Fig. 3** Endoscopic views showing: **a** the esophagogastric anastomosis with an 8-mm long dehiscence; **b** two double-pigtail plastic stents passing through the fistula orifice to drain the mediastinal collection; **c** the application of argon plasma coagulation at the fistula orifice.

A 69-year-old man underwent an upper gastrointestinal endoscopy, which revealed a large submucosal lesion in the distal esophagus with normal overlying mucosa. Endoscopic ultrasound (EUS) showed a hypoechoic lesion measuring 80×40×30 mm in the submucosa (● **Fig. 1**). A subtotal esophagectomy was performed. Histopathological examination of the resected tissue revealed a duplication cyst.

The patient developed sepsis 3 weeks after surgery. A computed tomography (CT) scan revealed evidence of dehiscence of the esophagogastric anastomosis with a large mediastinal collection (● **Fig. 2**). Upper gastrointestinal endoscopy confirmed the presence of a dehiscence affecting an 8-mm section of the esophagogastric anastomosis (● **Fig. 3 a**). Pus was aspirated through the fistula orifice and two double-pigtail plastic stents were placed endoscopically to drain the abscess


(● **Fig. 3 b**). There was rapid improvement in the patient's clinical condition.

After 1 week, the stents were removed and an over-the-scope clip (OTSC) was applied to close the fistula. Unfortunately, despite this, a Gastrografin swallow revealed persistence of the fistula.

A third upper gastrointestinal endoscopy was performed and after removing the OTSC, we applied argon plasma coagulation (APC) and three through-the-scope clips, which effectively sealed the fistula (● **Fig. 3 c**). Resolution of the mediastinal abscess as well as the fistula was confirmed on a further thoracic CT scan and Gastrografin swallow (● **Fig. 4**).

This case highlights the potentially useful role of endoscopic drainage and clipping devices in the management of a rare but serious adverse event of esophageal surgery.

Endoscopy\_UCTN\_Code\_TTT\_1AO\_2AI



**Fig. 4** Gastrografin swallow after placement of a metal clip confirming that the fistula orifice had been successfully closed.

**Competing interests:** None

**Samuel Costa<sup>1</sup>, Carlos Ferreira<sup>1</sup>,  
Rui Esteves<sup>2</sup>, Patricia Lages<sup>2</sup>,  
Paulo Costa<sup>2</sup>, Luis Ribeiro<sup>1</sup>, José Velosa<sup>1</sup>**

<sup>1</sup> Department of Gastroenterology and  
Hepatology, Hospital de Santa Maria,  
Lisbon, Portugal

<sup>2</sup> Department of Surgery 1, Hospital de  
Santa Maria, Lisbon, Portugal

#### **Bibliography**

**DOI** [http://dx.doi.org/  
10.1055/s-0034-1377352](http://dx.doi.org/10.1055/s-0034-1377352)  
Endoscopy 2014; 46: E371–E372  
© Georg Thieme Verlag KG  
Stuttgart · New York  
ISSN 0013-726X

#### **Corresponding author**

**Samuel Costa, MD**  
Hospital de Santa Maria  
Avenida Professor Egas Moniz  
1649-035 Lisboa  
Portugal  
samuelcosta@hotmail.com