


Y. KIMURA, N. FUKAWA, Y. MIYAUCHI, K. NOGUCHI, K. TANAKA\* (TOKYO INSTITUTE OF TECHNOLOGY, TOKYO UNIVERSITY OF AGRICULTURE AND TECHNOLOGY, AND JAPAN SCIENCE AND TECHNOLOGY AGENCY, SAITAMA, JAPAN)

Enantioselective Synthesis of [9]- and [11]Helicene-Like Molecules: Double Intramolecular [2+2+2] Cycloaddition  
*Angew. Chem. Int. Ed.* **2014**, *53*, 8480–8483.

## [2+2+2] to Twist: Synthesis of [9]- and [11]Helicene-Like Molecules



**Significance:** Higher order helicene-like molecules are difficult to synthesize enantioselectively due to steric constraints. The authors report the enantioselective synthesis of [9]- and [11]helicene-like molecules **4** and **6** via double intramolecular [2+2+2] cycloaddition of hexaynes, catalyzed by a cationic rhodium/chiral bis(phosphine) complex. Notably, molecules **4** and **6** both contain completely ortho-fused ring systems.

**Comment:** The authors report that the second cycloaddition is difficult to achieve because it proceeds through the highly sterically encumbered intermediate **3**. The diastereoselective synthesis of an [11]helicene-like molecule was reported previously (P. Sehnal et al. *Proc. Natl. Acad. Sci.* **2009**, *106*, 13169), but the reported molecule contained three para-fused rings.

**SYNFACTS Contributors:** Timothy M. Swager, Kathleen R. White  
Synfacts 2014, 10(9), 0926 Published online: 18.08.2014

**DOI:** 10.1055/s-0034-1378599; **Reg-No.:** S07914SF

2014 © THIEME STUTTGART • NEW YORK