Synlett

J. Qian et al.

Letter

A Facile and Mild Approach for Stereoselective Synthesis of α -Fluoro- α , β -unsaturated Esters from α -Fluoro- β -keto Esters via Deacylation

127

Jinlong Qian Wenbin Yi^{*} Meifang Lv Chun Cai

School of Chemical Engineering, Nanjing University of Science & Technology, Nanjing, Jiangsu 210094, P. R. of China yiwenbin@mail.njust.edu.cn B^{1} F $Cs_{2}CO_{3}$ H $Cs_{2}CO_{3}$ H



high yield and stereoselectiv
 simple material

Received: 02.08.2014 Accepted after revision: 09.10.2014 Published online: 05.11.2014 DOI: 10.1055/s-0034-1378917; Art ID: st-2014-w0647-1

Abstract The highly stereoselective olefination reaction of α -fluoro- β keto esters for the synthesis of α -fluoro- α , β -unsaturated esters has been developed. The olefination combines nucleophilic addition, intramolecular nucleophilic addition, and elimination in one step, as well as provides a facile synthetic approach to α -fluoro- α , β -unsaturated esters which are important units in many biologically active compounds and useful precursors in a variety of functional-group transformations.

Key words olefination, highly stereoselective, deacylation, fluoroolefins, carbon–carbon bond cleavage

Organofluorine compounds have experienced considerable growth in academic interest in recent years due to their growing importance in drug development purposes and crop protection.¹ In particular, α -fluoro- α , β -unsaturated esters are well known as precursors to biologically active compounds and have been successfully used to prepare a new generation of modified pheromones, herbicides, and medicines² (selected bioactive structures are shown in Figure 1). The traditional approaches for the preparation of these compounds are based on the Wittig,³ thia-Wittig,⁴ Horner-Wadsworth-Emmons (HWE),⁵ Peterson,⁶ or fluorous Julia⁷ olefination reactions (Scheme 1). Most of these procedures generally suffer from several major drawbacks including the requirement of metal catalysts,³ harsh reaction conditions,^{4,5} low selectivity,⁶ and the use of expensive or complex starting materials.⁵⁻⁸





Scheme 1 Traditional approach for the preparation of $\alpha\mbox{-fluoro-}\alpha\mbox{,}\beta\mbox{-unsaturated esters}$

F 18 examples

[.] Z/E up to 99:1



۸

Owing to the stability of carbon–carbon bonds, their cleavage has long remained a great challenge for organic chemists.⁹ Decarboxylation^{10b,c} and deacylation^{10a,d} are two types of the most prevailing methods to fulfill this purpose because of their efficiency informing reactive intermediates that successively promote the bond cleavage under mild conditions.¹⁰ The descriptions of this potentially useful and versatile molecule for the synthesis of α -functionalized α , β -unsaturated carbonyl compounds date back to 1978, in which Tsuboi's group reported the synthesis of 5,5,5-tri-chloro-3-penten-2-one by the reaction of chloral with 2,4-pentanedione via deacylation process.¹¹ In 2004, they continuously developed this method for the synthesis of α -chloro- α , β -unsaturated esters by the reaction of chlorinated ethyl acetoacetates with aldehydes.¹²

Along this line, we herein reported the first example of the synthesis of α -fluoro- α , β -unsaturated esters from α fluoro- β -keto esters and aldehydes through deacylation process (Scheme 2). This process successfully combines nu-



cleophilic addition, intramolecular nucleophilic addition, and elimination in one step. This protocol also provides a practical, simple, and mild synthetic approach to α -fluoro- α , β -unsaturated carbonyl compounds.

The starting material ethyl 2-fluoro-3-oxo-3-phenylpropanoate (1a) was easily prepared by stirring the corresponding β-keto ester with 1-chloromethyl-4-fluoro-1,4diazoniabicyclo[2.2.2]octane bis(tetrafluoroborate) (SelectfluorTM) according to the literature procedure.¹³ Our initial study on olefination study started with the reaction of **1a** and benzaldehyde (2a). A variety of parameters was summarized in Table 1. With regard to the influence of reaction temperature, it was found that the yield of the product 3aa increased from 56% at room temperature (Table 1, entry 1) to 80% at 40 °C (Table 1, entry 4) by using cesium carbonate as the base, whereas the yield had a significant reduction when the reaction conducted at 60 °C and 80 °C (Table 1, entries 2 and 3). Beside the cesium carbonate, other cesium salts (CsF and CsOAc) were tested, but only low efficiency were obtained (Table 1, entries 5 and 6). The reaction did not work in the presence of Na₂CO₃, NaOH, KOH, or KOt-Bu (Table 1, entries 7-13). Further studies indicated that the superior result was available by using acetonitrile compared with other solvents (Table 1, entries 4, 14-19). Based on the ¹H NMR data and comparison with the reported experimental data,^{3b,e} it was to our delight that a high ratio (up to 99:1) of Z stereoisomer was identified.

Syn lett

J. Qian et al.

 Table 1 Optimization of Reaction Conditions^a



^a Reaction conditions: **1a** (0.55 mmol), **2a** (0.5 mmol), base (1 mmol).

^b Isolated yields.

^c Relative ratio of the crude determined by ¹H NMR spectroscopy.

^d Reaction for 48 h. ^e GC yield based on **2a**.

With a set of optimized conditions in hand, the scope of α -fluoro- β -keto esters **1** and aldehydes **2** were investigated (Table 2).¹⁴ The reactions of α -fluoro- β -keto esters with aryl aldehydes bearing electron-withdrawing substituents (Table 2, entries 4-10) was more effective than electrondonating ones (Table 2, entries 2 and 3), and could be smoothly transformed into the desired products in excellent yields. Aromatic aldehydes with substituents at different positions of the aryl ring (para, meta, and ortho position) reacted well under the standard conditions (Table 2, entries 8-10). In addition, 1-naphthaldehyde, furfural, 2-thienaldehyde, and 2-pyridinecarboxaldehyde had good yields in this transformation, generating 3am, 3ak, 3ao and 3ap in 81%, 77%, 75%, and 93% yield, respectively (Table 2, entries 11-14). Alkyl aldehydes also worked well in high yields (Table 2, entries 15 and 16). α -Fluoro- β -keto esters derivates **1b**–**d** produced the corresponding α -fluoro- α , β -unsaturated esters in moderate to high yields (Table 2, entries 17–20), and indicated that electron-withdrawing substituents make deacylation proceed slightly more efficiently [NO₂/H/OMe = 87:80:63 (%)]. More economical α -fluoro- β -keto ester **1e** gave poor yields in the reaction (Table 2, entries 21, 22). It should be noteworthy that the *Z*/*E* ratios of this transformation are extremely high. X-ray crystal-structure analysis confirmed the structure and selectivity of product **3ag** (Figure 2).

Letter

Table 2 High Stereoselective Olefination Reactions of Different α -Fluoro- β -keto Esters **1** with Different Aldehydes **2**^a

R ¹		O R ³ H	Cs ₂ CO ₃	$\rightarrow R^{3}$	
	1	2			3
En- try	R ¹	R ²	R ³	Yield (%) [♭]	Z/E ^c
1	1a Ph	Et	2a Ph	3aa 80	96:4
2	1a Ph	Et	2b 4-MeC ₆ H ₄	3ab 50	93:7
3	1a Ph	Et	2c 4-Me ₃ OC ₆ H ₄	3ac 54	94:6
4	1a Ph	Et	2d 4-ClC ₆ H ₄	3ad 87	97:3
5	1a Ph	Et	2e 4-BrC ₆ H ₄	3ae 86	98:2
6	1a Ph	Et	2f 4-FC ₆ H ₄	3af 89	98:2
7	1a Ph	Et	2g 4-F ₃ CC ₆ H ₄	3ag 94	99:1
8	1a Ph	Et	2h 4-O ₂ NC ₆ H ₄	3ah 92	99:1
9	1a Ph	Et	2i 2-O ₂ NC ₆ H ₄	3ai 88	97:3
10	1a Ph	Et	2j 3-O ₂ NC ₆ H ₄	3aj 85	99:1
11	1a Ph	Et	2k 2-furfuryl	3ak 77	94:6
12	1a Ph	Et	2l 1-naphthyl	3al 81	96:4
13	1a Ph	Et	2m 2-thienyl	3am 75	94:6
14	1a Ph	Et	2n 2-pyridyl	3an 93	96:4
15	1a Ph	Et	20 PhCH ₂ CH ₂	3ao 95	95:5
16	1a Ph	Et	2p cyclohexyl	3ap 87	93:7
17	1b 4-C ₆ H ₄	Et	2a Ph	3aa 63	94:6
18	1c 4-0 ₂ NC ₆ H ₄	Et	2a Ph	3aa 87	99:1
19	1d 4-FC ₆ H ₄	Me	2h 4-O ₂ NC ₆ H ₄	3dh 82	99:1
20	1d 4-FC ₆ H ₄	Me	2g 4-F ₃ CC ₆ H ₄	3dg 84	99:1
21	1e Me	Et	2a Ph	3aa 21	93:7
22	1e Me	Et	2g 4-F ₃ CC ₆ H ₄	3ag 34	95:5

 $^{\rm a}$ Reaction conditions: 1 (0.55 mmol), 2 (0.5 mmol), Cs2CO3 (1.0 mmol). $^{\rm b}$ Isolated yields.

^c Relative ratio of the crude determined by ¹H NMR spectroscopy.

۸

129





We thought it could be possible to perform deacylation to produce α -fluoro- α . β -unsaturated ketones and amides. It was found that deacylation could be easily achieved by the same method using 2-fluoro-1,3-dione and α -fluoro- β -keto amide compounds (Scheme 3). Thus, reactions of 2-fluoro-1,3-diphenylpropane-1,3-dione (4a) or 2-fluoro-3-oxo-N,3diphenylpropanamide (4b) with benzaldehydes 2 gave olefination products in 61–88% yields with extremely high Z/Eratios.

These olefination reactions can be conducted without using Schlenk technique on a larger scale. The olefination of fluorous benzoylacetate 1a with benzaldehyde (2a) on a two-gram scale occurred in a high yield (81%) similar to that of the reaction conducted on a smaller scale (Scheme 4). The benzoic acid was collected for experimental use. Thus, these reactions should be practical for a number of applications in medicinal chemistry.

According to the reported literature¹¹ and experimental points a possible mechanism for this transformation is proposed in Scheme 5, in which Cs₂CO₃ plays an important role as a promoter of nucleophilic addition. Weak bases could not make nucleophilic addition happen. Strong base make the product decompose into (Z)-2-fluoro-3-phenylacrylic acid (see the Supporting Information). An intramolecular nucleophilic addition of intermediates ii preferentially adopts an antiperiplanar conformation, which is much more thermodynamicly and kineticly stable than its other conformation, and forms a four-membered-ring transition



130

J. Qian et al.



state **iii**. The final elimination of unstable transition state **iii** produces the designed product **3**.

In conclusion, a highly stereoselective olefination reaction of α -fluoro- β -keto esters for the synthesis of α -fluoro- α , β -unsaturated esters has been developed. This method provides a practical, simple, and mild synthetic approach to α -fluoro- α , β -unsaturated esters, which are important units in biologically active molecules. The protocol was also used to prepare α -fluoro- α , β -unsaturated ketones and amides. The high stereoselectivity and excellent yields makes this transformation very efficient and practical. Further studies to extend the synthetic applications for fluorinated compound are ongoing in our group.

Acknowledgment

We thank the Fundamental Research Funds for the Central Universities (30920130111002), National Natural Science Foundation of China (21476116), and Natural Science Foundation of Jiangsu (BK20141394). We also thank the Center for Advanced Materials and Technology for financial support.

Supporting Information

Supporting information for this article is available online at http://dx.doi.org/10.1055/s-0034-1378917.

References and Notes

 (a) Hiyama, T. Organofluorine Compounds: Chemistry Applications; Springer: Berlin, **2000**. (b) Iseki, K. Tetrahedron **1998**, 54, 13887. (c) Kirsch, P. Modern Fluoorganic Chemistry; Wiley-VCH: Weinheim, **2004**. (d) Adejare, A.; Ojima, I.; McCarthy, J. R.; Welch, J. T. J. Med. Chem. **1997**, 40, 2967.

- (2) (a) Van der Veken, P.; Kertèsz, I.; Senten, K.; Haemers, A.; Augustyns, K. Tetrahedron Lett. 2003, 44, 6231. (b) Nakamura, Y.; Okada, M.; Koura, M.; Tojo, M.; Saito, A.; Sato, A.; Taguchi, T. J. Fluorine Chem. 2006, 127, 627. (c) Guan, T.; Yoshida, M.; Ota, D.; Fukuhara, T.; Hara, S. J. Fluorine Chem. 2005, 126, 1185. (d) Pirrung, M. C.: Han, H.: Ludwig, R. T. I. Org, Chem. 1994, 59. 2430. (e) Laue, K. W.; Mück-Lichtenfeld, C.; Haufe, G. Tetrahedron 1999, 55, 10413. (f) Daubresse, N.; Chupeau, Y.; Francesch, C.; Lapierre, C.; Pollet, B.; Rolando, C. Chem. Commun. 1997, 1489. (g) Burkhart, J. P.; Weintraub, P. M.; Gates, C. A.; Resvick, R. J.; Vaz, R. J.; Friedrich, D.; Angelastro, M. R.; Bey, P.; Peet, N. P. Bioorg. Med. Chem. 2002, 10, 929. (h) Kaneko, T.; Clark, R.; Ohi, N.; Ozaki, F.; Kawahara, T.; Kamada, A.; Okano, K.; Yokohama, H.; Muramoto, K.; Arai, T.; Ohkuro, M.; Takenaka, O.; Sonoda, J. WO 9806720, 1998. (i) Jaeger, E. P.; Jurs, P. C.; Stouch, T. R. E. J. Med. Chem. 1993, 28, 275. (j) Honda, H.; Sato, S.; Isomae, K.; Ookawa, J.; Kuwamura, T. DE 3407806, 1984. (k) Hibi, S.; Kikuchi, K.; Yoshimura, H.; Nagai, M.; Tagami, K.; Abe, S.; Hishinuma, I.; Nagakawa, J.; Miyamoto, N. WO 9613478, 1996. (1) Wiedeman, P. E.; Djuric, S. W.; Pilushchev, M.; Sciotti, R. J.; Madar, D. J.; Kopecka, H. US 20020115669, 2002. (m) Sun, J.; Yang, Y.; Huang, Y. CN 103254053, 2013.
- (3) (a) Suzuki, Y.; Sato, M. Tetrahedron Lett. 2004, 45, 1679.
 (b) Lemonnier, G.; Zoute, L.; Dupas, G.; Quirion, J.-C.; Jubault, P. J. Org. Chem. 2009, 74, 4124. (c) Choudary, B. M.; Mahendar, K.; Kantam, M. L.; Kalluri, V. S.; Ranganath Athar, T. J. Adv. Synth. Catal. 2006, 348, 1977. (d) David, E.; Couve-Bonnaire, S.; Jubault, P.; Pannecoucke, X. Tetrahedron 2013, 69, 11039. (e) Lemonnier, G.; Poisson, T.; Couve-Bonnaire, S.; Jubault, P.; Pannecoucke, X. Eur. J. Org. Chem. 2013, 3278. (f) Zoute, L.; Dutheuil, G.; Quirion, J.-C.; Jubault, P.; Pannecoucke, X. Synthesis 2006, 3409.
- (4) (a) Satoh, T.; Itoh, N.; Onda, K.-I.; Kitoh, Y.; Yamakawa, K. Bull. Chem. Soc. Jpn. 1992, 65, 2800. (b) Chevrie, D.; Lequeux, T.; Pommelet, J.-C. Org. Lett. 1999, 1, 1539. (c) Chevrie, D.; Lequeux, T.; Pommelet, J.-C. Tetrahedron 2002, 58, 4759.
- (5) (a) Bergmann, E. D.; Shahak, I.; Appelbaum, J. Isr. J. Chem. 1968,
 6, 73. (b) Grison, C.; Genève, S.; Halbin, E.; Coutrot, P. Tetrahedron 2001, 57, 4903; and references cited therein. (c) Sano, S.; Yokoyama, M.; Shiro, Y.; Nagao, Y. Chem. Pharm. Bull. 2002, 50, 706. (d) Sano, S.; Ando, T.; Yokoyama, M.; Nagao, Y. Chem. Commun. 1997, 559. (e) Sano, S.; Ando, T.; Yokoyama, M.; Nagao, Y. Chem. Nagao, Y. Synlett 1998, 777. (f) Sano, S.; Teranishi, R.; Nagao, Y. Tetrahedron Lett. 2002, 43, 9183. (g) Sano, S.; Saito, K.; Nagao, Y. Tetrahedron Lett. 2003, 44, 3987.
- (6) (a) Welsh, J. T. J. Org. Chem. 1990, 55, 4782. (b) Lin, J.; Welsh, J. T. Tetrahedron Lett. 1998, 39, 9613.
- (7) (a) Zajc, B.; Kake, S. Org. Lett. 2006, 8, 4457. (b) Pfund, E.; Lebargy, C.; Rouden, J.; Lequeux, T. J. Org. Chem. 2007, 72, 7871.
 (c) Alonso, D. A.; Fuensanta, M.; Gómez-Bengoa, E.; Nájera, C. Adv. Synth. Catal. 2008, 350, 1823. (d) Arun, K. G.; Shaibal, B.; Saikat, S.; Soon, B. K.; Barbara, Z. J. Org. Chem. 2009, 74, 3689.
- (8) (a) Burton, D. J.; Greenlimb, P. E. J. Org. Chem. 1975, 40, 2796.
 (b) Veenstra, S. J.; Hauser, K.; Felber, P. Bioorg. Med. Chem. Lett. 1997, 7, 351. (c) Patrick, T. B.; Lanahan, M. V.; Yang, C.; Walker, J. K.; Hutchinson, C. L.; Neal, B. E. J. Org. Chem. 1994, 59, 1210.
 (d) Machleidt, H.; Wessendorf, R. Justus Liebigs Ann. Chem. 1964, 674, 1. (e) Etemad-Moghadam, G.; Seyden-Penne, J. Bull. Soc. Chim. Fr. 1985, 448. (f) van Steenis, J. H.; van der Gen, A. J. Chem. Soc., Perkin Trans. 1 2002, 2117. (g) Pfund, E.; Masson, S.; Vazeux, M.; Lequeux, T. J. Org. Chem. 2004, 69, 4670.
 (h) McCarthy, J. R.; Huber, E. W.; Le, T.; Laskovics, F. M.; Matthews, D. P. Tetrahedron 1996, 52, 45. (i) Kanai, M.; Percy, J.

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.

J. Qian et al.

M. *Tetrahedron Lett.* **2000**, *41*, 2453. (j) Pfund, E.; Lebargy, C.; Rouden, J.; Lequeux, T. *J. Org. Chem.* **2007**, *72*, 7871. (k) Calata, C.; Catel, J. M.; Pfund, E.; Lequeux, T. *Tetrahedron* **2009**, *65*, 3967.

- (9) (a) Ho, T. L. Heterolytic Fragmentation of Organic Molecules; Wiley: New York, 1993. (b) Crabtree, R. H. Nature (London, U.K.) 2000, 408, 415.
- (10) (a) Cai, S. J.; Wang, F.; Xi, C. J. J. Org. Chem. 2012, 77, 2331.
 (b) Yin, L.; Kanai, M.; Shibasaki, M. J. Am. Chem. Soc. 2009, 131, 9610. (c) Trost, B. M.; Xu, J.; Schmidt, T. J. Am. Chem. Soc. 2008, 130, 11852. (d) Chen, Y.; Wang, Y. J.; Sun, Z. M.; Ma, D. W. Org. Lett. 2008, 10, 625.
- (11) (a) Tsuboi, S.; Uno, T.; Takeda, A. Chem. Lett. **1978**, 1325.
 (b) Ueno, Y.; Setoi, H.; Okawara, M. Tetrahedron Lett. **1978**, 39, 3753.
- (12) Nakatsu, S.; Gubaidullin, A. T.; Mamedov, V. A.; Tsuboi, S. *Tetrahedron* **2004**, *60*, 2337.
- (13) Hornung, C. H.; Hallmark, B.; Baumann, M.; Baxendale, I. R.; Ley,
 S. V.; Hester, P.; Clayton, P.; Mackley, M. R. *Ind. Eng. Chem. Res.* **2010**, 49, 4576.

(14) **Typical Experimental Procedure for the Fluoroolefins** The reaction mixture of fluorinated substrates (0.55 mmol), aldehyde (0.5 mmol), Cs_2CO_3 (1 mmol) and MeCN (1.5 mL) was stirred at 40 °C for the indicated time until complete consumption of the starting material, which was monitored by TLC analysis (6–12 h). The solvents were removed by rotary evaporation to provide raw products. The residue was then chromatographed on silica gel (eluent: hexane–EtOAc), affording the desired fluoroolefins.

Ethyl (Z)-2-Fluoro-3-phenylacrylate (3aa)

Colorless oil. ¹H NMR (500 MHz, CDCl₃): δ = 7.64 (d, *J* = 6.8 Hz, 2 H), 7.43–7.35 (m, 3 H), 6.92 (d, *J* = 35.2 Hz, 1 H), 4.35 (q, *J* = 7.1 Hz, 2 H), 1.38 (t, *J* = 7.1 Hz, 3 H). ¹⁹F NMR (470 MHz, CDCl₃): δ = -125.31 (s). ¹³C NMR (126 MHz, CDCl₃): δ = 160.45 (d, *J* = 34.3 Hz), 146.07 (d, *J* = 267.5 Hz), 130.20 (s), 129.30 (d, *J* = 7.2 Hz), 128.68 (s), 127.82 (s), 116.48 (s), 60.89 (s), 13.23 (s). MS (EI): *m/z* = 194.12 [M⁺].

(Z)-2-Fluoro-1-phenyl-3-[2-(trifluoromethyl)phenyl]prop-2-en-1-one (5ar)

Colorless solid. ¹H NMR (500 MHz, CDCl₃): δ = 8.03 (d, *J* = 7.9 Hz, 1 H), 7.90 (d, *J* = 7.6 Hz, 2 H), 7.75 (d, *J* = 7.9 Hz, 1 H), 7.63 (t, *J* = 7.5 Hz, 2 H), 7.56–7.47 (m, 3 H), 7.19 (d, *J* = 33.6 Hz, 1 H). ¹⁹F NMR (470 MHz, CDCl₃): δ = -59.57 (s), -118.61 (s). ¹³C NMR (126 MHz, CDCl₃): δ = 187.69 (d, *J* = 28.3 Hz), 154.69 (d, *J* = 276.2 Hz), 135.67 (s), 133.32 (s), 132.07 (s), 131.51 (d, *J* = 12.1 Hz), 129.47 (d, *J* = 3.6 Hz), 129.32 (s), 129.11 (s), 128.59 (s), 126.22 (q, *J* = 5.5 Hz), 124.96 (s), 122.79 (s), 115.25 (s). MS (EI): *m/z* = 294.15 [M⁺].

(Z)-2-Fluoro-N-phenyl-3-[4-(trifluoromethyl)phenyl]acrylamide (5bg)

Colorless solid. ¹H NMR (500 MHz, DMSO): δ = 10.47 (s, 1 H), 7.90 (d, *J* = 8.2 Hz, 2 H), 7.80 (d, *J* = 8.3 Hz, 2 H), 7.74 (d, *J* = 7.7 Hz, 2 H), 7.35 (t, *J* = 7.9 Hz, 2 H), 7.20–7.07 (m, 2 H). ¹⁹F NMR (470 MHz, DMSO): δ = -61.30 (s), -121.52 (s). ¹³C NMR (126 MHz, DMSO): δ = 157.26 (d, *J* = 29.8 Hz), 50.89 (d, *J* = 281.5 Hz), 137.26 (s), 134.70 (s), 129.89 (d, *J* = 6.0 Hz), 128.52 (d, *J* = 32.1 Hz), 128.18 (s), 125.22 (s), 24.04 (s), 120.38 (s), 111.49 (s). MS (EI): *m/z* = 309.10 [M⁺].