


Pd-Catalyzed Synthesis of *N*-Aryl Indoles




Significance: Reported is a palladium-catalyzed synthesis of *N*-aryl indoles from the reaction of diaryl amines with olefins by a hydroamination–C–H activation route. In recent years, the traditional methods, for example, the venerable century-old Fischer indole synthesis, among other textbook reactions, have been superseded by a barrage of methodologies based on transition-metal-catalyzed heteroannulation processes originating with Larock, with significant contributions by Buchwald, Ackermann, Glorius, Fagnou, among others (see: *Angew. Chem. Int. Ed.* **2013**, 52, 3434; *J. Am. Chem. Soc.* **2012**, 134, 9098; *Angew. Chem. Int. Ed.* **2013**, 52, 12426). The present methodology takes advantage of the Fujiwara–Moritani reaction (hydroamination of olefins) together with a precedented aryl C–H activation to achieve the *N*-aryl indole synthesis. Although a comparison to another ruthenium-based photocatalytic method is given, the significance of the present route in assessment to plethora of recently similar established methods is unfortunately not made and would require reading of about 30 references cited in this report.


Comment: The scope of the reaction is defined mainly with styrenes with a few cases of other aliphatic olefins included. The yields of products are modest to good with noticeably lower yields with sterically demanding styryl systems. *N*-Aryl variation to heterocyclic substituted aryls and pyridyl may be obtained albeit also with lower yields. For unsymmetrical diaryl amines, the regioselectivity of C–H activation–annulations occurs towards the EWG aryl ring. Based on the achieved regioselectivity pattern, a mechanism of initial orthopalladation (for EWG aryls) and initial N-palladation (for EDG aryls) is proposed. The reaction of prototype diphenylamine in the absence of styrene reactant partner in D_2O afforded diphenylamine deuterated in all *ortho*- and *para*-positions (75–80% *d*-incorporation), which suggests reversible electrophilic palladation and raises the question of validity to name this a C–H activation process. However, the absence of *para*-deuterated product in the presence of the styrene co-reactant is good evidence for Pd being trapped by initial N-coordination. Further, putative 2-aminostilbene and indolenine intermediates were shown to be converted into indole products under the given conditions. This is a useful beginning of a mechanistic study whose elaboration is promised.

H. XIANG, C. YANG* (SHANGHAI INSTITUTE OF MATERIA MEDICA, P. R. OF CHINA)
A Facile and General Approach to 3-[(Trifluoromethyl)thio]-4*H*-chromen-4-one
Org. Lett. **2014**, *16*, 5686–5689.

Direct Cyclization–Trifluoromethylthiolation to 4*H*-Chromen-4-ones



$R^1 = 4\text{-H, 4-Me, 4-Et, 4-Pr, 4-}i\text{-Pr, 4-OMe, 4-F, 4-Cl, 4-Br, 4-Ph, 4-(4'\text{-MeOC}_6\text{H}_4), 4\text{-NO}_2, 5\text{-OMe, 5-Et, 5-Br, 5-Cl, 4,5-(OMe)}_2, 4,5\text{-Me}_2, 4\text{-Cl-5-Me, 4,6-Me}_2, 4\text{-Cl-6-Br, 4-(4'\text{-MeO}_2\text{C-C}_6\text{H}_4), 4,5\text{-(CH=CH)}_2$



Significance: Reported is a general synthesis of 3-[(trifluoromethyl)thio]-4*H*-chromen-4-ones **2** from 1-(2'-hydroxyphenyl)-3-dimethylaminoprop-2-enones **1** and the active electrophilic trifluoromethylthio moiety generated from AgSCF_3 and trichloroisocyanuric acid (TCCA). This synthetic approach leads to the direct introduction of the SCF_3 group into chromen-4-ones, which represents a very desirable technique to produce diverse molecules for the medicinal chemist. The screening of reaction conditions showed that the reaction is highly dependent on the solvent; for example, yields in THF were better than in DMF; however, no product was obtained for reactions that were carried out in MeCN, CH_2Cl_2 , DMSO, MTBE or toluene. The reaction also showed dependency on the optimized amount of TCCA (1.5 equiv), and remarkable insensitivity to air and moisture. The synthetic utility of this reaction to synthesize other SCF_3 -containing heterocycles such as a toppyryrone C analogue was also investigated.

Comment: The trifluoromethylthio group is incorporated in many bioactive molecules such as the hypotensive agents losartan and nifedipine (L. M. Yagupolskii et al. *J. Fluorine Chem.* **2001**, *109*, 87). Classic approaches of introducing this group include an indirect, multistep method (A. E. Feiring *J. Org. Chem.* **1979**, *44*, 2907). Herein, an easy and direct synthetic route was used to accomplish the same result. A mechanism was presented without evidence. It includes the possibility of an intramolecular Michael addition–cyclization of **1**, followed by enolate nucleophilic attack on the $^+\text{SCF}_3$ species, then a *N,N*-dimethylamine elimination process to produce **2**.

SYNFACTS Contributors: Victor Snieckus, Mona Gamal-Eldin (Snieckus Innovations)

Synfacts 2015, 11(1), 0014 Published online: 15.12.2014

DOI: 10.1055/s-0034-1379731; Reg-No.: V15714SF

2015 © THIEME STUTTGART • NEW YORK

Rhodium(I)-Catalyzed [4+2]-Cycloaddition Route to 1,2-Dihydropyridines

Category

Synthesis of Heterocycles

Key words


bridged bicycles

C–H activation


dihydropyridines

piperidines

rhodium catalysis



Representative examples:




Significance: Piperidine heterocycles are abundant in numerous pharmaceuticals such as oxycotin, plavix, ritalin, and tofacitinib. The piperidine ring can contain multiple stereocenters, quaternary carbons, or be highly substituted. One method to prepare chiral substituted piperidines is the asymmetric hydrogenation of pyridine or tetrahydropyridine precursors; therefore, a robust, scalable route to a diverse range of substituted pyridines or tetrahydropyridines would be of synthetic value to the pharmaceutical industry. The known Diels–Alder approach to substituted pyridines is compromised by significant electronic and steric constraints (J. A. Varela, C. Saá *Chem. Rev.* **2003**, *103*, 3787; D. L. Boger *Chem. Rev.* **1986**, *86*, 781; A. Saito et al. *Tetrahedron Lett.* **2007**, *48*, 6852).

SYNFACTS Contributors: Victor Snieckus, Brandon P. Schuff (Pfizer)
Synfacts 2015, 11(1), 0015 Published online: 15.12.2014
DOI: 10.1055/s-0034-1379737; **Reg-No.:** V15914SF

Comment: In the presented methodology, a broad scope of 1,2-dihydropyridines was prepared through a rhodium(I)-catalyzed C–H activation–alkenylation–electrocyclization sequence of reactions. Treatment with an electrophile under kinetic or thermodynamic control conditions afforded the corresponding iminiums, which then were treated with nucleophiles to afford the desired tetrahydropyridines with high diastereoselectivity. The 1,2-dihydropyridine intermediates were also shown to react in [3+2] or [4+2] cycloaddition modes to afford bridged bicyclic isoquinuclidines and tropane cores. The catalyst loading was low (<1%), conditions were robust (air-stable), and the tetrahydropyridines were prepared on scale (>100 mmol) in good yield.

A. ALIZADEH,* V. SADEGHI, F. BAYAT, L.-G. ZHU (TARBIAT MODARES UNIVERSITY, TEHRAN, IRAN AND ZHEJIANG UNIVERSITY, HANGZHOU, P. R. OF CHINA)
Highly Efficient Diastereoselective Synthesis of Azabicyclo[2.2.2]octanes
Synlett **2014**, 25, 2609–2612.

One-Pot Diastereoselective Synthesis of Azabicyclo[2.2.2]octanes



Significance: Reported is the base-mediated synthesis of azabicyclo[2.2.2]octanes through the reaction of dibenzalacetone and malononitrile.

The products were unambiguously identified using NMR and IR spectroscopy as well as mass spectrometry and, for one example, by X-ray diffraction analysis. The starting benzalacetones are easily synthesized from benzaldehyde and acetone (C. R. Conard, M. A. Dolliver *Org. Syn.* **1932**, 12, 22). The method was optimized with respect to solvent, temperature and base, and the substrate scope was modestly examined using electron-rich and electron-poor aromatic substitution. The electronics of the ring had surprisingly little effect on the yield of the reaction; however, *ortho*-substituted dibenzalacetones did not provide products, probably due to steric hindrance effects. Also unsuccessful were dibenzalacetones with *para*-hydroxyaryl substituents, possibly due to strong electron-donating effects.


Comment: General interest in the azabicyclo[2.2.2]octane (isoquinuclidine) structure is fueled by its presence in many interesting natural products, and also its wide array of both pharmaceutical and synthetic uses. Interestingly, the cinchona alkaloids contain quinuclidine part structures and have many pharmacological properties, such as muscle relaxant, analgesic, and antimalarial. They are also very useful as organocatalysts (see Book below). Of the several methods for the synthesis of azabicyclo[2.2.2]octanes outlined in the introduction to this new work, many suffer from substrate scope issues or poor *endo/exo* selectivity. The current method should prove useful as it uses easily accessible or readily available starting materials to diastereoselectively provide moderate to good yields of new compounds in an efficient, operationally simple one-pot process.

Book: C. E. Song *An Overview of Cinchona Alkaloids in Chemistry*, In *Cinchona Alkaloids in Synthesis and Catalysis: Ligands, Immobilization and Organocatalysis*, C. E. Song, Ed.; Wiley-VCH: Weinheim, Germany, **2009**, 1–10.

SYNFACTS Contributors: Victor Snieckus, Johnathan Board
Synfacts 2015, 11(1), 0016 Published online: 15.12.2014
DOI: 10.1055/s-0034-1379729; **Reg-No.:** V15514SF

2015 © THIEME STUTTGART • NEW YORK

Aziridine and β -Lactam Synthesis by Amination and Carbonylation Reactions




Significance: Although significant progress has been made in the functionalization of unactivated C–H bonds, the transformation of aliphatic C–H bonds remains a challenging endeavor (H. M. L. Davies et al. *Chem. Soc. Rev.* **2011**, *40*, 1855). Typically, such transition-metal-catalyzed reactions utilize a directing group within the substrate, which serves to position the metal catalyst in the proximity of a particular C–H bond, and proceed through the kinetically favored five-membered cyclometalated intermediate (L. Ackermann *Chem. Rev.* **2011**, *111*, 1315). The current report highlights a different mode of reactivity with the reaction being directed by a simple secondary amine and proceeding through a strained four-membered cyclometalated intermediate. A broad substrate scope is demonstrated for the efficient syntheses of a series of aziridines. In addition, the ring opening of these products provides a range of secondary amines presenting substitution patterns which would be difficult to access by more conventional methods. The methodology was extended to incorporate a palladium-mediated C–H carbonylation to allow access to potentially biologically active β -lactams.


Comment: 3,3,5,5-Tetramethylmorpholin-2-one was initially utilized as the model substrate given that this presents a range of C–H bonds that cannot participate in the classical five-membered cyclopalladation. Treatment with stoichiometric Pd(OAc)₂ led to the isolation of the four-membered ring complex (characterized by X-ray) resulting from the amine-directed cyclopalladation. Oxidation of this intermediate with PhI(OAc)₂ led to the formation of the corresponding aziridine through reductive elimination from a palladium(IV) intermediate. Optimization enabled the reaction to be carried out in a catalytic manner. A broad substrate scope was shown demonstrating excellent selectivity for the formation of the strained heterocycles and the ability to differentiate between methyl groups based on subtle stereoelectronic effects. Replacement of the oxidant PhI(OAc)₂ with CO allowed the development of a carbonylation process proceeding through a palladium(II)–palladium(0) catalytic cycle enabling the synthesis of β -lactams for a range of cyclic and acyclic amines.

A. R. REDDY, C.-Y. ZHOU, Z. GUO, J. WEI, C.-M. CHE* (HKU SHENZHEN INSTITUTE OF RESEARCH AND INNOVATION AND THE UNIVERSITY OF HONG KONG, P. R. OF CHINA) Ruthenium–Porphyrin-Catalyzed Diastereoselective Intramolecular Alkyl Carbene Insertion into C–H Bonds of Alkyl Diazomethanes Generated In Situ from *N*-Tosylhydrazones
Angew. Chem. Int. Ed. **2014**, *53*, 14175–14180.


Tetrahydrofurans and Pyrrolidines from *N*-Tosylhydrazones via C–H Insertion



Representative examples:



Application:




Significance: Reported is a method for synthesizing substituted tetrahydrofurans and pyrrolidines from acyclic *N*-tosylhydrazones via an intramolecular ruthenium–porphyrin catalyzed C–H insertion process. Key to the success of this method was the identification of conditions for minimizing decomposition (elimination) of the alkylcarbene or carbenoid intermediate, which was realized through careful optimization of base, solvent and catalyst. The ruthenium–porphyrin catalyst used in this study is not commercially available, but is readily prepared in a single step. A one-pot method for accessing the tetrahydrofuran or pyrrolidine products from the corresponding acyclic ketone was also reported and proceeded with comparable yields. The power of this methodology is demonstrated through a concise synthesis of the alkaloid (±)-pseudoheliotridane from pyrrolidine.

Comment: Pyrrolidines and tetrahydrofurans are two of the most prevalent heterocycles found in drugs today. The present method demonstrates good scope for accessing these heterocycles, tolerating both alkenes and free alcohols, and provides access to structurally diverse products including spiro, attached ring, and fused ring systems from easily accessible starting materials. *N*-Tosylhydrazones derived from either aldehydes or ketones were viable substrates, allowing for flexible substitution at the 3 position of the heterocycle. In those cases in which two or more stereocenters were introduced, a high level of 2,3-*cis* diastereoselectivity was observed, especially in the synthesis of pyrrolidines. Kinetic isotope effect experiments and the observed stereospecificity of the present reaction provide strong support for a concerted C–H insertion process.

SYNFACTS Contributors: Victor Snieckus, Matt Dowling (Pfizer)
Synfacts 2015, 11(1), 0018 Published online: 15.12.2014
DOI: 10.1055/s-0034-1379701; Reg-No.: V14814SF

2015 © THIEME STUTTGART • NEW YORK

Copper(I)-Catalyzed Synthesis of 3-Amino-2-pyrones and 2,5-Dihydrofurans




Significance: Reported is a three-component reaction of propargyl alcohols, aldehydes, and amines under copper(I) catalysis to form 3-amino-2-pyrones and 2,5-dihydrofurans. It follows a previous report from the Ma laboratories in which the same components afforded allylic amines (*Nat. Commun.* **2014**, *5*, 3884). The difference appears to be that the aldehyde component is ethyl glyoxylate; there is little if any modification of the reaction conditions or the catalyst. Consideration of the readily derived mechanism of the reaction allows formulation of the 2-pyrone **1** as the expected product resulting from mild (silica gel) acidic treatment of isolated intermediate **2**, $R = \text{OEt}$, which in turn results from an allene precursor. A rational consideration that EWG in the aldehyde component would encourage formation of the furan ring was confirmed by using glyoxal and the isolation of dihydrofurans **2**, $R = \text{Ph}$. A reasonable mechanism is given which, although proceeding via isolated **2**, lacks further definition especially in terms of graphical representation of copper involvement. Both processes were tested with at least ten examples each, thus sufficient to justify giving these new processes general status.

SYNFACTS Contributors: Victor Snieckus
Synfacts 2015, 11(1), 0019 Published online: 15.12.2014
DOI: 10.1055/s-0034-1379705; Reg-No.: V15214SF


Comment: The significance of 2-pyrones and dihydrofurans as units possessing biological activity, selective COX-1 inhibition, and as structural units of bioactive natural and unnatural products, respectively, is noted. Fourteen optimization experiments established the best conditions as shown. Scope for $R^1 =$ alkyl, benzyl, aryl, and $R^2, R^3 =$ substituted pyrrolidinyl, piperidinyl, and one case of dibutyl was established for the 2-pyrone synthesis, while mainly $R^2, R^3 =$ piperidinyl derivatives for the dihydrofurans were obtained. In the glyoxal study, as expected, EWG = NO_2 led to poor yields of product. Experiments were carried out in a flame-dried Schlenk tube under argon atmosphere. Significance of this work would have been increased if further transformation of products **1** and **2** would have been carried out or noted (for example, for **1**, use for Diels–Alder chemistry).

C. HUO,* Y. YUAN, M. WU, X. JIA, X. WANG, F. CHEN, J. TANG (NORTHWEST NORMAL UNIVERSITY, LANZHOU, P. R. OF CHINA)
Auto-Oxidative Coupling of Glycine Derivatives
Angew. Chem. Int. Ed. **2014**, *53*, 13544–13547.


Auto-Oxidation–Povarov Annulation Approach to Quinolines



Selected examples:



Other reactions also reported:



Significance: The reported auto-oxidation of *N*-aryl glycines **1** afforded a versatile imine intermediate, which underwent the Povarov annulation affording **3**; further oxidation led to **4**, while nucleophilic addition afforded **5**. When 2,3-dihydrofuran was employed in the Povarov reaction, a subsequent ring opening–ring closing sequence was observed, resulting in the isolation of compound **3c**. The reported Povarov reaction, in particular, is complementary to existing reactions of anilines with glyoxalates and terminal alkynes under acidic catalysis (for example, see: J. B. Bharate et al. *Org. Biomol. Chem.* **2014**, *12*, 6267) and avoids the selectivity issues associated with the functionalization of bishalogenated quinolines.

SYNFACTS Contributors: Victor Snieckus, Benjamin N. Roche (Pfizer)
Synfacts 2015, 11(1), 0020 Published online: 15.12.2014
DOI: 10.1055/s-0034-1379702; Reg-No.: V14914SF

Comment: Although a more extensive demonstration of substrate scope or functional group tolerance would have been appreciated in the presented synthesis, highlights include the tolerance of *N*-acyl glycines, allyl esters, and halides. Oxidation side products **4** were always observed in the Povarov annulations, but were minimized when R² was an electron-rich substituent. The reaction rates were found to be highly dependent on solvent choice, with a 5:1 mixture of MeCN and DCE required for optimal reactivity. The reaction efficiency was also noted to depend on the use of 3–5 week old DCE or on the addition of a trace quantity of HCl, an observation that will impact the reproducibility of these results without due attention.

Synthesis of Indoles, Pyrazoles, and Pyridazinones

Category

Synthesis of Heterocycles

Key words


indoles

pyridazinones

pyrazoles

diazonium salts

alkenes



Significance: Reported is a one-pot synthesis of indoles, pyrazoles, and pyridazinones by a variation of the Japp–Klingemann Fischer indole synthesis, involving a trifluoromethylation. The reaction was found to well-tolerate a variety of functionalized arenediazonium salts and aryl allyl ketones. *meta*-Substituted arenediazonium salts provided mixtures of regioisomeric indoles (**A** and **B**). *para*-Substituted arenediazonium salts were also used with methyl pent-4-enoate to provide dihydropyridazinones in good yields.

Comment: The indole and pyrazole heterocyclic core is found in a number of top-selling drugs, such as sumatriptan, zolmitriptan, rizatriptan, tadalafil, and celecoxib (M. Baumann et al. *Beilstein J. Org. Chem.* **2011**, *7*, 442). Therefore, a simple and efficient synthesis of these heterocyclic cores is a worthwhile quest. The developed method gives access to various trifluoromethylated heterocycles. Previously, a similar methodology has been used to synthesize pyrazoles (A. Citterio et al. *J. Heterocycl. Chem.* **1981**, *18*, 763). Unexplained is the fact that all examples of dihydropyridazinone synthesis use *para*-substituted arenediazonium salt precursors.

SYNFACTS Contributors: Victor Snieckus, M. Selim Hossain (Snieckus Innovations)

Synfacts 2015, 11(1), 0021 Published online: 15.12.2014


DOI: 10.1055/s-0034-1379730; Reg-No.: V15614SF

D. GHORAI, J. CHOUDHURY* (INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH BHOPAL, INDIA)

Exploring a Unique Reactivity of N-Heterocyclic Carbenes (NHC) in Rhodium(III)-Catalyzed Intermolecular C–H Activation/Annulation

Chem. Commun. **2014**, *50*, 15159–15162.

NHCs in Rhodium-Catalyzed C–H Activation–Annulation to Fused Imidazolium Salts



Significance: The increasing interest in N-heterocyclic carbenes (NHCs) arises arguably from their unprecedented stereoelectronic properties, strong metal–NHC bonding, and great stability of their metal complexes. NHCs act as both ligands and directing groups. These properties make NHCs useful in C–H functionalization as well as C–C and C–heteroatom bond-forming catalysis (see Review below). Reported here is the first directed intramolecular C–H functionalization–annulation reaction using a NHC–rhodium(III) complex as catalytic system. Thus, reaction of imidazolium salts **1** with internal alkynes **2**, bearing different types of substituent groups, furnishes a variety of imidazo[1,2-*a*]quinolinium derivatives **3** in a one-pot process in 35–93% yield.

Review: M. N. Hopkinson, C. Richter, M. Schedler, F. Glorius *Nature* **2014**, *510*, 485–496.

Comment: This is the first report of this kind of reaction, which is formally a [4+2]-cycloaddition process. A range of substrates was evaluated and a relationship between the electronic demand of the rings and the alkyne was established. Another pleasant surprise is the mild room-temperature reaction conditions. The NHC–rhodium(III) complex was isolated and characterized, thus supporting strongly the proposed mechanism. The main disadvantage of this methodology is the use of a large amount of AgOTf, which can make its scalability expensive. Further applications are anticipated.

SYNFACTS Contributors: Victor Snieckus, Keller G. Guimarães
Synfacts 2015, 11(1), 0022 Published online: 15.12.2014
DOI: 10.1055/s-0034-1379740; **Reg-No.:** V16214SF


2015 © THIEME STUTTGART • NEW YORK

Z. ZHANG,* W. ZHANG, J. LI, Q. LIU, T. LIU, G. ZHANG* (HENAN NORMAL UNIVERSITY, XINXIANG, P. R. OF CHINA)

Synthesis of Multisubstituted Pyrroles from Doubly Activated Cyclopropanes Using an Iron-Mediated Oxidation Domino Reaction

J. Org. Chem. **2014**, *79*, 11226–11233.

Iron-Mediated Synthesis of Pyrroles from Cyclopropanes



Significance: The synthesis of highly substituted pyrroles **3** from cyclopropanes **1** and amines **2** via an iron-mediated sequential ring opening–cyclization–dehydrogenation reaction is reported. The conditions were optimized using different solvents, reaction times, and iron catalysts. The scope was studied and cyclopropanes **1** bearing methyl and aryl substituents at R^1 were tested with the latter giving better yields. EWG- and EDG-substituted aryl amines as well as $\text{R}^2 = \text{OEt}$ were well tolerated. Also, a methyl group in the cyclopropane ($\text{R}^4 = \text{Me}$) was suitable, furnishing 1,2,3,5-substituted pyrrole **3** in good yield. A series of aromatic and aliphatic amines were screened as well: aliphatic amines furnished pyrroles **3** in low yields, while aromatic amines gave better results, although the yields decrease according to the position of the phenyl substituent. A reaction mechanism involving a radical process was suggested based on radical trapping experiments.

Comment: Cyclopropane derivatives can be used as precursors to synthesize a variety of useful heterocyclic motifs (C. A. Carson, M. A. Kerr *Chem. Soc. Rev.* **2009**, *38*, 3051; J. R. Green, V. Snieckus *Synlett* **2014**, *25*, 2258). The present work reports an efficient synthesis of tri- and tetra-substituted pyrroles in moderate to good yields from readily available cyclopropanes and amines. The reaction shows a broad substrate scope; both EWG and EDG in **1** and **2** affording pyrroles **3** in comparable yields. While anilines with different patterns of substitution furnished **3** in good yields, a large excess of amine was required when benzylamines were used. The readily available cyclopropanes can be easily prepared in high yields by the reaction of a dicarbonyl compound with 1,2-dibromoethane (Z. Zhang et al. *Angew. Chem. Int. Ed.* **2007**, *46*, 1726).

SYNFACTS Contributors: Victor Snieckus, Livia Cristina Frota
Synfacts 2015, 11(1), 0023 Published online: 15.12.2014
DOI: 10.1055/s-0034-1379704; **Reg-No.:** V15114SF

2015 © THIEME STUTTGART • NEW YORK

Category

Synthesis of Heterocycles

Key words

pyrroles

cyclopropanes


iron catalysis

L.-S. GE, Z.-L. WANG, X.-L. AN, X. LUO,* W.-P. DENG* (EAST CHINA UNIVERSITY OF SCIENCE AND TECHNOLOGY, SHANGHAI, P. R. OF CHINA)

Direct Synthesis of Polysubstituted 2-Aminothiophenes by Cu(II)-Catalyzed Addition/Oxidative Cyclization of Alkynoates with Thioamides

Org. Biomol. Chem. **2014**, *12*, 8473–8479.

Synthesis of 2-Aminothiophenes from Alkynoates and Thioamides



Significance: Reported is the synthesis of 2-aminothiophenes **3** via a copper-catalyzed addition–oxidative cyclization of alkynoates **2** with thioamides **1**. Among a series of tested copper salts, $\text{Cu}(\text{OAc})_2$ proved to be the most efficient under aerobic conditions. There was no improvement in yields when the reaction was carried out under an oxygen atmosphere. When the reaction was done under a nitrogen atmosphere, only a trace amount of the product was obtained. Instead, compound **4** was isolated in 46% yield, suggesting that the reaction proceeds through a Michael addition–oxidative cyclization pathway. The reaction also proceeded with catalytic FeCl_3 , albeit in comparatively lower yield. The reaction failed at 50 °C and a lower yield was observed at 100 °C. A wide variety of EWGs in **1** were tested to synthesize **3** in moderate to good yields. The reaction of N-mono-substituted β -ketothioamide with dimethyl acetylenedicarboxylate gave a mixture of compounds, which were not characterized.

Comment: 2-Aminothiophene is a useful building block for synthetic chemists as well as a key structural unit in biologically active molecules, such as olanzapine which is used for the treatment of schizophrenia and bipolar disorder. Substituted 2-aminothiophenes are generally synthesized by the direct functionalization of thiophenes or by the cyclization of suitable precursors. A number of methods using the latter strategy are known (see Review below). The current method utilizes readily available starting materials and is easy to set up due to its tolerance to aerobic conditions. However, the reaction scope is limited. Also, it would have been interesting to see the reaction carried out with unsymmetrical alkynoates.

Review: A. El-Mekabaty *Synth. Commun.* **2014**, *44*, 1–31.

SYNFACTS Contributors: Victor Snieckus, Suneel P. Singh (Snieckus Innovations)

Synfacts 2015, 11(1), 0024 Published online: 15.12.2014

DOI: 10.1055/s-0034-1379728; Reg-No.: V15414SF

2015 © THIEME STUTTGART • NEW YORK

A. J. ROSENBERG, I. AHMED, R. J. WILSON, T. M. WILLIAMS, L. KAMINSKY, D. A. CLARK*
(SYRACUSE UNIVERSITY, USA)

An Improved Synthesis of Imidazo[4,5-*b*]pyridines and Imidazo[4,5-*b*]pyrazines by Palladium Catalyzed Amidation Using Xantphos in a 1,4-Dioxane:*tert*-Amyl Alcohol Solvent System

Adv. Synth. Catal. **2014**, *356*, 3465–3470.

An Improved Procedure for a Pd-Catalyzed Amidation Approach to Imidazopyridines

Category

Synthesis of Heterocycles


Key words

imidazo[4,5-*b*]pyridines

imidazo[4,5-*b*]pyrazines

palladium catalysis

amidation



Z = CH, N, CHMe (1 example)

R¹ = *i*-Pr, *c*-Pent, Cy, Ph, Bn, CH₂(4-MeOC₆H₄), CH₂(2,4-(MeO)₂C₆H₃), CH₂(2,5-(MeO)₂C₆H₃), CH₂(4-Me₂NC₆H₄), CH₂(4-FC₆H₄), CH₂(3-FC₆H₄), CH₂(4-ClC₆H₄), CH₂(2-ClC₆H₄), CH₂(4-PhC₆H₄), CH₂(3,5-(MeO)₂C₆H₄), CH₂(4-F₃CC₆H₄), CH₂(4-NCC₆H₄), CH₂(3-O₂NC₆H₄), CH₂Cy, (*R*)-CHMePh

R² = H, Me, Ph, Cy, 2-styryl, furan-2-yl

Significance: The present report is a full paper on the previous communication concerned with the synthesis of imidazopyridines using a palladium-catalyzed coupling reaction of amides and 3-amino-2-chloropyridines (A. J. Rosenberg, J. Zhao, D. A. Clark *Org. Lett.* **2012**, *14*, 1764). The current study aimed to expand the substrate scope, and to find more economical phosphine-based ligands instead of the previously utilized biaryl ligand, Me₄-*t*-Bu-XPhos. The reaction was tested with the Xantphos ligand, despite the fact that the reaction had previously failed when run in *t*-BuOH. Re-examination of the reaction with Xantphos in aprotic solvents led to the desired imidazopyridine together with an unwanted protodehalogenated pyridine byproduct. In order to minimize the formation of this byproduct, a mixed protic–aprotic solvent system using dioxane–*t*-AmOH was devised giving rise to a 93% yield of the desired imidazopyridine. The new reaction conditions improved the yields in some cases, and in others, allowed for product formation, where previously only decomposition had been observed [Z = CH, R¹ = CH₂(4-ClC₆H₄), R² = H; and any substituent containing a nitro group].



Comment: The presented procedure is an improvement over the one reported previously. With the current procedure, chlorinated substrates survive the reaction better, and electron-poor substrates afford higher yields. Further, the authors rightly claim economic benefits and a quick comparison of the prices (Sigma-Aldrich website, 10/11/2014) indicates the following prices: Xantphos: 61 CAD\$/mmol, and Me₄-*t*-Bu-XPhos: 135 CAD\$/mmol (1 gram quantity). No drawbacks of the procedure are mentioned, but brominated and iodinated substrates are evident in their absence. Furthermore, the scope of pyridine starting materials was not well studied; this is most likely due to poor availability of substituted 3-amino-2-chloropyridines.

SYNFACTS Contributors: Victor Snieckus, Toni Rantanen
Synfacts 2015, 11(1), 0025 Published online: 15.12.2014
DOI: 10.1055/s-0034-1379736; **Reg-No.:** V15814SF

2015 © THIEME STUTTGART • NEW YORK

S. CHEN, J. PAN, Y. WANG, Z. ZHOU* (NANKAI UNIVERSITY, TIANJIN, P. R. OF CHINA)
Stereocontrolled Construction of the 3,4-Dihydrothiacarbazol-2(9*H*)-one Skeleton by Using Bifunctional Squaramide-Catalyzed Cascade Reactions
Eur. J. Org. Chem. **2014**, 7940–7947.

Enantioselective Synthesis of 3,4-Dihydrothiacarbazol-2(9*H*)-ones



Significance: Reported is the enantioselective synthesis of 3,4-dihydrothiacarbazol-2(9*H*)-ones **3** by reaction of indoline-2-thiones **1** with *N*-alkenylphthalimides **2** catalyzed by the chiral squaramide **4**. Screening of organocatalysts with double hydrogen-bond donor ability led to squaramide **4** derived from *L*-*tert*-leucine as the best catalyst for this transformation affording high enantioselectivity. The reaction conditions were optimized in terms of solvent, temperature, and catalyst loadings. Lower temperatures (0 °C) culminated in lengthy reaction time and lower yield but equivalent ee, while higher temperatures (40 °C) provided equivalent reaction yields but loss of stereocontrol. The study of the reaction scope showed that the presence of different substituents on both **1** and **2** were tolerated, but in some cases loss of stereocontrol without following a pattern was observed.


Comment: The indole skeleton is an important class of heterocycles present in many natural products with broad biological activities, and can be synthesized by many well-described methodologies (see Review below). The thiopyran indole **3** was obtained by an activation process promoted by two hydrogen-bonding interactions of **2** with the squaramide organocatalyst, followed by a Michael addition step and a thiolysis reaction. The starting materials **1** and **2** are readily available. Although a mild process, the reported approach has long reaction times and the study of the reaction scope is narrow.

Review: G. R. Humphrey, J. T. Kuethe *Chem. Rev.* **2006**, *106*, 2875–2911.













SYNFACTS Contributors: Victor Snieckus, Sara Gomes
Synfacts 2015, 11(1), 0026 Published online: 15.12.2014
DOI: 10.1055/s-0034-1379703; **Reg-No.:** V15014SF

2015 © THIEME STUTTGART • NEW YORK

Copper-Catalyzed Amidation of π -Excessive Heterocycles



Representative examples:



Significance: Amino-substituted heterocycles are prevalent as subunits in a large number of pharmaceutical drug targets. As a result, transition-metal-catalyzed aryl C–N bond formation has been studied extensively. Traditionally, these compounds are synthesized by coupling heteroaryl halides with amine surrogates. Pan and co-workers report an efficient method for direct amidation of π -electron-rich heterocycles with *N*-fluorobenzenesulfonimide (NFSI). Interestingly, even though NFSI is typically used as a fluorinating reagent, no fluorinated byproducts are observed in this reaction. The procedure uses mild conditions to provide the desired functionalized products in high yields.

Comment: A variety of electron-deficient and electron-rich thiophenes, furans, and pyrroles undergo regioselective amidation under these reaction conditions. Although the reaction does proceed at ambient temperature, higher yields are obtained at slightly elevated temperature (60 °C). A mechanistic pathway involving copper(I), copper(II), and copper(III) species is proposed, and further studies to investigate the mechanism of this transformation are underway. For a recent method of 2-aminothiophene and other π -excessive heterocycle synthesis, see the copper(II)-catalyzed synthesis of 2-aminothiophenes from alkynoates and thioamides (*Org. Biomol. Chem.* **2014**, *12*, 8473; *Synfacts* **2015**, *11*, 24).