



M. KAUSHIK, K. BASU, C. BENOIT, C. M. CIRTU, H. VALI, A. MOORES* (MCGILL UNIVERSITY, MONTRÉAL AND INSTITUT NATIONAL DE SANTÉ PUBLIQUE DU QUÉBEC, QUEBEC CITY, CANADA)

Cellulose Nanocrystals as Chiral Inducers: Enantioselective Catalysis and Transmission Electron Microscopy 3D Characterization

J. Am. Chem. Soc. **2015**, *137*, 6124–6127.

Enantioselective Hydrogenation of Ketones on Palladium/Cellulose Nanocrystals



Significance: Palladium(0) deposited on cellulose nanocrystals (Pd@CNCs) was prepared by treatment of CNCs with palladium(II) chloride in aqueous hydrochloric acid, followed by the reduction with hydrogen gas. The Pd@CNCs was used in an enantioselective hydrogenation of prochiral ketones under hydrogen pressure in water to give the corresponding alcohols in 67–91% yield and 5–65% ee.

Comment: The aqueous suspension of Pd@CNCs was reused twice without significant loss of the catalytic activity or enantioselectivity. CSCs did not catalyze the carbonyl hydrogenation in the absence of palladium. Hydrogenation of 2-methoxyacetophenone on PdNPs@C under similar conditions gave the corresponding racemic alcohol in 38% yield.

SYNFACTS Contributors: Yasuhiro Uozumi, Yoichi M. A. Yamada, Aya Ohno

Synfacts 2015, 11(8), 0879 Published online: 20.07.2015

DOI: 10.1055/s-0034-1381113; Reg-No.: Y08515SF

2015 © THIEME STUTTGART • NEW YORK

Category

Polymer-Supported Synthesis

Key words

palladium

cellulose

nanocrystals

asymmetric catalysis

hydrogenation

ketones

Synfact
of the month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.