

Specialized clinical cytology may improve the results of EUS (endoscopic ultrasound)-guided fine-needle aspiration (FNA) from pancreatic tumors

Authors

Arne R. Schneider¹, Andreas Nerlich², Theodoros Topalidis³, Wolfgang Schepp¹

Institutions

Institutions are listed at the end of article.

submitted 31. August 2014
accepted after revision
22. September 2014

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1390886>
Published online: 5.12.2014
Endoscopy International Open 2015; 03: E134–E137
© Georg Thieme Verlag KG
Stuttgart · New York
E-ISSN 2196-9736

Corresponding author

Arne Schneider, MD
Department of
Gastroenterology, Hepatology,
and Gastroenterological
Oncology
Bogenhausen Academic
Teaching Hospital
Englschalkinger Str. 77
D-81925 Munich
Germany
Fax: +4989/9270-2486
arne.schneider@lrz.tum.de

Background and study aims: A variety of factors (needle type, needle passes, tumor location, cytological assessment, etc.) may influence the diagnostic accuracy of endoscopic ultrasound-guided fine-needle aspiration cytology (EUS-FNAC) from pancreatic tumors. Whereas most published studies report a diagnostic accuracy of >80% for EUS-FNAC, the results in routine settings are often considerably lower. This retrospective study aimed to define the effect of switching microscopic assessment from a standard pathology department to a highly specialized institute of cytology.

Patients and methods: A total of 63 patients underwent EUS-FNAC of solid or semisolid pancreatic masses. Specimens of the first consecutive 20 cases (Phase 1) were assessed by the local department of pathology. Then in Phase 2, involving another 43 subsequent cases, a specialized cytology laboratory examined all aspirates.

All EUS-FNACs were performed in the same manner, using a 22-gauge needle. After cytological evaluation, all patients either underwent surgery or were followed up for at least 6 months.

Results: Of the tumors, 56 were solid and 7 semisolid; the mean size was 30 mm. Sensitivity (sens.), specificity (spec.), positive predictive value (PPV), and negative predictive value (NPV) of EUS-FNAC were 38.5% (95%CI [confidence interval] 13.9–68.4%), 100% (59.0–100%), 100% (47.8–100%), and 46.7% (21.3–73.4%) during Phase 1 versus 91.4% (95%CI 76.9–98.2%), 100% (63.1–100%), 100% (89.1–100%), and 72.7% (39.0–94.0%) during Phase 2.

Conclusion: These results emphasize the considerable impact of a dedicated cytological evaluation on the results of EUS-FNAC.

Introduction

Endoscopic ultrasound-guided fine-needle aspiration (EUS-FNA) is an established means for the diagnostic workup of pancreatic masses [1, 2]. Since its introduction into clinical medicine, the technique has been refined and different needle types have been compared with regard to their diagnostic gain. Numerous studies report accuracy and sensitivity of EUS-FNA cytology to reach up to 95%. However, there is a large variation in these statistical results, ranging from 68.1 to 95% [3–10]. The reasons for this discrepancy remain ambiguous, although the reality in many endoscopy departments is that many physicians are dissatisfied with the accuracy of EUS-FNAC in their clinical routines. The technique of tissue sampling via EUS-guided FNA is well defined and largely standardized. The question of preference for small (25-gauge), intermediate (22-gauge), or large-size (19-gauge) needles – or whether to use core biopsy needles, with a chance of obtaining a

histological diagnosis – seems to remain a matter of ongoing debate [3, 11–15].

Once successful aspiration has been achieved, further processing of the specimen by endoscopists and assessment by cytopathologists seem highly variable: Some physicians prefer formalin fixation to enable histopathological embedding, whereas the majority prepare a varying number of slides that undergo either air drying or chemical fixation. Additionally, cytopathologists employ a wide range of staining methods, depending on individual preferences.

During past years, we have modified several of the abovementioned aspects to improve the sensitivity of EUS-FNAC in our department. However, none of these merely technical modifications yielded a detectable increase in sensitivity.

We therefore agreed on switching the microscopic assessment of the slides from our in-house department of pathology to a dedicated laboratory of clinical cytology. The present study compares the results of EUS-FNAC of solid or semiso-

License terms



lid pancreatic masses performed at our department before and after this modification.

Patients and methods

The study population comprised all patients undergoing EUS-FNA at our department for diagnostic workup of solid or semisolid (cystic with solid fractions) pancreatic masses between January 2009 and September 2012. Patients were identified retrospectively by using the electronic medical information database of the hospital. Most patients underwent EUS-FNA for cytological confirmation of suspected unresectable adenocarcinoma of the pancreas. In contrast, patients with highly suspicious but potentially resectable tumors were sent directly for surgery without pursuing a preoperative microscopic diagnosis. Individual courses of disease were followed for a minimum of 6 months, particularly to rule out false-negative results of cytology where pancreatic cancer might evolve despite nonmalignant cytology. In the latter case, lesions were considered malignant either if a carcinoma was histologically confirmed after resection, if the size of the lesion increased during follow-up, or if its size decreased during chemotherapy.

EUS-FNA was performed using a curvilinear array echoendoscope (Pentax EG-3870UTK, Pentax Medical, Hamburg, Germany) attached to a Hitachi EUB 8500 ultrasound processor (Hitachi Medical Systems, Wiesbaden, Germany). The procedure was performed by experienced endoscopists (>30 previous procedures) with patients in the left lateral position under sedation with intravenous propofol. For tumors of the pancreatic head, puncture was done from the duodenal position in the majority of patients. The remaining patients were visualized and biopsied using a transgastric approach. All EUS-FNA were performed with a 22-gauge needle (Sonotip II, Medi-Globe, Aachenmühle, Germany). After visualization of the tumor in a stable position of the endoscope, the needle was advanced into the tumor under continuous EUS guidance, then the stylet was entirely removed and suction applied. A minimum of six needle passes to and fro were performed inside of the lesion during two to four separate punctures, depending on the macroscopic appearance of the specimen with respect to cellularity and bloody additives. Further punctures were abandoned when gross inspection augured tissue without relevant bloody admixture. The aspirate was then recovered by reintroduction of the stylet and subsequent sparging of the needle with air from a 5-mL luer lock syringe. Then smears on glass slides were readily prepared from aspirated fluid with corpuscular fractions. Additionally, cellular clots or spools were gently rolled over glass slides to allow for a detachment of cells on the slide surface. The slides were finally air dried and sent for further cytopathological evaluation.

During study Phase 1 from January 2009 through October 2010, dried aspirates were sent for microscopic diagnosis to the hospital's institute of pathology. There, the slides were stained with Giemsa and analyzed for adequacy of the specimen, cellularity, blood cell additives, and final diagnosis. After this period (Phase 2, November 2010–September 2012), the slides were sent to a specialized institute of clinical cytology. In contrast, May-Grünwald-Giemsa was the standard staining method of the cytologist, whereas further microscopic assessment of cytological aspirates corresponded to the workflow outlined above. Sensitivity (sens.), specificity (spec.), positive predictive value (PPV), and negative predictive value (NPV) as well as corresponding 95%

confidence interval (CI) were calculated with GraphPad Prism 6 (GraphPad Software Inc., La Jolla, CA, USA). The study was approved by the local ethics committee (Ethics Committee of the Faculty of Medicine, Technical University Munich; approval no. 588/13).

Results

Of 64 consecutive patients who had undergone EUS-FNAC, aspirates of 63 patients (24 male [38%], 39 female [62%]; mean age 70, range 28–85) showed cytological features of pancreatic parenchyma and were included in the study. During Phase 1, 20 patients underwent EUS-FNAC; in Phase 2, patients undergoing the procedure numbered 43. The tumor sites of the entire group were: 35 tumors (56%) located in the head; 21 tumors (33%) in the body; and 7 tumors (11%) in the tail of the pancreas. When the study population was differentiated according to affiliation (Phase 1/Phase 2 of the study period), the distribution of the sites broke down to: 5/16 (25/37%) in the head; 11/22 (55/51%) in the body; and 4/5 (20/12%) in the tail. Lesions were classified hypoechoic in 56 cases (89%) and hypoechoic with cystic areas in 7 cases (11%). The mean size of the tumors in the entire population was 30 mm (range 12–80 mm), with 39 mm (range 12–80 mm) in the Phase 1 group, and 30 mm (range 13–70 mm) in the Phase 2 group ($P=0.12$, Mann-Whitney test). During EUS-FNAC, 2 to 5 separate needle passes provided 4 to 27 slides with smears. Mean follow-up of all patients was 17.8 months (range 6.2–34.7 mo).

Surgical resection was done on 9 patients and 54 patients were solely followed up. On final diagnosis, 48 lesions were malignant (44 adenocarcinoma, 3 neuroendocrine tumors [NET], and 1 metastasis of breast carcinoma); and 15 lesions were of benign origin (2 focal post-inflammatory residues after acute pancreatitis, 2 focal inflammatory changes within chronic pancreatitis, 1 autoimmune pancreatitis, 2 intraductal papillary mucinous neoplasms [IPMN], 3 hemorrhagic pancreatic cysts, and 5 undefined).

Accordingly, sens., spec., PPV, and NPV of EUS-FNAC were 38.5% (95%CI 13.9–68.4%), 100% (59.0–100%), 100% (47.8–100%), and 46.7% (21.3–73.4%) during Phase 1 versus 91.4% (95%CI 76.9–98.2%), 100% (63.1–100%), 100% (89.1–100%), and 72.7% (39.0–94.0%) during Phase 2 (Table 1). Accuracy was 60 and 93%, respectively.

Discussion

Published data report on a high accuracy of EUS-FNAC for the diagnosis of pancreatic tumors. According to a variety of studies, mean sensitivity, specificity, and accuracy can be expected to be around 83, 100, and 88% – although these statistical parameters may vary from 54 to 95%, 71 to 100%, and 65 to 96%, respectively [16–18].

However, clinical experience among physicians applying the technique in routine diagnostics is often considerably lower than these study results. Some endoscopists may even have abandoned EUS-guided fine-needle aspiration because, due to ambiguous cytology, they lack faith in the results. In this regard, several factors may have an effect on the diagnostic yield of EUS-FNAC.

Table 1 Summary of statistical results.

Phase 1 (n=20)				Phase 2 (n=43)			
	Real result		Σ		Real result		Σ
	Benign	Malignant			Benign	Malignant	
FNAC +	0	5	5	FNAC +	0	32	32
FNAC-	7	8	15	FNAC-	8	3	11
Σ	7	13	20	Σ	8	35	43
Sens.	38.5 % (13.9 – 68.4 %)			Sens.	91.4 % (76.9 – 98.2 %)		
Spec.	100 % (59.0 – 100 %)			Spec.	100 % (63.1 – 100 %)		
PPV	100 % (47.8 – 100 %)			PPV	100 % (89.1 – 100 %)		
NPV	46.7 % (21.3 – 73.4 %)			NPV	72.7 % (39.0 – 94.0 %)		

Abbreviations: FNAC, fine-needle aspiration cytology benign; FNAC +, fine-needle aspiration cytology malignant; NPV, negative predictive value; PPV, positive predictive value; Sens., sensitivity; Spec., specificity.
95 % confidence interval.

The endoscopic technique of EUS-FNA has been well defined and even outlined in clinical guidelines. A minimum of two to five needle passes per lesion, depending on a gross macroscopic assessment of the aspirate by the endoscopist, are considered mandatory for adequate microscopic diagnosis [18]. The question of whether to use smaller (25-gauge) or larger size (19-, 22-gauge) or even core biopsy needles is still a matter of debate [3, 11, 12]. However, conflicting results of these studies suggest a minor effect of this issue. Immediate microscopic assessment of aspirates by an on-site cytopathologist may further improve EUS-FNAC and reduce the need for multiple needle passes; however, this is rarely available in the routine setting [19–21]. If the initial EUS-FNAC is not diagnostic, repeating the procedure may improve results [19, 20]. Alternatively, percutaneous computed tomography (CT) or ultrasound-guided puncture can be performed, but implies a higher risk for peritoneal tumor seeding. There is no evidence that larger tumors facilitate FNAC assessment [22]. This may derive from the fact that the larger the lesion, the more often nondiagnostic aspirates from cystic or necrotic areas are encountered.

Regarding microscopic diagnosis, cytopathologists generally consider pancreatic adenocarcinoma a challenging task because scarce cellularity, inflammation, and desmoplastic changes may mimic both well-differentiated carcinoma in chronic pancreatitis and fibrosing pancreatitis in cancers [17, 23, 24]. Since standardized guidelines defining the cytological workup are lacking, preparation and further processing of slides remain highly individual. Whereas some cytopathologists prefer air-dried samples, others recommend different methods of fixation. Moreover, staining methods largely differ and imply Papanicolaou, hematoxylin-eosin, or May-Grünwald-Giemsa [3, 10, 22–27].

During a continuous effort to improve the results of EUS-FNAC in our own clinic, we changed several parameters (e.g., needles of different sizes and shapes, number of needle passes and smears, puncture with and without vacuum, etc.) regarding the technical process of specimen acquisition. None of these measures, however, yielded substantial improvement. As a consequence, we discussed the issue with the local pathologist and agreed upon sending FNAC smears to an external specialized cytologist. Although we continue to obtain excellent histopathological results of core biopsies from the local pathologist, the evaluation of FNAC smears by a specialized cytologist had a striking effect on the diagnostic performance. In particular, sensitivity exceeded 90 % and therefore equaled published data.

In most cases, we perform EUS-FNAC of solid pancreatic lesions to prove malignancy prior to initiating palliative chemotherapy or chemoradiotherapy. For this reason, the pretest probability of adenocarcinoma was quite high in the relatively large tumors (mean size 30 mm) in this group of patients. Nevertheless, previous publications that addressed the statistical issues of EUS-FNAC investigated tumors of similar sizes and are therefore comparable with our study [3–10].

Due to the fact that the technical aspects of EUS-FNAC are largely standardized (or have only a minor impact on sensitivity and specificity), preparation of smears and microscopic assessment remain the most variable factors for EUS-FNAC results. A retrospective study by Alsibai et al. has shown that an expert cytopathologist may considerably improve the sensitivity of EUS-FNAC as compared with local cytopathologists [27]. A similar effect has been demonstrated for the training of on-site cytotechnicians present during EUS: Diagnostic accuracy of smear evaluation increased from 74.8 to 90.5 % after 1 year of training, involving more than 100 patients [25]. Recently, a German survey among more than 100 EUS centers demonstrated sobering results regarding the outcome of EUS-FNAC – more than half of the institutions reported a success rate of 75 % and lower [28].

Conclusions

Summing up, specialized cytologists should be involved if ambiguous or inadequate results are encountered despite optimization of specimen-acquisition technique. A sensitivity and specificity exceeding 90 % can be achieved, but involves a well-balanced system of cell acquisition, preparation of smears, and further processing in the cytological laboratory. The Papanicolaou Society of Cytopathology has recently formulated guidelines for the terminology and nomenclature in pancreatobiliary cytology [29]. In the future, further consensual standards are mandatory for both endoscopists and cytologists to ensure a widespread, consistent quality of EUS-FNAC.

Competing interests: None

Institutions

- ¹ Department of Gastroenterology, Hepatology, and Gastroenterological Oncology, Bogenhausen Academic Teaching Hospital, Munich Municipal Hospital Holding, Munich, Germany
- ² Department of Pathology, Bogenhausen Academic Teaching Hospital, Munich Municipal Hospital Holding, Munich, Germany
- ³ Institute of Cytopathology, Hannover, Germany

References

- 1 Eltoun IA, Alston EA, Roberson J. Trends in pancreatic pathology practice before and after implementation of endoscopic ultrasound-guided fine-needle aspiration: an example of disruptive innovation effect? *Arch Pathol Lab Med* 2012; 136: 447–453
- 2 Dumonceau JM, Polkowski M, Larghi A et al. Indications, results, and clinical impact of endoscopic ultrasound (EUS)-guided sampling in gastroenterology: European Society of Gastrointestinal Endoscopy (ESGE) clinical guideline. *Endoscopy* 2011; 43: 897–912
- 3 Itoi T, Itokawa F, Sofuni A et al. Puncture of solid pancreatic tumors guided by endoscopic ultrasonography: a pilot study series comparing Trucut and 19-gauge and 22-gauge aspiration needles. *Endoscopy* 2005; 37: 362–366
- 4 Moller K, Papanikolaou IS, Toerner T et al. EUS-guided FNA of solid pancreatic masses: high yield of 2 passes with combined histologic-cytologic analysis. *Gastrointest Endosc* 2009; 70: 60–69
- 5 Takahashi K, Yamao K, Okubo K et al. Differential diagnosis of pancreatic cancer and focal pancreatitis by using EUS-guided FNA. *Gastrointest Endosc* 2005; 61: 76–79
- 6 Iglesias-Garcia J, Dominguez-Munoz E, Lozano-Leon A et al. Impact of endoscopic ultrasound-guided fine needle biopsy for diagnosis of pancreatic masses. *World J Gastroenterol* 2007; 13: 289–293
- 7 Varadarajulu S, Tamhane A, Eloubeidi MA. Yield of EUS-guided FNA of pancreatic masses in the presence or the absence of chronic pancreatitis. *Gastrointest Endosc* 2005; 62: 728–736; quiz 751, 753
- 8 Eloubeidi MA, Varadarajulu S, Desai S et al. A prospective evaluation of an algorithm incorporating routine preoperative endoscopic ultrasound-guided fine needle aspiration in suspected pancreatic cancer. *J Gastrointest Surg* 2007; 11: 813–819
- 9 Hwang CY, Lee SS, Song TJ et al. Endoscopic ultrasound guided fine needle aspiration biopsy in diagnosis of pancreatic and peripancreatic lesions: a single center experience in Korea. *Gut Liver* 2009; 3: 116–121
- 10 Fisher L, Segarajasingam DS, Stewart C et al. Endoscopic ultrasound guided fine needle aspiration of solid pancreatic lesions: performance and outcomes. *J Gastroenterol Hepatol* 2009; 24: 90–96
- 11 Fabbri C, Polifemo AM, Luigiano C et al. Endoscopic ultrasound-guided fine needle aspiration with 22- and 25-gauge needles in solid pancreatic masses: a prospective comparative study with randomisation of needle sequence. *Dig Liver Dis* 2011; 43: 647–652
- 12 Sakamoto H, Kitano M, Komaki T et al. Prospective comparative study of the EUS guided 25-gauge FNA needle with the 19-gauge Trucut needle and 22-gauge FNA needle in patients with solid pancreatic masses. *J Gastroenterol Hepatol* 2009; 24: 384–390
- 13 Varadarajulu S, Bang JY, Hebert-Magee S. Assessment of the technical performance of the flexible 19-gauge EUS-FNA needle. *Gastrointest Endosc* 2012; 76: 336–343
- 14 Bang JY, Hebert-Magee S, Trevino J et al. Randomized trial comparing the 22-gauge aspiration and 22-gauge biopsy needles for EUS-guided sampling of solid pancreatic mass lesions. *Gastrointest Endosc* 2012; 76: 321–327
- 15 Varadarajulu S, Fraig M, Schmulewitz N et al. Comparison of EUS-guided 19-gauge Trucut needle biopsy with EUS-guided fine-needle aspiration. *Endoscopy* 2004; 36: 397–401
- 16 Hartwig W, Schneider L, Diener MK et al. Preoperative tissue diagnosis for tumours of the pancreas. *Br J Surg* 2009; 96: 5–20
- 17 Yoshinaga S, Suzuki H, Oda I et al. Role of endoscopic ultrasound-guided fine needle aspiration (EUS-FNA) for diagnosis of solid pancreatic masses. *Dig Endosc* 2011; 23: 29–33
- 18 Jenssen C, Dietrich CF. Endoscopic ultrasound-guided fine-needle aspiration biopsy and trucut biopsy in gastroenterology – an overview. *Best Pract Res Clin Gastroenterol* 2009; 23: 743–759
- 19 Eloubeidi MA, Varadarajulu S, Desai S et al. Value of repeat endoscopic ultrasound-guided fine needle aspiration for suspected pancreatic cancer. *J Gastroenterol Hepatol* 2008; 23: 567–570
- 20 Tadic M, Kujundzic M, Stoos-Veic T et al. Role of repeated endoscopic ultrasound-guided fine needle aspiration in small solid pancreatic masses with previous indeterminate and negative cytological findings. *Dig Dis* 2008; 26: 377–382
- 21 Cherian PT, Mohan P, Douiri A et al. Role of endoscopic ultrasound-guided fine-needle aspiration in the diagnosis of solid pancreatic and peripancreatic lesions: is onsite cytopathology necessary? *HPB (Oxford)* 2010; 12: 389–395
- 22 Uehara H, Ikezawa K, Kawada N et al. Diagnostic accuracy of endoscopic ultrasound-guided fine needle aspiration for suspected pancreatic malignancy in relation to the size of lesions. *J Gastroenterol Hepatol* 2011; 26: 1256–1261
- 23 Lin F, Staerckel G. Cytologic criteria for well differentiated adenocarcinoma of the pancreas in fine-needle aspiration biopsy specimens. *Cancer* 2003; 99: 44–50
- 24 Mitsuhashi T, Ghafari S, Chang CY et al. Endoscopic ultrasound-guided fine needle aspiration of the pancreas: cytomorphological evaluation with emphasis on adequacy assessment, diagnostic criteria and contamination from the gastrointestinal tract. *Cytopathology* 2006; 17: 34–41
- 25 Petrone MC, Arcidiacono PG, Carrara S et al. Does cytotechnician training influence the accuracy of EUS-guided fine-needle aspiration of pancreatic masses? *Dig Liver Dis* 2012; 44: 311–314
- 26 Eloubeidi MA, Chen VK, Eltoun IA et al. Endoscopic ultrasound-guided fine needle aspiration biopsy of patients with suspected pancreatic cancer: diagnostic accuracy and acute and 30-day complications. *Am J Gastroenterol* 2003; 98: 2663–2668
- 27 Alsibai KD, Denis B, Bottlaender J et al. Impact of cytopathologist expert on diagnosis and treatment of pancreatic lesions in current clinical practice. A series of 106 endoscopic ultrasound-guided fine needle aspirations. *Cytopathology* 2006; 17: 18–26
- 28 Barreiros AP, Burmester E, Will U et al. Clinical procedures and outcomes of endoscopic fine needle biopsies (EUS-FNB): results of a German survey of the working group endoscopic ultrasound of the DEGUM. *Ultraschall Med* 2013; 34-WS_S17_02
- 29 Pitman MB, Centerno BA, Ali SZ et al. Standardized terminology and nomenclature for pancreatobiliary cytology: the Papanicolaou Society of Cytopathology guidelines *Diagn Cytopathol* 19. 02. 2014: DOI 10.1002/dc.23092