

Does Bilateral ITA Grafting Increase Perioperative Complications? Outcome of 6,476 Patients with Bilateral versus 5,020 Patients with Single ITA Bypass

Oliver Deutsch^{1,*} Laura Gansera^{2,*} Markus Wunderlich¹ Walter Eichinger¹ Brigitte Gansera¹

¹Department of Cardiovascular Surgery, Klinikum Bogenhausen, Munich, Germany

²Department of Cardiology, Klinikum Augsburg, Augsburg, Germany

* Both authors contributed equally to the work.

Address for correspondence Brigitte Gansera, MD, PhD, Department of Cardiovascular Surgery, Klinikum Bogenhausen, Engelschalkingerstrasse 77, Munich 81925, Germany (e-mail: brigitte_gansera@web.de).

Thorac Cardiovasc Surg 2016;64:188–194.

Abstract

Objectives Despite the superior patency of internal thoracic artery (ITA) grafting compared with saphenous veins, frequency of bilateral ITA (BITA) grafting in Europe is still approximately 10%. The aim of the present study was to compare the early outcome of patients receiving either BITA or single ITA (SITA) grafting.

Methods A total of 11,496 patients with isolated coronary artery bypass grafting (CABG), operated between January 1996 and December 2012, were analyzed retrospectively; 6,476 patients (mean age 65.2 years, 81.3% males) received BITA and 5,020 patients (mean age 66.6 years, 76.7% males) SITA grafting. Mean body mass index (BMI) was 27.2 versus 27.4, $p = 0.017$. Incidence of diabetes was 28.9 versus 28.4%, $p = 0.08$. Ejection fraction (EF) > 50 was 71.3% (BITA) versus 66.3% (SITA), $p < 0.001$. Elective operations were performed in 88.4% (BITA) versus 83.3% (SITA), and urgent/emergent surgery was necessary in 11.6% (BITA) versus 16.7% (SITA), $p < 0.001$.

Results Number of grafts was 3.76 (BITA) versus 3.06, $p < 0.001$. Duration of surgery (194.4 vs. 180.4 minutes) as well as X-clamp time (60.4 vs. 51.7 minutes) was prolonged for BITA, $p < 0.001$. Perioperative infarction rate revealed 3.2% (BITA) versus 3.6%, $p = 0.54$. Frequency of rethoracotomy due to bleeding was higher in the BITA group (3.8 vs. 2.1%), $p < 0.001$. Sternal instabilities occurred in 2.3% (BITA) versus 2.2%, $p = 0.749$. Duration of mechanical ventilation < 12 hours was 74.6 versus 77.1%, $p = 0.09$ and duration of in-hospital stay was 10.5 versus 10.4 days, $p = 0.68$. Thirty-day mortality was 2.4% (BITA) versus 3.0%, $p = 0.09$. Multivariate analysis identified prolonged duration of surgery, BMI > 30 , emergent operations, advanced age, and BITA grafting as predictor for sternal instabilities. EF $< 30\%$, advanced age plus emergency were associated with increased 30-day mortality.

Conclusion CABG using BITA can be performed routinely with good clinical results and low mortality. Compared with SITA grafting, bleeding complications were enhanced.

Keywords

- ▶ bilateral ITA
- ▶ outcome

received
February 23, 2015
accepted after revision
May 29, 2015
published online
August 13, 2015

© 2016 Georg Thieme Verlag KG
Stuttgart · New York

DOI <http://dx.doi.org/10.1055/s-0035-1558992>.
ISSN 0171-6425.

Introduction

Coronary artery bypass grafting (CABG) with bilateral internal thoracic artery (ITA) (BITA) grafting can improve long-term results in cardiac morbidity and mortality. Numerous clinical investigations could demonstrate an evident benefit in survival for BITA compared to single ITA (SITA) grafting.¹⁻³ An enhanced incidence of bleeding and wound complications, compared with patients with SITA remains an ongoing debate. Particularly, in patients suffering from diabetes, pedicled harvest of both ITAs is described to increase the risk of postoperative deep sternal wound infections.⁴ In contrary, other series, even including small patients cohorts, did not identify pedicled BITA harvesting as a risk factor for mediastinitis.⁵ The aim of the study was to compare the postoperative outcomes (30 days) in a large cohort of patients who had undergone CABG with BITA or SITA in situ or T-grafts, especially to identify patient-related risk factors, such as obesity and advanced age.

Methods

A total of 11,496 patients with isolated CABG, operated between January 1996 and December 2012, were analyzed retrospectively; 6,476 patients (mean age 65.2 years, 81.3% males) received BITA and 5,020 patients (mean age 66.6 years, 76.7% males) SITA grafting. Mean body mass index (BMI) was 27.2 versus 27.4, $p = 0.017$. Incidence of diabetes mellitus was 28.9 versus 28.4%, $p = 0.08$. Ejection fraction (EF) > 50 was 71.3% (BITA) versus 66.3% (SITA), $p < 0.001$. Elective operations were performed in 88.4% (BITA) versus 83.3% (SITA), and urgent/emergent surgery was necessary in 11.6% (BITA) versus 16.7% (SITA), $p < 0.001$. Patients who were under preoperative resuscitation were excluded from the study.

Statistical Analysis

Demographic and clinical data are presented as frequency distribution and simple percentages. Values of continuous variables are expressed as mean \pm standard deviation. In unmatched patient cohorts, univariate analysis of selected preoperative and postoperative discrete variables was accomplished by the chi-square test with the appropriate degrees of freedom or the Fishers exact test to assess the equality of proportions. Conditional logistic regression models with ordinal variables were used to determine the independent effects of selected demographic/clinical variables on hospital complications and mortality. All analyses were performed with SPSS (Statistical Package for Social Science version 18, SPSS Inc., Chicago, Illinois). A p -value < 0.05 was considered to indicate statistical significance.

Results

Demographic data are depicted in ►Table 1. Operative data are depicted in ►Table 2. Perioperative complications and mortality are depicted in ►Table 3. The multivariate regression analysis for variables influencing complications and mortality is given in ►Table 4. ►Table 5 shows a division of sternal instabilities related to the surgical ITA-harvesting approach (pedicled or skeletonized), different graft technique (in situ or T-graft) in dependence of diabetes/no diabetes. Number of grafts was higher in the BITA group, as well as duration of surgery and X-clamp time was prolonged for BITA grafting. Perioperative infarction rate revealed no statistical significant difference between both groups. Frequency of rethoracotomy due to bleeding was significantly higher in the BITA group. Sternal instabilities occurred in 2.3% (BITA) versus 2.2% (SITA), $p = 0.749$. Sternal instabilities without

Table 1 Demographic data and preoperative variables

Variables	SITA	BITA	p-Value
Patients, n (%)	5,020 (100.0)	6,476 (100.0)	
Sex, n (%)			
Male	3,850 (76.7)	5,265 (81.3)	< 0.001
Female	1,170 (23.3)	1,211 (18.7)	
Age, mean \pm SD, y	66.6 \pm 9.2	65.2 \pm 9.3	< 0.001
Body mass index, mean \pm SD	27.4 \pm 5.2	27.2 \pm 3.2	0.017
EF (%)			
EF $> 50\%$	66.3%	71.3%	< 0.001
EF 30–50%	28.8%	24.7%	
EF $< 30\%$	4.9%	4.1%	
Elective/urgent/emergent operation			
Elective (%)	83.3	88.4	< 0.001
Urgent (%)	9.6	6.3	
Emergent (%)	7.8	5.2	

Abbreviations: BITA, bilateral internal thoracic artery; EF, ejection fraction; SD, standard deviation; SITA, single internal thoracic artery.

Table 2 Comparison of operative data

Variables	SITA	BITA	p-Value
Number of grafts (mean ± SD)	3.06 ± 0.92	3.76 ± 0.92	< 0.0001
Duration of surgery (min, mean)	180.4 ± 55.8	194.4 ± 48.5	< 0.0001
Bypass time (min, mean)	79.8 ± 33.5	84.8 ± 28.3	< 0.0001
X-clamp time (min, mean)	51.7 ± 22.2	60.4 ± 18.9	< 0.001

Abbreviations: BITA, bilateral internal thoracic artery; SD, standard deviation; SITA, single internal thoracic artery.

infection were 0.6% (SITA) versus 1.0% (BITA), $p = 0.03$, but incidence of sternal instabilities with infection was lower in the BITA group, 1.6% (SITA) versus 1.3% (BITA), $p = 0.26$. Division of sternal instabilities related to the surgical ITA-harvesting approach (pedicled or skeletonized) and different graft technique (in situ or T-graft) result in a comparable incidence for patients with or without diabetes in the pedicled SITA in situ group (2.9 vs. 2.8%, $p = 0.838$); but in a higher incidence of sternal complications in patients with diabetes of the pedicled in situ BITA group (0.9 vs. 2.4%, $p = 0.001$). In patients receiving skeletonized SITA grafts ($n = 827$ of all 5,020 SITAs), or skeletonized BITA T-grafts ($n = 263$ of all 6,476 BITAs), sternal instabilities occur in a higher frequency compared with the former group, but without statistical differences between patients suffering from diabetes or not.

Duration of mechanical ventilation < 12 and > 24 hours were comparable between the groups. Mean period of in-hospital stay was 10.5 versus 10.4 days. Thirty-day mortality was higher in the SITA group, without statistically significance. Multivariate analysis identified prolonged duration of surgery, BMI > 30, advanced age, and BITA grafting as predictors for sternal instabilities without infections. Sternal wound infections were strongly associated with prolonged duration of cardiopulmonary bypass. EF < 30%, advanced age, emergent operations, duration of surgery, bypass, and X-clamp time were independent predictors for increased 30-day mortality.

Discussion

Multiple cohort studies have demonstrated that long-term outcome after CABG can be improved remarkably when both ITAs are used for revascularization.^{1,3,6-8} There is evidence about superior patency of ITA grafting to saphenous veins.⁹ The use of the left ITA bypass preferentially to the left anterior descending artery (LAD) has become surgical standard, providing reduced cardiac events, and enhanced survival rates, compared with patients receiving isolated vein grafts. Although the evidence that the long-term outcome after CABG using both ITAs results in convincing beneficial advantages,^{1,3,6-8} the frequency of BITA grafting in Europe is still approximately 10%.¹⁰ The National US Database of the Society of Thoracic Surgeons reports that < 4% of patients, who underwent CABG currently receive the benefits of BITA grafting.¹¹ There is an ongoing discussion of pros and cons for this surgical approach. Although there is consensus that BITA grafting can improve long-term survival,³ the reluctance to use BITA seems to be based on the fear of deep sternal wound infections, especially among diabetic patients. Some reports describe a higher incidence of wound-healing and bleeding complications after BITA grafting,^{4,12} in particular, when pedicled harvesting is performed in diabetic patients.^{4,12,13} A large series, consisting of 126,235 diabetic patients (122,465 SITA and 3,770 BITA) published by Deo et

Table 3 Comparison of postoperative complications

Variables	SITA	BITA	p-Value
Myocardial infarction (%)	3.6	3.2	0.54
Rethoracotomy due to bleeding (%)	2.1	3.8	< 0.001
Sternal instabilities, all (%)	2.2	2.3	0.749
Sternal instabilities without infection (%)	0.6	1.0	0.03
Sternal instabilities with infection (%)	1.6	1.3	0.26
Mechanical ventilation < 12 h (%)	77.1	74.6	0.09
Mechanical ventilation > 24 h (%)	8.6	6.7	0.07
Reintubation (%)	3.7	3.1	0.002
Tracheotomy (5%)	0.7	0.5	0.002
In-hospital stay, d (mean ± SD)	10.4 ± 9.5	10.5 ± 9.3	0.68
30-day mortality (%)	3.0	2.4	0.09

Abbreviations: BITA, bilateral internal thoracic artery; SD, standard deviation; SITA, single internal thoracic artery.

Table 4 Multivariate regression analysis for variables influencing complications and mortality^a

Variables	p-Value	95% confidence interval	
		Lower bound	Upper bound
Sternal instability with infection			
Age	0.630	– 0.021	0.013
BMI	0.515	– 0.016	0.032
LVEF < 30%	0.572	– 1.063	0.588647
Emergency	0.843	– 0.245	0.300
Duration of operation	0.110	– 0.008	0.01
Bypass time	0.032	0.001	0.020
X-clamp time	0.810	– 0.015	0.012
Use of BITA	0.317	– 0.160	0.493
Sternal instability without infection			
Age	0.000	0.037	0.087
BMI	0.034	0.001	0.039
Emergency	0.026	0.040	0.624
LVEF < 30%	0.065	– 0.042	1.418
Duration of operation	0.000	0.004	0.012
Bypass time	0.381	– 0.018	0.007
X-clamp time	0.954	– 0.017	– 0.108
Use of BITA	0.015	– 1.017	– 0.108
30-day survival			
Age	0.000	– 0.066	– 0.037
BMI	0.780	– 0.015	0.20
Emergency	0.000	– 0.784	– 0.492
Duration of operation	0.000	– 0.011	– 0.007
EF < 30%	0.000	0.779	1.502
Bypass time	0.000	– 0.023	– 0.12
X-clamp time	0.000	0.012	0.028
Use of BITA	0.983	– 0.247	0.252

Abbreviations: BITA, bilateral internal thoracic artery; BMI, body mass index; EF, ejection fraction; LVEF, left ventricular ejection fraction; SD, standard deviation.

Note: Bold values show *p* values <0.05.

^aPLUM (polytomous universal model)—ordered logistic regression model.

al,⁴ identified the risk of deep sternal wound infections to be minimized when performing ITA harvesting in a skeletonized manner. With this approach, patients after BITA grafting had a similar risk of deep sternal infections compared with SITA grafting.

A comparison of these results to our own seems to be difficult, as our cohort contributes in only 28% of patients suffering from diabetes. Nevertheless, incidence of deep wound infection was at least 3.1% for BITA and 1.6% for the SITA in the mentioned investigation of 3,770 BITAs,⁴ whereas deep sternal wound infection occurred in only 1.3% of 6,476 BITA grafts in our own series. Moreover, we performed a mixed approach of pedicled harvesting (until the year 2010) and skeletonized technique (since the year 2010); therefore, simple comparison to the results in the literature seems

problematic. After the performance of a subgroup analysis within our collective, the use of pedicled BITA harvest did lead to increased sternal wound complication rates than the use of skeletonized BITAs (2.4 vs. 3.3%), even in diabetic patients (3.9 vs. 5.2%).

Nevertheless, patients suffering from diabetes who received pedicled in situ BITA grafts were identified to experience significant higher rates of sternal instabilities compared with those without diabetes (3.9 vs. 2.4%, *p* = 0.001).

Choo et al⁵ identified pedicled BITA grafting not to be associated with an increased incidence of infectious sternal complications, albeit in only 162 patients with or without existing diabetes. Randomized clinical trials concerning this issue are rarely available, most of the studies are of observational character or deal with small collectives. Taggart et al¹²

Table 5 Sternal wound complications

Variables	Diabetes	No diabetes	p-Value
SITA (<i>n</i> = 4,193) and BITA (<i>n</i> = 6,213), pedicled, in situ grafts (1996–2010), related to diabetes			
Sternal instabilities (all) BITA (%)	3.9%	2.4%	0.001
Sternal instabilities (all) SITA (%)	2.9%	2.8%	0.838
SITA (<i>n</i> = 827) and BITA (<i>n</i> = 263), skeletonized, T-grafts (2010–2012), related to diabetes			
Sternal instabilities (all) BITA (%)	5.2%	3.4%	0.463
Sternal instabilities (all) SITA (%)	4.4%	4.0%	0.852

Abbreviations: BITA, bilateral internal thoracic artery; SITA, single internal thoracic artery.

published the results of the unique randomized investigation, a multicenter study including 1,554 patients with SITA grafting and 1,548 patients with BITA grafting. Perioperative complications as well as results after 1 year were similar between both groups in terms of mortality, infarction rate, and repeated revascularization. A small increase for the need of sternal wound reconstructions was observed (0.6% SITA vs. 1.9% BITA), around half of these patients had a history of diabetes in comparison with, similar to our portion, around one-quarter of all patients in the whole trial. Advocates of BITA grafting regularly survey larger cohorts than their opponents. One of the most important recently published study concerning this subject was performed by Dorman et al⁶: a 30-year follow-up of propensity score-matched cohorts, which consist of diabetic patients receiving either SITA (*n* = 414) or BITA (*n* = 414). This sophisticated and well-done study identified BITA grafting providing enhanced survival without any increase in perioperative morbidity or mortality. Although ITA harvest was performed exclusively in a pedicled manner, incidence of deep sternal infections was nearly similar between SITA and BITA groups. These data are supported by numerous prior investigations.^{5,14–16}

Other groups strongly recommend the skeletonized approach to preserve sternal blood flow.^{4,13,17–19} Kamiya et al¹⁷ could verify in a small collective (*n* = 24) that the damage of the tissue microcirculation in the middle and lower retro-sternal area was significantly less after ITA skeletonization compared with that after the pedicled ITA-harvesting technique. In this context, the authors of the present trial would like to emphasize that in case of pedicled ITA harvesting, the pedicle was always kept very small.

Toumpoulis et al¹⁹ analyzed 140 articles related to this subject. Twenty four articles were identified to give best evidence. In general, BITA grafting carries a 2.5- to 5-fold higher risk for mediastinitis, which does raise up to 10% in diabetic patients, but can be reduced to 0.4–2.6% when skeletonized techniques were used. The data of Momin et al²⁰ do not support the perception that BITA grafting increases wound complications. This trial enfolds a 10-year period of 95 diabetic patients receiving BITA, and deep sternal wound infection was 1.2 versus 3.2% (*p* = 0.36).

Interestingly, reoperation for bleeding in the 414 patients of Dorman cohort⁶ was lower in the BITA group (2.4 vs. 3.1%, *p* = 0.513). Our data do not confirm these results, as frequen-

cy of rethoracotomy due to bleeding was significantly higher in the BITA group (3.8 vs. 2.1%, *p* < 0.001). This fact might be explainable by the unselected collective in our large series, as we included all patients, for example emergent cases, who were under antiplatelet therapy and, in a not negligible number, patients with obesity (BMI mean was at least 27.2). On the contrary, one of our own previous studies,¹⁶ consisting of 2,251 patients with extreme obesity (BMI 30–50), could demonstrate that obesity per se is not a risk factor for increased bleeding or wound-healing complications, but the combination of obesity, diabetes mellitus, and BITA grafting increases the risk for sternal complications. Nevertheless, the blood loss of mean 1,125.5 ± 850.3 mL (*p* < 0.0001) in the BITA group seems to be high. In this context, it might be of interest, whether chest-tube output has changed after conversion from pedicled to skeletonized ITA grafts. Other groups likewise describe slightly higher bleeding complications after BITA grafting,^{12,20} commonly associated with a higher number of anastomoses and prolonged X-clamp time.²⁰

As expected, perfusion and X-clamp times were significantly prolonged in the BITA group, in our cohort as well as in numerous other publications, and had a significant influence on mortality in the present study. These results compare favorably to other investigations,⁶ which clearly defined prolonged perfusion or X-clamp time as an independent risk factor for increased morbidity or mortality. One cause of prolonged perfusion- and X-clamp time in our series might be the significant higher number of anastomoses within the BITA group. In this context, one will raise up the justified question about a selection bias in our cohort. Doubtlessly, patients with the need for single bypass or, for example, grafts to the LAD/diagonal branch understandably cannot receive BITA grafting. That means that selection criteria in a predetermined extent are subjected to restrictions depending on vessel status and morphology.

Comparing mortality is another difficult issue. Study designs are varying widely. For example, within the randomized trial of Taggart et al¹² mortality was 1.2% for BITA as well as for SITA, therefore much lower than in our series. Around 40% were performed off-pump, whereas we mostly performed on-pump procedures. Instead of EF, New York Heart Association (NYHA), NY or Canadian Cardiovascular Society (CCS) classification was analyzed and different

categories for elective/urgent/emergent operations were not presented.

Other series include or exclude emergent operations, a fact that self-evidently must lead to different survival rates. Momin et al²⁰ reported a higher 30-day mortality of 5.0 versus 6.1% ($p = 0.46$), despite in a comparatively low-aged population (mean 62 years in the BITA group), but included urgent operations in more than 33%. EF was $< 30\%$ in around 10%, more than twofold higher compared with our series, which may easily explain the poorer results. In-hospital mortality within the propensity score-pair-matched collective of diabetic patients published by Dorman et al⁶ was 4.6% (SITA) versus 3.1% (BITA). In the large review of Deo et al,⁴ certainly in diabetic patients, early mortality was 2.5% in the SITA cohort and 2.3% in the BITA cohort ($p = 0.8$), therefore analogical to our results. Even, a straight comparability of the numerous studies seems to be difficult, the main focus of the present study is on early postoperative outcome, and we report, to our knowledge, with the experience of 6,476 patients receiving BITA grafting, the largest single-center study concerning this subject.

Limitations

The retrospective, nonrandomized character might be regarded as a substantial limitation of the present study. Changes of surgical strategies, concerning ITA harvesting lead to a "historical bias." The fact that most of the patients (from 1996 until the year 2010) received pedicled in situ grafts, but only a small number (from the year 2010 to 2012) received skeletonized T-grafts, must result in subgroups, which are hard to compare. Therefore, these results are of a less meaningful statistic power, restricted clinical relevance, and limited validity.

The within the absolute values slight, but nonnegligible bias concerning statistically significant differences in terms of age, BMI, EF, and elective versus emergent cases between both groups might be considered as a result of the large number of patients.

Conclusion

CABG using BITA can be performed routinely in large patient cohorts with good clinical results and low mortality. Compared with SITA grafting, bleeding complications were enhanced. Prolonged duration of surgery, advanced age, emergent operations, BITA grafting, and obesity could be identified as predictors for sternal instabilities. Sternal wound infections were associated with duration of cardiopulmonary bypass, but not BITA grafting per se, neither obesity had any influence on sternal wound infections. Number of sternal instabilities was higher in the subgroup receiving skeletonized BITA T-grafts compared with those with pedicled in situ BITA grafts. Pedicled BITA grafting plus diabetes was associated with a higher incidence of sternal instabilities than pedicled BITA grafting without diabetes. Advanced age, EF $< 30\%$, duration of surgery, prolonged bypass, and X-clamp time plus emergency oper-

ation status were independent risk factors for increased 30-day mortality.

Acknowledgments

The first two authors (O.D. and L.G.) contributed equally to this article. O.D. performed the statistics and evaluation. L. G. prepared the setting, content of the work, and wrote the article.

References

- 1 Taggart DP, D'Amico R, Altman DG. Effect of arterial revascularisation on survival: a systematic review of studies comparing bilateral and single internal mammary arteries. *Lancet* 2001; 358(9285):870–875
- 2 Rizzoli G, Schiavon L, Bellini P. Does the use of bilateral internal mammary artery (IMA) grafts provide incremental benefit relative to the use of a single IMA graft? A meta-analysis approach. *Eur J Cardiothorac Surg* 2002;22(5):781–786
- 3 Kurlansky PA, Traad EA, Dorman MJ, Galbut DL, Zucker M, Ebra G. Thirty-year follow-up defines survival benefit for second internal mammary artery in propensity-matched groups. *Ann Thorac Surg* 2010;90(1):101–108
- 4 Deo SV, Shah IK, Dunlay SM, et al. Bilateral internal thoracic artery harvest and deep sternal wound infection in diabetic patients. *Ann Thorac Surg* 2013;95(3):862–869
- 5 Choo SJ, Lee SK, Chung SW, et al. Does bilateral pedicle internal thoracic artery harvest increase the risk of mediastinitis? *Yonsei Med J* 2009;50(1):78–82
- 6 Dorman MJ, Kurlansky PA, Traad EA, Galbut DL, Zucker M, Ebra G. Bilateral internal mammary artery grafting enhances survival in diabetic patients: a 30-year follow-up of propensity score-matched cohorts. *Circulation* 2012;126(25):2935–2942
- 7 Lytle BW, Loop FD. Superiority of bilateral internal thoracic artery grafting: it's been a long time comin'. *Circulation* 2001;104(18): 2152–2154
- 8 Rubens FD, Boodhwani M. Skeletonization of the internal thoracic artery for coronary artery bypass grafting. *Curr Opin Cardiol* 2009; 24(6):559–566
- 9 Gansera B, Schmidtler F, Angelis I, Kiask T, Kemkes BM, Botzenhardt F. Patency of internal thoracic artery compared to vein grafts - postoperative angiographic findings in 1189 symptomatic patients in 12 years. *Thorac Cardiovasc Surg* 2007;55(7): 412–417
- 10 Bridgewater B. Cardiac registers: the adult cardiac surgery register. The Society for Cardiothoracic Adult Cardiac Surgery Database. *Heart* 2010;96:1441–1443
- 11 Tabata M, Grab JD, Khalpey Z, et al. Prevalence and variability of internal mammary artery graft use in contemporary multivessel coronary artery bypass surgery. *Circulation* 2009;120:935–940
- 12 Taggart DP, Lees B, Gray A, Altman DG, Flather M, Channon K. ART Investigators. A randomized trial to compare survival following bilateral versus single internal mammary grafting in coronary revascularization. *Trials* 2006;7:7–17
- 13 De Paulis R, de Notaris S, Scaffa R, et al. The effect of bilateral internal thoracic artery harvesting on superficial and deep sternal infection: The role of skeletonization. *J Thorac Cardiovasc Surg* 2005;129(3):536–543
- 14 Gansera B, Günzinger R, Angelis I, Eichinger W, Neumaier-Prauser P, Kemkes BM. End of the millenium- end of the single thoracic artery graft. Two thoracic arteries- standard for the next millenium? Early clinical results and analysis of risk factors in 1487 patients with bilateral internal thoracic artery grafts. *Thorac Cardiovasc Surg* 2001;49:10–15

- 15 Gansera B, Schmidtler F, Gillrath G, et al. Does bilateral ITA grafting increase perioperative complications? Outcome of 4462 patients with bilateral versus 4204 patients with single ITA bypass. *Eur J Cardiothorac Surg* 2006;30(2):318–323
- 16 Syrakas CA, Neumaier-Prauser P, Angelis I, Kiask T, Kemkes BM, Gansera B. Is extreme obesity a risk factor for increased in-hospital mortality and postoperative morbidity after cardiac surgery? Results of 2251 obese patients with BMI of 30 to 50. *Thorac Cardiovasc Surg* 2007;55(8):491–493
- 17 Kamiya H, Akhyari P, Martens A, Karck M, Haverich A, Lichtenberg A. Sternal microcirculation after skeletonized versus pedicled harvesting of the internal thoracic artery: a randomized study. *J Thorac Cardiovasc Surg* 2008;135(1):32–37
- 18 Boodhwani M, Lam BK, Nathan HJ, et al. Skeletonized internal thoracic artery harvest reduces pain and dysesthesia and improves sternal perfusion after coronary artery bypass surgery: a randomized, double-blind, within-patient comparison. *Circulation* 2006;114(8):766–773
- 19 Toumpoulis IK, Theakos N, Dunning J. Does bilateral internal thoracic artery harvest increase the risk of mediastinitis? *Interact Cardiovasc Thorac Surg* 2007;6(6):787–791
- 20 Momin AU, Deshpande R, Potts J, et al. Incidence of sternal infection in diabetic patients undergoing bilateral internal thoracic artery grafting. *Ann Thorac Surg* 2005;80(5):1765–1772, discussion 1772