


F.-X. WANG, J.-Y. DU, H.-B. WANG, P.-L. ZHANG, G.-B. ZHANG, K.-Y. YU, X.-Z. ZHANG, X.-T. AN, Y.-X. CAO, C.-A. FAN\* (LANZHOU UNIVERSITY AND COLLABORATIVE INNOVATION CENTER OF CHEMICAL SCIENCE AND ENGINEERING (TIANJIN), P. R. OF CHINA)  
 Total Synthesis of *Lycopodium* Alkaloids Palhinine A and Palhinine D  
*J. Am. Chem. Soc.* **2017**, *139*, 4282–4285.

## Synthesis of Palhinine A and D



**Significance:** Palhinine A and D are members of the *Lycopodium* alkaloid family isolated from the whole plant of *Palhinhaea cernua* L. Fan and co-workers describe the first total syntheses relying on a microwave-assisted regio- and stereoselective nitron-alkene [3+2] cycloaddition as the key step to access the nine-membered azanone ring system.

**Comment:** Treatment of hydroxylamine **E** with formalin in  $\text{CH}_2\text{Cl}_2$  resulted in the formation of nitron **F**, which underwent an intramolecular 1,3-dipolar cycloaddition to give isoxazolidine **G**. N-Alkylation followed by reductive N–O cleavage gave access to the challenging azanone ring system, key structural motif of palhinine A and D.

**SYNFACTS Contributors:** Erick M. Carreira, Marco Brandstätter  
 Synfacts 2017, 13(06), 0565 Published online: 16.05.2017  
 DOI: 10.1055/s-0036-1590422; Reg-No.: C02317SF

2017 © THIEME STUTTGART • NEW YORK

Category

Synthesis of Natural Products and Potential Drugs

Key words

palhinine A and D

*Lycopodium* alkaloids

nitron

isooxazolidine

1,3-dipolar cycloaddition

Synfact  
 of the month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.