





A. R. HAJIPOUR*, F. REZAEI, Z. KHORSANDI* (ISFAHAN UNIVERSITY OF TECHNOLOGY, IRAN AND UNIVERSITY OF WISCONSIN, MADISON, USA)
 Pd/Cu-Free Heck and Sonogashira Cross-Coupling Reaction by Co Nanoparticles Immobilized on Magnetic Chitosan as Reusable Catalyst
Green Chem. **2017**, *19*, 1353–1361.


Heck and Sonogashira Reactions on Magnetic Cobalt Nanoparticles



Selected examples:



Selected examples:



Significance: A magnetic, chitosan-supported, methyl salicylate–cobalt complex (Co-MS@MNPs/CS) was prepared as shown in eq. 1. Co-MS@MNPs/CS catalyzed the Heck reaction of aryl halides with terminal olefins to give the corresponding internal alkenes in $\leq 92\%$ yield (eq. 2, 26 examples). Co-MS@MNPs/CS also promoted the Sonogashira reaction of aryl iodides or bromides with arylacetylenes to give the corresponding diarylacetylenes in $\leq 80\%$ yield (eq. 3, 15 examples).

Comment: The catalyst was characterized by means of FT-IR, TGA, EA, XRD, FE-SEM, SEM-EDX, TEM, magnetization curve, and ICP analyses. In the Heck reaction of iodobenzene with methyl acrylate, the catalyst was recovered and reused four times without loss of its catalytic activity.

SYNFACTS Contributors: Yasuhiro Uozumi, Shuo Yan
 Synfacts 2017, 13(06), 0659 Published online: 16.05.2017
 DOI: 10.1055/s-0036-1590479; Reg-No.: Y05717SF