G. POVIE, Y. SEGAWA*, T. NISHIHARA, Y. MIYAUCHI, K. ITAMI* (JAPAN SCIENCE AND TECHNOLOGY AGENCY, EXPLORATORY RESEARCH FOR ADVANCED TECHNOLOGY, NAGOYA, NAGOYA UNIVERSITY, AND KYOTO UNIVERSITY, JAPAN)

Synthesis of a Carbon Nanobelt

Science 2017, 356, 172-175.

Nanobelt Mission Accomplished: A Loop of 12 Fused Edge-Sharing Benzenes

Significance: The synthesis of a fully fused carbon nanobelt (7) — a belt segment of a (6,6)carbon nanotube — has been a long-lasting goal in materials science. Segawa, Itami, and co-workers have now reported on the successful bottom-up synthesis of such a carbon nanobelt, a closed loop of 12 fully fused edge-sharing benzene rings. The synthesis starts with several consecutive Wittig reactions. In the final step, a Ni(cod)₂-mediated aryl–aryl coupling reaction prompts the formation of the new fully fused arene system.

SYNFACTS Contributors: Timothy M. Swager, Lukas Zeininger Synfacts 2017, 13(07), 0697 Published online: 19.06.2017 **DOI:** 10.1055/s-0036-1590570; **Reg-No.:** S05417SF

Comment: The authors studied in detail the optoelectronic and structural properties of this 'shortest armchair carbon nanotube'. Besides verifying the rigid cylindrical shape of this carbon nanobelt through X-ray crystallography, its photophysical characterization combined with theoretical calculations confirmed the structural rigidity of this fully conjugated arene system. The synthesis of this belt of 12 edge-sharing benzene rings may pave the way towards a precisely controlled singlechirality bottom-up synthesis of (6,6)carbon nanotubes.

Category

Synthesis of Materials and Unnatural Products

Key words

aryl-aryl coupling
Wittig reaction
carbon nanobelt



697