

Do Knee Bracing and Delayed Weight Bearing Affect Mid-Term Functional Outcome after Anterior Cruciate Ligament Reconstruction?

Riccardo Di Miceli¹ Carlotta Bustos Marambio¹ Alessandro Zati¹ Maria Grazia Benedetti¹

Messandro Zati¹ Roberta Monesi¹

¹ SC Medicina Fisica e Riabilitativa, Istituto Ortopedico Rizzoli di Bologna, Bologna, Italy Address for correspondence Maria Grazia Benedetti, MD, SC Medicina Fisica e Riabilitativa, Istituto Ortopedico Rizzoli di Bologna, Via Pupilli 1, Bologna 40136, Italy (e-mail: benedetti@ior.it).

Joints 2017;5:202-206.

Abstract	Purpose The aim of this study was to assess the effect of knee bracing and timing of full weight bearing after anterior cruciate ligament reconstruction (ACLR) on functional outcomes at mid-term follow-up. Methods We performed a retrospective study on 41 patients with ACLR. Patients were divided in two groups: ACLR group, who received isolated ACL reconstruction and ACLR-OI group who received ACL reconstruction and adjunctive surgery. Information about age at surgery, bracing, full or progressive weight bearing permission after surgery were collected for the two groups. Subjective IKDC score was obtained at follow-up. Statistical analysis was performed to compare the two groups for IKDC score. Subgroup analysis was performed to assess the effect of postoperative regimen (knee bracing and weight bearing) on functional outcomes. Results The mean age of patients was 30.8 ± 10.6 years. Mean IKDC score was
	87.4 ± 13.9 . The mean follow-up was 3.5 ± 1.8 years. Twenty-two (53.7%) patients underwent ACLR only, while 19 (46.3%) also received other interventions, such as meniscal repair and/or collateral ligament suture. Analysis of overall data showed no differences between the groups for IKDC score. Patients in the ACLR group exhibited a significantly better IKDC score when no brace and full weight bearing after 4 weeks from surgery was prescribed in comparison with patients who worn a brace and had delayed full weight bearing. No differences were found with respect to the use of brace
Keywords	and postoperative weight bearing regimen in the ACLR-OI group.
 rehabilitation 	Conclusion Brace and delayed weight bearing after ACLR have a negative influence on
 anterior cruciate 	long-term functional outcomes. Further research is required to explore possible
ligament	differences in the patients operated on ACLR and other intervention with respect to
 weight-bearing 	strategies
 weight-bearing functional outcome 	sualegies.
	Level of Evidence Level III, recrospective observational study.

Introduction

Anterior cruciate ligament (ACL) rupture is a common injury, mainly affecting young, physically active individuals.¹ It is often caused by a noncontact mechanism, as sudden deceleration combined with changing in direction, pivoting, or landing with the knee in nearly full extension after a jump.² ACL lesion is usually combined with other injuries, typically to collateral ligaments, subchondral bone, and menisci.³ ACL reconstruction (ACLR) is the predominant treatment in

published online September 15, 2017 DOI https://doi.org/ 10.1055/s-0037-1606617. ISSN 2282-4324. Copyright © 2017 Georg Thieme Verlag License terms KG Stuttgart · New York



current practice, and the relative merits of hamstring and patellar tendon (PT) grafts have been well established.⁴ Rehabilitation interventions after surgery must be both safe and effective, with the ultimate goal of returning the patients to their activities often at a high functional level.

Successful outcomes have been consistently achieved with the rehabilitation principles of early weight bearing, using a combination of weight bearing and non-weightbearing exercise focused on guadriceps and lower extremity strength recover and meeting specific objective requirements for return to activity.⁵ However, high variability in the composition and time ranges of rehabilitation components has been described. Consensus over the most effective rehabilitation protocol is lacking,⁶ especially on ACL injuries combined with collateral ligament and meniscal injuries in adults.⁷ This may lead to confusion among patients and therapists regarding the best protocol to adopt.⁸ In particular, knee bracing and timing for weight bearing have been widely investigated.^{1,9–15} Previous systematic reviews found that accelerated rehabilitation, early weight bearing, and early range of motion (ROM) gaining are possibly safe and beneficial to patient outcomes.^{9,11} Recent literature showed no benefits of knee bracing after ACLR with respect to pain, stability, ROM, and protection from additional injuries.¹²⁻¹⁵

The aim of the present study was to assess the effects of knee bracing and timing of full weight bearing after ACLR on the functional outcomes at mid-term follow-up.

Methods

This study was designed as a retrospective observational study. The study was approved by the local ethical committee.

Patients aged between 15 and 70 years and residents of Italy, who underwent ACLR with semitendinosus and gracilis tendon grafts from 2008 to 2013 and who, 4 weeks after surgery, received a rehabilitation program for 2 weeks at our Institution were considered eligible for the study.

The rehabilitation program during the 2 weeks was based on a combination of ROM exercises, muscle strengthening, gait training, physical therapy modalities, muscles stretching, and balance and proprioception exercises.

At the beginning of the rehabilitation program, different prescriptions were provided by orthopaedic surgeons in terms of use of brace to immobilize the knee in extension and timing of weight bearing.

One hundred eight patients were contacted by means of a letter. Forty-one patients accepted to participate in the study. Their clinical records were retrieved and information about age at surgery, bracing, and timing to partial and/or full weight bearing after surgery were collected. Subjective IKDC score was obtained by directly interviewing the patient at follow-up.

The patients were divided into two groups: ACLR and ACLR-other intervention (ACLR-OI) groups. Differences between the groups for age at surgery, duration of follow-up, and IKDC score were analyzed using ANOVA for normally distributed data, otherwise with the Mann–Whitney U-test.

The subgroup analysis was performed to assess the differences in IKDC score according to knee bracing (yes/no) and weight-bearing regimen (full/progressive) in the two groups with the Mann–Whitney U-test using Monte Carlo method for small samples. An analysis of the effect of postoperative regimen (bracing and weight bearing) on IKDC score at follow-up regardless of surgical treatment was performed with the Kruskal–Wallis test and the Mann–Whitney U-test for post-hoc analysis using the Monte Carlo method for small samples and the Sidak correction for multiple comparisons. The significance was considered for p < 0.05.

Results

The mean age of the study population (41 patients) was 30.8 ± 10.6 years. The mean IKDC score was 87.4 ± 13.9 . The mean follow-up period from surgery was 3.5 ± 1.8 years. Twenty-two patients (53.7%) underwent ACL reconstruction only (ACL group), while 19 (46.3%) also received other interventions, such as total or partial meniscectomy, collateral ligament suture, and microfractures (ACL-OI group). Analysis of data showed no differences between the two groups in terms of age at the time of surgery, duration of follow-up, and IKDC score (**-Table 1**).

Twenty patients (48.8%) used knee bracing for 4 weeks, while 21 (51.2%) did not. All the patients had the prescription of partial weight bearing for 4 weeks postoperatively. Thirty-two patients (78%) were allowed to regain full weight bearing (FWB) within 2 weeks of rehabilitation, while nine patients (22%) were prescribed to continue progressive weight bearing (PWB) up to the orthopaedic visit after the rehabilitation period. In the ACLR group, 9 patients (40.9%) received a brace and 13 (59.1%) did not; 16 patients (72.7%) had FWB and 6 (27.3%) had PWB. In the ACL-OI group, 12 patients (63.2%) received a brace and 7 (36.8%) did not; 16 patients (84.2%) had FWB and 3 (15.8%) had PWB.

Subgroup analysis according to postoperative regimen demonstrated significant differences only for the ACLR group in favor of no bracing and FWB (p = 0.001 and p = 0.018, respectively), while no differences were found in ACLR-OI group. The overall results demonstrated that patients without brace and FWB had better IKDC score than subjects with brace and PWB, with a statistically significant difference (p = 0.017 and p = 0.036, respectively; **~Table 2**).

Table 1 Comparison between groups

Variables	Group (N)	p-Value	
	ACLR (22)	ACLR-OI (19)	
Age (mean \pm SD)	31.4 ± 10.8	30.2 ± 10.5	ns
Follow-up, y (mean \pm SD)	3.4 ± 1.6	3.6 ± 2	ns
IKDC score (mean \pm SD)	89	85.6	ns

Abbreviations: ACLR, anterior cruciate ligament reconstruction; ACLR-OI, anterior cruciate ligament reconstruction + other interventions; IKDC, International Knee Documentation Committee; SD, standard deviation; ns, nonsignificant.

Groups (N)	Variables	Brace		p-Value	WB		p-Value
		No	Yes		Progressive	Full	
ACLR (22)	N (%)	13 (59.1%)	9 (40.9%)		6 (27.3%)	16 (72.7%)	
	IKDC score (mean \pm SD)	95.1 ± 8	80.2 ± 15.6	0.001	78.0 ± 15.8	93.1 ± 10.5	0.018
ACLR-OI (19)	N (%)	7 (36.8%)	12 (63.2%)		3 (15.8%)	16 (84.2%)	
	IKDC score (mean \pm SD)	83.3 ± 19.1	86.9 ± 11.7	ns	84.7 ± 12.8	85.7 ± 15.1	ns
Overall (41)	N (%)	20 (48.8%)	21 (51.2%)		32 (78%)	9 (22%)	
	IKDC score (mean \pm SD)	90.9 ± 13.7	84 ± 13.6	0.017	80.3 ± 14.4	89.4 + 13.3	0.036

Table 2 Subgroup analysis for IKDC score at follow-up according to postoperative regimen

Abbreviations: ACLR, anterior cruciate ligament reconstruction; ACLR-OI, anterior cruciate ligament reconstruction + other interventions; IKDC, International Knee Documentation Committee; ns, nonsignificant; SD, standard deviation; WB, weight bearing.

Independently from the type of surgical treatment, significantly better results were reported in the subgroup without brace and FWB in comparison with bracing and PWB, while patients wearing a brace had worse IKDC score when FWB was not allowed (**~Table 3**).

Discussion

The overall mean IKDC score in this study appeared consistent with literature.^{16,17} In the mid-term follow-up, patients without brace and who had reached FWB 6 weeks after surgery generally exhibited better IKDC scores than patients who wore a knee brace and continued to have PWB 6 weeks after surgery. This finding confirmed the previous literature that showed no advantage of a postoperative treatment with a stabilizing knee brace after ACLR.^{10–15,18} However, when we analyzed separately the patients who had concomitant surgery (ACLR-OI group), no differences were found in IKDC score between the braced and unbraced subjects.

The use of a brace is widespread, because it is considered to have positive effects on joint stability and protects the graft by minimizing the stress forces across the knee. However, disadvantages of bracing have been claimed, including the potential muscle atrophy, loss of knee extension at the removal, decreased patient's perception of maximal performance, increased fatigability during exercise, and additional costs.^{12–15,18} Unfortunately, no evidence exists on effectiveness of bracing in patients with ACLR and concomitant surgery,¹⁸ and this issue needs to be confirmed by further research.

Regarding the timing of full weight-bearing prescription, great differences among orthopaedic surgeons are common, both for isolated ACLR and ACLR combined with meniscal repair/resection or other additional surgery. Two literature reviews concluded that 2 years postoperatively, the outcomes in patients with a lateral meniscus repair, medial meniscus resection, or lateral meniscus resection, were not significantly different from that of patients with an isolated ACLR.¹⁹ Also, the concomitant meniscal injuries had no significant influence on the outcomes up to 8-year followup.²⁰ This point is particularly relevant, since the association of meniscal resection and ACL reconstruction has been demonstrated to increase the rate of osteoarthritis over the long term, at a minimum 10 years after ACL injury.¹⁹ In this study, the type of intervention did not influence IKDC score at a mean follow-up of 3.5 years postoperatively. However, when separately analyzed, patients with ACLR and adjunctive surgery demonstrated no difference in functional outcomes with respect to the weight-bearing timing. The different provisions of surgeons stressed in this study

Table 3 Effect of	postoperative regimen	on IKDC score at follow-up	regardless of	f surgical treatment
-------------------	-----------------------	----------------------------	---------------	----------------------

Subgroups	Bracing	WB	Surgical treatment (N)	IKDC score (mean \pm SD)	<i>p</i> -Value	Post-hoc test (p-Values)							
(N)						Subgroups (pairwise comparisons)							
						1 vs. 4	2 vs. 4	3 vs. 4	1 vs. 2	1 vs. 3	2 vs. 3		
Subgroup 1 (17)	No	Full	ACLR (11) ACLR-OI (6)	90.9 ± 14.8	0.02	0.032	ns	0.03	ns	ns	ns		
Subgroup 2 (3)	No	Progressive	ACLR (2) ACLR-OI (1)	91.0 ± 6.2									
Subgroup 3 (15)	Yes	Full	ACLR (5) ACLR-OI (10)	87.6 ± 11.7									
Subgroup 4 (6)	Yes	Progressive	ACLR (4) ACLR (2)	74.9 ± 14.7									

Abbreviations: ACLR, anterior cruciate ligament reconstruction; ACLR-OI, anterior cruciate ligament reconstruction + other interventions; IKDC, International Knee Documentation Committee; ns, nonsignificant; SD, standard deviation; WB, weight bearing.

well describe the need for further research in the area to gain evidences for optimal and shared postoperative management in ACLR.

Also, rehabilitation programs after ACL reconstruction, with or without additional surgery, lack the standards.^{6,7} In this study, the protocol followed during 2 weeks of supervised rehabilitation was designed according to the evidence obtained from literature. Both open and closed kinetic chain exercises were utilized,^{21,22} emphasizing weight bearing when possible and ROM and progressive muscular strengthening.^{9–11,23} Neuromuscular electrical stimulation (NMES) combined with exercise was employed for quadriceps strength^{24–26} together with proprioceptive and balance exercises, which have been demonstrated to improve joint position sense, muscle strength, and perceived knee joint function.^{27,28}

Although preoperative rehabilitation seems to be effective in improving the outcomes after surgery,^{29,30} this was not carried out in the present cohort of patients. Furthermore, with respect to the duration of the rehabilitation, there is evidence that a postoperative one year rehabilitation, individually tailored based on possible concomitant surgery, graft source, and the patient's functional status, and associated to a preoperative rehabilitation, improves the postoperative outcomes.³¹

This study has some limitations. According to the rules of the National Health System, we were unable to follow patients for a long period in hospital and usually, after the first 2 weeks of postoperative rehabilitation in hospital, patients were advised to continue rehabilitation in the outpatient clinics. This could represent a bias in the study since, due to the retrospective nature of the study, the information on the timing and type of rehabilitation that patients had after the period in our institute and that could have affected the final outcome, was not available. In addition, the small number of patients in subgroups is a relevant limitation of this study. Unfortunately, only 38% of eligible patients for the study accepted to participate. It is possible that since these patients were young and easily returned to their usual daily activities after surgery without any further problem at this long-term follow up, they were not interested in answering to the call.

In conclusion, data in this study evidenced a better midterm functional outcome when patients with isolated ACLR were not immobilized with a knee brace after surgery and had full weight bearing in 6 weeks after surgery. In the case of ACLR with concomitant intervention on menisci or ligaments, data were not clearly in favor of any of the condition explored in terms of bracing and weight bearing. Caution should be exercised in interpreting the results in ACLR-OI group due to the small number of cases, asking for further research in the field.

References

- 1 Lobb R, Tumilty S, Claydon LS. A review of systematic reviews on anterior cruciate ligament reconstruction rehabilitation. Phys Ther Sport 2012;13(04):270–278
- 2 Hernández LM, Micheo WF, Amy E. Rehabilitation update for the anterior cruciate ligament injured patient: current concepts. Bol Asoc Med P R 2006;98(01):62–72

- 3 Bowers AL, Spindler KP, McCarty EC, Arrigain S. Height, weight, and BMI predict intra-articular injuries observed during ACL reconstruction: evaluation of 456 cases from a prospective ACL database. Clin J Sport Med 2005;15(01):9–13
- 4 Mohtadi NG, Chan DS, Dainty KN, Whelan DB. Patellar tendon versus hamstring tendon autograft for anterior cruciate ligament rupture in adults. Cochrane Database Syst Rev 2011;7(09):CD005960
- 5 Adams D, Logerstedt DS, Hunter-Giordano A, Axe MJ, Snyder-Mackler L. Current concepts for anterior cruciate ligament reconstruction: a criterion-based rehabilitation progression. J Orthop Sports Phys Ther 2012;42(07):601–614
- 6 Negus J, Fransen M, Chen JS, Parker DA, March L. Exercise-based interventions for conservatively or surgically treated anterior cruciate ligament injuries in adults. Cochrane Database Syst Rev 2012;(10):CD010128
- 7 Trees AH, Howe TE, Grant M, Gray HG. WITHDRAWN: Exercise for treating anterior cruciate ligament injuries in combination with collateral ligament and meniscal damage of the knee in adults. Cochrane Database Syst Rev 2011;18(05):CD005961
- 8 Makhni EC, Crump EK, Steinhaus ME, et al. Quality and variability of online available physical therapy protocols from academic orthopaedic surgery programs for anterior cruciate ligament reconstruction. Arthroscopy 2016;32(08):1612–1621
- 9 Kruse LM, Gray B, Wright RW. Rehabilitation after anterior cruciate ligament reconstruction: a systematic review. J Bone Joint Surg Am 2012;94(19):1737–1748
- 10 Wright RW, Preston E, Fleming BC, et al. A systematic review of anterior cruciate ligament reconstruction rehabilitation: part I: continuous passive motion, early weight bearing, postoperative bracing, and home-based rehabilitation. J Knee Surg 2008;21(03): 217–224
- 11 Wright RW, Preston E, Fleming BC, et al. A systematic review of anterior cruciate ligament reconstruction rehabilitation: part II: open versus closed kinetic chain exercises, neuromuscular electrical stimulation, accelerated rehabilitation, and miscellaneous topics. J Knee Surg 2008;21(03):225–234
- 12 Bordes P, Laboute E, Bertolotti A, et al. No beneficial effect of bracing after anterior cruciate ligament reconstruction in a cohort of 969 athletes followed in rehabilitation. Ann Phys Rehabil Med 2017;60(04):230–236
- 13 Mayr HO, Stüeken P, Münch EO, et al. Brace or no-brace after ACL graft? Four-year results of a prospective clinical trial. Knee Surg Sports Traumatol Arthrosc 2014;22(05):1156–1162
- 14 Lindström M, Wredmark T, Wretling ML, Henriksson M, Felländer-Tsai L. Post-operative bracing after ACL reconstruction has no effect on knee joint effusion. A prospective, randomized study. Knee 2015;22(06):559–564
- 15 Wright RW, Fetzer GB. Bracing after ACL reconstruction: a systematic review. Clin Orthop Relat Res 2007;455:162–168
- 16 Zaffagnini S, Marcacci M, Lo Presti M, Giordano G, Iacono F, Neri MP. Prospective and randomized evaluation of ACL reconstruction with three techniques: a clinical and radiographic evaluation at 5 years follow-up. Knee Surg Sports Traumatol Arthrosc 2006; 14(11):1060–1069
- 17 Schulz AP, Lange V, Gille J, et al. Anterior cruciate ligament reconstruction using bone plug-free quadriceps tendon autograft: intermediate-term clinical outcome after 24-36 months. Open Access J Sports Med 2013;4(04):243–249
- 18 Lowe WR, Warth RJ, Davis EP, Bailey L. Functional bracing after anterior cruciate ligament reconstruction: a systematic review. J Am Acad Orthop Surg 2017;25(03):239–249
- 19 Filbay SR, Ackerman IN, Russell TG, Macri EM, Crossley KM. Health-related quality of life after anterior cruciate ligament reconstruction: a systematic review. Am J Sports Med 2014; 42(05):1247–1255
- 20 LaPrade CM, Dornan GJ, Granan LP, LaPrade RF, Engebretsen L. Outcomes after anterior cruciate ligament reconstruction using the norwegian knee ligament registry of 4691 patients: how does

meniscal repair or resection affect short-term outcomes? Am J Sports Med 2015;43(07):1591-1597

- 21 Glass R, Waddell J, Hoogenboom B. The effects of open versus closed kinetic chain exercises on patients with acl deficient or reconstructed knees: a systematic review. N Am J Sports Phys Ther 2010;5(02):74–84
- 22 Uçar M, Koca I, Eroglu M, et al. Evaluation of open and closed kinetic chain exercises in rehabilitation following anterior cruciate ligament reconstruction. J Phys Ther Sci 2014;26(12):1875–1878
- 23 Wilk KE, Macrina LC, Cain EL, Dugas JR, Andrews JR. Recent advances in the rehabilitation of anterior cruciate ligament injuries. J Orthop Sports Phys Ther 2012;42(03):153–171
- 24 Kim KM, Croy T, Hertel J, Saliba S. Effects of neuromuscular electrical stimulation after anterior cruciate ligament reconstruction on quadriceps strength, function, and patient-oriented outcomes: a systematic review. J Orthop Sports Phys Ther 2010;40 (07):383–391
- 25 Lepley LK, Wojtys EM, Palmieri-Smith RM. Combination of eccentric exercise and neuromuscular electrical stimulation to improve quadriceps function post-ACL reconstruction. Knee 2015;22(03):270–277
- 26 Imoto AM, Peccin S, Almeida GJ, Saconato H, Atallah ÁN. Effectiveness of electrical stimulation on rehabilitation after ligament

and meniscal injuries: a systematic review. Sao Paulo Med J 2011; 129(06):414–423

- 27 Cooper RL, Taylor NF, Feller JA. A systematic review of the effect of proprioceptive and balance exercises on people with an injured or reconstructed anterior cruciate ligament. Res Sports Med 2005; 13(02):163–178
- 28 Ordahan B, Küçükşen S, Tuncay İ, Salli A, Uurlu H. The effect of proprioception exercises on functional status in patients with anterior cruciate ligament reconstruction. J Back Musculoskeletal Rehabil 2015;28(03):531–537
- 29 Alshewaier S, Yeowell G, Fatoye F. The effectiveness of preoperative exercise physiotherapy rehabilitation on the outcomes of treatment following anterior cruciate ligament injury: a systematic review. Clin Rehabil 2017;31(01):34–44
- 30 Kim DK, Hwang JH, Park WH. Effects of 4 weeks preoperative exercise on knee extensor strength after anterior cruciate ligament reconstruction. J Phys Ther Sci 2015;27(09):2693–2696
- 31 Grindem H, Granan LP, Risberg MA, Engebretsen L, Snyder-Mackler L, Eitzen I. How does a combined preoperative and postoperative rehabilitation programme influence the outcome of ACL reconstruction 2 years after surgery? A comparison between patients in the Delaware-Oslo ACL Cohort and the Norwegian National Knee Ligament Registry. Br J Sports Med 2015;49(06):385–389