


M. W. GRIBBLE, JR, S. GUO, S. L. BUCHWALD\* (MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, USA)


Asymmetric Cu-Catalyzed 1,4-Dearomatization of Pyridines and Pyridazines without Preactivation of the Heterocycle or Nucleophile

*J. Am. Chem. Soc.* **2018**, *140*, 5057–5060.

## Enantioselective Dearomatization of Pyridines with Styrene-Derived Nucleophiles



Selected examples:



**Significance:** Dearomatization of heteroaromatics is a valuable transformation in organic chemistry to create new carbon–carbon bonds. Herein, Buchwald and co-workers report the dearomatization of pyridines with a chiral benzylic copper species generated through an enantioselective copper hydride addition to the corresponding styrene.

**SYNFACTS Contributors:** Mark Lautens, Andrew Whyte  
*Synfacts* 2018, 14(07), 0707 Published online: 18.06.2018  
**DOI:** 10.1055/s-0037-1609533; **Reg-No.:** L06218SF

**Comment:** The intermediate dihydropyridine could be oxidized to regenerate the pyridine or reduced to give the piperidine scaffold. A variety of substituents could be tolerated on either the styrene or pyridine, including halogens and alkyl or alkenyl groups. The alkene could also be a  $\beta$ -substituted styrene containing a heteroatom in the alkyl chain.