

Paper

Substituent-Dependent Chemoselective Synthesis of Highly Functionalized Benzo[*h*]quinolines and 4-Benzylpyrans from 2-Methyl-5-nitro-benzonitrile

Rahul Panwar Shally Ranjay Shaw Amr Elagamy Chandan Shah Ramendra Pratap^{*}

Department of Chemistry, University of Delhi, North campus, Delhi, India-110007, India ramendrapratap@gmail.com rpratap@chemistry.du.ac.in

Dedicated to Prof. Uli Kazmaier on his 59th birthday

Received: 01.10.2018 Accepted after revision: 04.10.2018 Published online: 30.10.2018 DOI: 10.1055/s-0037-1610665; Art ID: so-2018-d0051-op

License terms: cc

Abstract A facile, efficient and atom-economic synthesis of highly substituted benzo[*h*]qninolines was established by reaction of 2-methyl-5-nitrobenzonitrile with suitably functionalized 2*H*-pyran-2-ones under basic conditions. We observed that the presence of a thiomethyl group at the C-4 position of pyran provides 6-aryl-4-(2-cyano-4-nitrobenzyl)-2-oxo-2*H*-pyran-3-carbonitrile exclusively without any trace of benzo[*h*]quinolines. Depending on the nature of the functional group at C-4 of the pyran ring, different products were achieved. To probe the mechanism, we performed control experiments and isolated 3-(1-amino-7-nitro-3-thiophen-2-yl-naphthalen-2-yl)-3-piperidin-1-yl-acrylonitrile, which, on further treatment with base, provided the benzo[*h*]quinolines. The structure of one the products was characterized by singlecrystal X-ray diffraction.

Key words 2*H*-pyran-2-one, 2-methyl-5-nitro-benzonitrile, benzo[*h*]quinoline, 4-benzylpyran, chemoselective

Polycyclic *N*- and *O*-heterocycles such as amsacrine, benzo[*c*]phenanthridines, ellipticines, intoplicine and coralyne exhibit excellent biological properties such as DNA topoisomerase inhibition and anticancer activity.^{1–5} Among the *N*-heterocycles, fused and isolated quinolines are very important and exhibit a wide range of biological properties including anaesthetic,^{6a} antiestrogenic,^{6b-c} antimalarial,⁷ anti-HIV,⁸ anticancer,⁹ antitubercular,¹⁰ and antimicrobial activities.¹¹ They have also been widely used in agrochemical areas¹² and in material chemistry.¹³ Recently, our research group has reported the synthesis and anticancer activity of functionalized benzo[*h*]quinolines.¹⁴ This class of compounds has also been used in the construction of nano and meso structures that exhibit novel electronic and photonic properties.¹⁵



Various synthetic methodologies have been reported for the synthesis of fused and isolated quinolines, from Skraup, Doebner and von Miller syntheses, ^{16–19} through Diels–Alder reactions ²⁰ and Friedlander condensations reactions.²¹ Quinolines have also been synthesized by palladium,²² copper,²³ nickel,²⁴ and zinc²⁵ catalyzed inter- and intramolecular cyclization reactions. The reaction of aryl isothiocyanates, alkynes, and alkyl triflates also provides guinolines and fused guinolines.²⁶ In another approach, partially reduced benzo[h]quinolines were obtained by reaction of arylidenes and 1-tetralone in the presence of ammonium acetate and sodium methoxide.²⁷ Benzo[*h*]quinolines have also been afforded by reaction of 6-methoxy-1-tetralone and methyl propiolate in saturated ammonical methanol.²⁸ Ram and co-workers reported the synthesis of partially reduced benzo[h]quinolines by reaction of 5,6-dihydro-2oxobenzo[*h*]chromenes with formamidine or benzamidine and S-methylisothiourea under basic conditions.²⁹ Recently, our group established a one-pot chemoselective synthesis of benzo[*h*]quinolines by reaction of 2-pyranones and 2-cyanomethylbenzonitrile in DMF in the presence of sodamide. This reaction requires extended reaction times (up to 36-50 h) with conventional heating but 55 minutes under microwave irradiation.³⁰⁻³² In this connection, we wanted to use 2-methyl-5-nitrobenzonitrile as the carbanion source instead of 2-cyanomethylbenzonitrile to study its effect on reactivity with 2-pyranone and to achieve the benzo[h]quinolines without a nitrile group, because regioselective removal of a nitrile group is difficult and requires additional steps.

To initiate our studies, the precursor 6-aryl-4-methylsulfanyl-2-oxo-2*H*-pyran-3-carbonitriles **4** were synthesized by reaction of 2-cyano-3,3-bismethylthioacrylate and a range of aryl/heteroaryl methyl ketones in dimethylsulfoxide under basic conditions at room temperature^{33a-c} and R. Panwar et al.

amination of 6-aryl-4-methylthio-2-oxo-2*H*-pyran-3-carbonitriles with secondary amines in ethanol provided 6-aryl-4-amino-2-oxo-2*H*-pyran-3-carbonitriles **1**.^{34a-e}

To study the optimization, 6-(4-chlorophenyl)-2-oxo-4-(piperidin-1-yl)-2H-pyran-3-carbonitrile (1) and 2-methyl-5-nitrobenzonitrile (2) were selected as a model substrates. Initially, we performed the reaction using triethylamine as base in dimethyl sulfoxide (DMSO) and DMF at 100 °C, but no reaction was observed (Table 1, entries 1 and 2). Similarly, the use of sodamide in DMF or DMSO at either room temperature or higher temperatures led mainly to starting material being recovered (entries 3–6): while use of sodium hydride afforded complex mixtures (entries 7-9) containing starting materials and desired product, along with unidentified decomposition products. When the reaction was performed in DMF using KOH as base at room temperature for 12 h, 45% yield of the desired product was isolated (entry 10) and when the same reaction mixture was heated at 100 °C for 6 h, 93% of the desired benzo[h]quinoline was isolated (entry 11). Use of DMSO instead of DMF, lowered the yield of product (entry 12). Use of potassium tert-butoxide as a base in DMF at 100 °C provided the desired product in 65% yield (entry 13); whereas using cesium carbonate as base was did not afford the desired product (entry 14). By using lithium hydroxide and sodium hydroxide under the same conditions noted in entry 11, a lower yield and complex reaction mixture were observed, respectively (entries 15 and 16). Use of KOH in water gave no reaction and starting materials were recovered (entry 17) and using KOH in DMSO for 6 hours provided only 40% yield of the desired product (entry 18). Thus, we chose the reaction of 2pyranone and 2-methyl-5-nitrobenzonitrile with potassium hydroxide in DMF at 100 °C as the optimal reaction conditions.

The generality of the optimized protocol was then tested by the synthesis of a range of functionalized benzo[h]quinolines (Scheme 1). We have used pyrans functionalized with different secondary amine and aryl groups and the yield was generally not affected significantly by their nature, although the presence of a 2-thienyl group lowered the yield of product.

The structure of one of the products (**3g**) was confirmed by single-crystal X-ray analysis (Figure 1).³⁵ It was found that benzo[h]quinoline ring is completely planar and one unit cell contains eight molecules. It is interesting to note that the C8-C7-C4 angle is 115.29° and not 120°, probably due to steric repulsion by the piperidine ring at C17. Bond angles for C12-C7-C4, C18-C17-N4 are 124.81° and 120.76°, respectively. This repulsion also leads to a higher torsion angle of 59° between the benzo[h]quinoline ring and the pfluorophenyl ring. Paper



277

THIEME



1	NEt ₃	DMSO	100	12	_d
2	NEt ₃	DMF	100	12	_d
3	$NaNH_2$	DMSO	r.t. ^b	12	_d
4	$NaNH_2$	DMSO	70	12	_d
5	$NaNH_2$	DMF	70	12	_ ^d
6	$NaNH_2$	DMF	100	12	_ ^d
7	NaH	DMF	r.t.	12	_ ^e
8	NaH	DMF	100	12	_ ^e
9	NaH	DMSO	100	12	_ ^e
10	КОН	DMF	r.t.	12	45
11	КОН	DMF	100	6	93
12	КОН	DMSO	100	12	65
13	t-BuOK	DMF	100	12	65
14	CS ₂ CO ₃	DMF	100	12	_d
15	LiOH	DMF	100	8	55
16	NaOH	DMF	100	8	_ ^e
17	КОН	H ₂ O	100	12	_d
18	КОН	DMSO	100	6	40

^a All the reactions were performed by stirring 6-(4-chlorophenyl)-4-(piperidin-1-yl)-2*H*-pyran-2-one-3-carbonitrile (**1**; 0.5 mmol), 2-methyl-5-nitrobenzonitrile (**2**; 0.6 mmol), base (1.0 mmol) in solvent (5.0 mL).

^b Room temperature (25–30 °C).

^c Isolated yield reported.

^d Starting materials recovered.

^e Complex mixture formation.

complex mixture formation.





R. Panwar et al.

278

THIEME



Scheme 1 Synthesis of 2-amino-5-aryl-4-sec-amino-benzo[h]quinoline-6-carbonitriles 3. Reagents and conditions: 6-aryl-4-sec-amino-2Hpyran-2-one-3-carbonitriles 1 (0.5 mmol), 2-methyl-5-nitrobenzonitrile 2 (0.6 mmol), KOH (1.0 mmol) in DMF (5.0 mL). Isolated yield reported.

To expand the scope of the reaction, 6-aryl-4-methylsulfanyl-2-oxo-2H-pyran-3-carbonitriles **4** were reacted under similar reaction conditions with 2. The usual workup and purification afforded the 6-aryl-4-(2-cyano-4-nitrobenzyl)-2-oxo-2H-pyran-3-carbonitriles 5a-d chemoselectively (Scheme 2). Interestingly, it was observed that the presence of a thiomethyl group at C-4 diverts the chemoselectivity and provides 6-aryl-4-(2-cyano-4-nitrobenzyl)-2oxo-2H-pyran-3-carbonitriles exclusively, without forma-



Scheme 2 Synthesis of 6-aryl-4-(2-cyano-4-nitrobenzyl)-2-oxo-2Hpyran-3-carbonitrile 5a-d

Mechanistically, the reaction is probably initiated by the generation of the carbanion of 2-methyl-5-nitrobenzonitrile (Scheme 3), which reacts at C-4 or C-6 of the 2-pyranone, depending on the substrate selected. If 6-aryl-4-secamino-2-oxo-2H-pyran-3-carbonitriles are the substrate, the carbanion attacks at C-6 through Michael addition to afford intermediate A. The latter intermediate can then undergo ring opening and protonation to afford two possible intermediate regioisomers, **B** and **D**. If reaction proceeds via *cis* intermediate **B**, then, in the presence of excess base, the benzylic carbanion attacks the aromatic nitrile group to afford intermediate C. Intermediate C undergoes decarboxylation followed by cyclization onto the aliphatic nitrile group with tautomerization to afford the benzo[h]quinolines 3.

Alternatively, if intermediate **D** were to be formed, then benzylic carbanion generated by the excess base could cyclize by involvement of vinylic nitrile to provide intermediate E. Subsequently, the imine group of intermediate E can cyclize onto the aromatic nitrile followed by decarboxylation to afford F, which, on tautomerization, provides functionalized phenanthridines. However, the reaction provides benzo[*h*]quinolines chemoselectively without any trace of phenanthridine.

tion of benzo[*h*]quinolines. The structures of these condensation products were confirmed unambiguously by NMR and IR spectroscopy and by HRMS.



279

Scheme 3 Plausible mechanism for the synthesis of 2-amino-5-aryl-4- *sec*-amino-benzo[*h*]quinoline-6-carbonitriles and 6-aryl-4-(2-cyano-4-nitroben-zyl)-2-oxo-2*H*-pyran-3-carbonitriles

On the other hand, if 6-aryl-4-methylsulfanyl-2-oxo-2*H*-pyran-3-carbonitriles are used as precursors then the carbanion from 2-methyl-5-nitrobenzonitrile attacks at C-4 of the pyran ring via Michael addition with formation of intermediate **G**. Loss of the methylsulfanyl group then affords the 6-aryl-4-(2-cyano-4-nitrobenzyl)-2-oxo-2*H*-pyran-3carbonitriles **5** exclusively.

It is interesting to note that, if 6-aryl-4-methylsulfanyl-2-oxo-2*H*-pyran-3-carbonitriles were used as precursors, the carbanion generated from 2-methyl-5-nitrotoluene attacks at C-4 rather than C-6, possibly due to the greater electrophilicity at C-4 caused by the vacant d-orbitals of sulfur and the better leaving group ability of the SMe group compared with the secondary amine.

To probe the mechanism, we performed the reaction of 2-oxo-4-piperidin-1-yl-6-thiophen-2-yl-2*H*-pyran-3-carbonitrile and 2-methyl-5-nitrotoluene using a lower concentration of base and isolated the 3-(1-amino-7-nitro-3thiophen-2-yl-naphthalen-2-yl)-3-(piperidin-1-yl)-acrylonitrile in 81% yield, which supports the intermediacy of **C** (Scheme 4). Furthermore, we used compound **7** and performed the cyclization under optimized reaction conditions and isolated the benzo[*h*]quinoline. R. Panwar et al.

280

THIEME



Scheme 4 Synthesis of 3-(1-amino-7-nitro-3-(thiophene-2-yl)naph-thalene-2-yl)-3-(piperidin-1-yl)acrylonitrile (**7**) and benzo[*h*]quinoline (**30**)

It is interesting to note that reaction of 2-pyranone with 2-cyanomethylbenzonitrile³⁰ requires 35–50 hours for completion of the reaction, whereas the use of 2-methyl-5-nitrobenzonitrile as a carbanion source provides the ben-zo[h]quinolines in 4–6 hours. We propose that the presence of a nitrile group reduces the rate of reaction, probably by stabilizing the carbanion formed and slowing the cyclization step due to steric hindrance.

In summary, we have developed a simple and efficient approach for the chemoselective synthesis of 5-aryl-4-sec-amino-benzo[*h*]quinoline-6-carbonitriles **3** and 6-aryl-4-(2-cyano-4-nitrobenzyl)-2-oxo-2*H*-pyran-3-carbonitriles **5** by reaction of 2-methyl-5-nitrobenzonitrile with 6-aryl-4-sec-amino-2*H*-pyran-2-one-3-carbonitriles **1** and 6-aryl-4-methylsulfanyl-2*H*-pyran-2-one-3-carbonitriles **4**, respectively, under basic conditions. We also investigated mechanistic aspects of the reaction and found that the reaction can be stopped at the functionalized naphthalene stage. We confirmed the structure of compound **3g** by single-crystal X-ray analysis and found that the aryl group present at position 5 is distorted from the usual sp² angle of 120°.

Commercially available reagents were used directly. NMR spectra were recorded at 400 MHz for ¹H and 100 MHz for ¹³C and chemical shifts (δ) are given as parts per million. Spectra were referenced to the residual ¹H signal of CDCl₃ at δ = 7.24 ppm, ¹³C of CDCl₃ at δ = 77.00 ppm, the residual ¹H of DMSO-*d*₆ at δ = 2.49 ppm and the ¹³C of of DMSO-*d*₆ at δ = 39.50 ppm as the internal standards. Splitting patterns in ¹H NMR data are described as s, singlet; d, doublet; dd, doubled doublet; t, triplet; bs, broad singlet; m, multiplet and coupling constants given in hertz (Hz). HRMS was recorded using an Agilent LCMS with Quadropole time of flight using the ESI mode of ionization. All the required starting materials were synthesized by following reported procedures.^{33a-c}

Synthesis of Benzo[h]quinolines (3a-p); General Procedure

A mixture of 6-aryl-2-oxo-4-*sec*-amino-2*H*-pyran-3-carbonitrile **1** (0.5 mmol), 2-methyl-5-nitrobenzonitrile **2** (0.6 mmol) and KOH (1.0 mmol) in DMF (5.0 mL) was stirred at 100–110 °C for 6 h. After completion of reaction, the reaction mixture was poured onto crushed ice and neutralized with 10% HCl to pH ~6–7. The solid product was filtered, washed with cold water, dried and purified by silica gel column chromatography, eluting with hexane–EtOAc (70:30).

Paper

Synthesis of 6-Aryl-4-(2-cyano-4-nitrobenzyl)-2-oxo-2*H*-pyran-3-carbonitriles (5a-d); General Procedure

A mixture of 6-aryl-4-(methylthio)-2-oxo-2H-pyran-3-carbonitrile **4** (0.5 mmol), 2-methyl-5-nitrobenzonitrile **2** (0.6 mmol) and KOH (1.0 mmol) in DMF (5.0 mL) was stirred at 100–110 °C for 1 h. When the reaction was complete, the mixture was poured onto crushed ice and neutralized with 10% HCl to pH ~6–7. The solid product was filtered, washed with water, dried, and purified by silica gel column chromatography, eluting with hexane–EtOAc (60:40).

3-(1-Amino-7-nitro-3-(thiophen-2-yl)naphthalen-2-yl)-3-(piperidin-1-yl)acrylonitrile (7)

A mixture of 2-oxo-4-(piperidin-1-yl)-6-(thiophen-2-yl)-2H-pyran-3-carbonitrile **10** (0.5 mmol), 2-2-methyl-5-nitrobenzonitrile **1** (0.6 mmol), and KOH (0.6 mmol) in DMF (5.0 mL) was stirred at 100– 110 °C for 2–3 h. After completion of reaction, the reaction mixture was poured onto crushed ice followed by neutralization with 10% HCI to pH ~6–7. The crude material was filtered, washed with water, dried, and purified by silica gel column chromatography eluting with hexane–EtOAc (70:30).

9-Nitro-5-phenyl-4-(piperidin-1-yl)benzo[h]quinolin-2-amine (3a)

Yield: 88% (175 mg); red solid; mp 190-192 °C.

IR (KBr): 3356, 1334 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 0.75–0.85 (m, 2 H), 0.92–1.00 (m, 2 H), 1.21–1.26 (m, 2 H), 2.33–2.39 (m, 2 H, CH₂-N-CH₂), 2.97 (d, *J* = 12 Hz, 2 H, CH₂-N-CH₂), 5.22 (s, 2 H, -NH₂), 6.30 (s, 1 H, ArH), 7.25–7.32 (m, 3 H, ArH), 7.36 (s, 1 H, ArH), 7.41 (d, *J* = 8 Hz, 2 H, ArH), 7.76 (d, *J* = 8 Hz, 1 H, ArH), 8.24 (dd, *J* = 8 Hz, 1 H, ArH), 9.96 (s, 1 H, ArH).

 ^{13}C NMR (100 MHz, CDCl₃): δ = 23.5, 24.3, 52.5, 98.5, 113.1, 121.2, 122.0, 124.5, 127.0, 127.2, 128.3, 129.1, 129.4, 136.3, 141.5, 141.6, 145.2, 148.5, 157.7, 160.2.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₄H₂₃N₄O₂: 399.1816; found: 399.1819.

4-Morpholino-9-nitro-5-phenylbenzo[h]quinolin-2-amine (3b)

Yield: 85% (170 mg); red solid; mp >250 °C.

IR (KBr): 3449, 1334 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 2.52 (t, *J* = 20 Hz, 2 H), 2.66 (t, *J* = 20 Hz, 2 H), 2.92 (d, *J* = 12 Hz, 2 H), 3.49 (d, *J* = 12 Hz, 2 H), 4.91 (s, 2 H, -NH₂), 6.34 (s, 1 H, ArH), 7.40–7.42 (m, 2 H, ArH), 7.45 (s, 1 H, ArH), 7.48–7.50 (m, 3 H, ArH), 7.86 (d, *J* = 8 Hz, 1 H, ArH), 8.34 (dd, *J* = 12 Hz, 1 H, ArH), 10.0–10.0 (m, 1 H, ArH).

 ^{13}C NMR (100 MHz, DMSO- d_6): δ = 51.1, 64.6, 99.2, 111.4, 120.8, 121.0, 123.3, 127.0, 127.1, 128.6, 129.1, 129.3, 136.0, 141.0, 141.7, 144.4, 148.8, 157.8, 159.3.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₃H₂₁N₄O₃: 401.1608; found: 401.1609.

_

281 THIEME

9-Nitro-4-(piperidin-1-yl)-5-(p-tolyl)benzo[h]quinolin-2-amine (3c)

Yield: 86% (180 mg); red solid; mp 150-152 °C.

IR (KBr): 3480, 1337 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 1.18–1.26 (m, 3 H), 1.39–1.40 (m, 3 H), 2.33 (s, 3 H, -CH₃), 2.78–2.90 (m, 3 H, CH₂-N-CH₂), 3.01–3.09 (M, 1 H, CH₂-N-CH₂), 4.84 (s, 2 H, -NH₂), 7.13 (d, *J* = 8 Hz, 2 H, ArH), 7.19 (s, 1 H, ArH), 7.23 (s, 1 H, ArH), 7.29 (d, *J* = 8 Hz, 2 H, ArH), 7.8 (d, *J* = 8 Hz, 1 H, ArH), 8.81 (s, 1 H, ArH).

¹³C NMR (100 MHz, CDCl₃): δ = 21.2, 23.6, 24.3, 52.5, 98.3, 113.2, 121.2, 122.0, 124.2, 127.8, 128.2, 129.2, 129.3, 136.4, 136.8, 138.8, 141.6, 145.1, 148.7, 157.7, 160.1.

HRMS (ESI): m/z [MH]⁺ calculated for C₂₅H₂₅N₄O₂: 418.1972; found: 418.1973.

4-Morpholino-9-nitro-5-(p-tolyl)benzo[h]quinolin-2-amine (3d)

Yield: 82% (170 mg); red solid; mp 240–242 °C.

IR (KBr): 3354, 1332 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 2.44 (s, 3 H, -CH₃), 2.54–2.62 (m, 2 H), 2.65–2.69 (m, 2 H), 2.93 (d, *J* = 8 Hz, 2 H, CH₂-N-CH₂), 3.5 (d, *J* = 8 Hz, 2 H, CH₂-N-CH₂), 4.88 (s, 2 H, -NH₂), 6.32 (s, 1 H, ArH), 7.22 (d, *J* = 8 Hz, 2 H, ArH), 7.37 (d, *J* = 4 Hz, 2 H, ArH), 7.43 (s, 1 H, ArH), 7.84 (d, *J* = 8 Hz, 1 H, ArH), 8.33 (dd, *J* = 8 Hz, 1 H, ArH), 10.01–10.02 (m, 1 H, ArH).

 ^{13}C NMR (100 MHz, CDCl₃): δ = 21.7, 52.1, 70.0, 114.6, 116.0, 121.0, 122.5, 125.2, 127.0, 127.1, 127.4, 127.5, 128.4, 129.7, 130.1, 130.2, 130.8, 130.8, 133.1, 156.4.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₄H₂₃N₄O₃: 415.1765; found: 415.1762.

5-(4-Methoxyphenyl)-9-nitro-4-(piperidin-1-yl)benzo[*h*]quinolin-2-amine (3e)

Yield: 78% (167 mg); red solid; mp 245–257 °C.

IR (KBr): 3401, 1333 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 0.81–0.86 (m, 2 H), 1.20–1.31 (m, 4 H), 2.38–2.43 (m, 2 H, CH₂-N-CH₂), 3.01–3.04 (m, 2 H, CH₂-N-CH₂), 3.85 (s, 3 H, -OMe), 5.27 (s, 2 H, -NH₂), 6.31 (s, 1 H, ArH), 6.9 (d, *J* = 8 Hz, 2 H, ArH), 7.36 (d, *J* = 8 Hz, 2 H, ArH), 7.39 (s, 1 H, ArH), 7.77 (d, *J* = 8 Hz, 1 H, ArH), 8.26 (dd, *J* = 8 Hz, 1 H, ArH), 9.94–9.95 (m, 1 H, ArH).

 ^{13}C NMR (100 MHz, CDCl₃): δ = 24.4, 29.6, 52.4, 55.4, 98.1, 112.6, 113.0, 121.2, 122.0, 124.1, 128.1, 128.3, 130.4, 130.5, 134.2, 136.4, 141.1, 145.0, 157.6, 159.0, 160.1.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₅H₂₅N₄O₃; 429.1921; found: 429.1929.

5-(4-Methoxyphenyl)-4-morpholino-9-nitrobenzo[*h*]quinolin-2-amine (3f)

Yield: 79% (170 mg); red solid; mp >250 °C.

IR (KBr): 3383, 1332 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 2.59–2.69 (m, 2 H), 2.91–2.93 (m, 2 H), 3.49–3.51 (m, 2 H), 3.86 (s, 3 H, -OMe), 4.90 (s, 2 H, -NH₂), 6.30 (s, 1 H, ArH), 6.93 (d, *J* = 8 Hz, 2 H, ArH), 7.37–7.40 (m, 3 H, ArH), 7.81 (d, *J* = 8 Hz, 1 H, ArH), 8.3 (dd, *J* = 8 Hz, 1 H, ArH), 9.97–9.98 (m, 1 H, ArH).

¹³C NMR (100 MHz, DMSO- d_6): δ = 51.3, 55.7, 65.2, 99.4, 112.0, 113.0, 121.1, 121.5, 123.1, 128.1, 129.0, 129.1, 129.7, 131.0, 134.4, 139.2, 141.1, 149.4, 159.2, 159.8.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₄H₂₃N₄O₄, 431.1714; found: 431.1715.

5-(4-Fluorophenyl)-9-nitro-4-(piperidin-1-yl)benzo[h]quinolin-2-amine (3g)

Yield: 81% (168 mg); red solid; mp 200–202 °C.

IR (KBr): 3358, 1334 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 1.19–1.23 (m, 3 H), 1.30–1.34 (m, 3 H), 2.37–2.43 (m, 2 H, CH₂-N-CH₂), 3.01–3.04 (m, 2 H, CH₂-N-CH₂), 4.83 (s, 2 H, -NH₂), 6.33 (s, 1 H, ArH), 7.04–7.09 (m, 2 H, ArH), 7.37 (s, 1 H, ArH), 7.43–7.46 (m, 2 H, ArH), 7.82 (d, *J* = 12 Hz, 1 H), 8.31 (dd, *J* = 8 Hz, 1 H), 7.98 (d, *J* = 4 Hz, 1 H).

¹³C NMR (100 MHz, CDCl₃): δ = 24.6, 29.7, 52.7, 98.8, 114.1 (d, J_{C-F} = 20 Hz), 121.7 (d, J_{C-F} = 80 Hz), 124.5, 128.3, 129.8, 131.0 (d, J_{C-F} = 10 Hz), 136.3, 138.0, 140.4, 145.3, 149.4, 158.0, 160.1.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₄H₂₂FN₄O₂: 417.1721; found: 417.1724.

5-(4-Fluorophenyl)-4-morpholino-9-nitrobenzo[*h*]quinolin-2amine (3h)

Yield: 80% (167 mg); red solid; mp >250 °C.

IR (KBr): 3484, 1332 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 2.53–2.69 (m, 4 H), 2.89 (d, *J* = 8 Hz, 2 H, CH₂-N-CH₂), 3.52 (d, *J* = 12 Hz, 2 H, CH₂-N-CH₂), 4.90 (s, 2 H, -NH₂), 6.32 (s, 1 H, ArH), 7.1 (t, *J* = 12 Hz, 2 H, ArH), 7.38 (s, 1 H, ArH), 7.42–7.46 (m, 2 H, ArH), 7.84 (d, *J* = 8 Hz, 1 H, ArH), 8.31 (dd, *J* = 4 Hz, 1 H), 9.98 (d, *J* = 4 Hz, 1 H).

¹³C NMR (100 MHz, CDCl₃): δ = 51.3, 65.4, 98.5, 112.5, 114.2 (d, J_{C-F} = 30 Hz), 121.6 (d, J_{C-F} = 20 Hz), 125.0, 128.5, 130.8, 130.8, 136.2, 139.6, 145.4, 157.7, 159.0, 161.1, 163.5.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₃H₂₀FN₄O₃: 419.1514; found: 419.1513.

5-(4-Chlorophenyl)-9-nitro-4-(piperidin-1-yl)benzo[*h*]quinolin-2-amine (3i)

Yield: 93% (201 mg); red solid; mp 215-217 °C.

IR (KBr): 3489, 1328 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 0.94–1.02 (m, 1 H), 1.17–1.20 (m, 1 H), 1.26–1.29 (m, 3 H), 1.38–1.41 (m, 1 H), 2.35 (t, *J* = 24 Hz, 2 H, CH₂-N-CH₂), 2.95 (d, *J* = 12 Hz, 2 H, CH₂-N-CH₂), 4.81 (s, 2 H, -NH₂), 6.28 (s, 1 H, ArH), 7.28–7.30 (m, 3 H, ArH), 7.36 (d, *J* = 8 Hz, 2 H, ArH), 7.76 (d, *J* = 8 Hz, 1 H, ArH), 8.25 (dd, *J* = 8 Hz, 1 H, ArH), 9.92–9.93 (m, 1 H, ArH).

 ^{13}C NMR (100 MHz, CDCl₃): δ = 24.1, 26.2, 53.7, 104.5, 115.6, 115.8, 116.1, 121.2, 128.6, 129.2, 130.7, 130.8, 134.8, 135.2, 138.3, 145.0, 147.0, 158.8, 162.0, 164.3.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₄H₂₂ClN₄O₂: 433.1426; found: 433.1435

5-(4-Chlorophenyl)-4-morpholino-9-nitrobenzo[*h*]quinolin-2-amine (3j)

Yield: 91% (197 mg); red solid; mp 150–152 °C.

IR (KBr): 3353, 1329 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 2.60–2.71 (m, 4 H), 2.90 (d, *J* = 12 Hz, 2 H, CH₂-N-CH₂), 3.55 (d, *J* = 12 Hz, 2 H, CH₂-N-CH₂), 4.97 (s, 2 H, -NH₂), 6.35 (s, 1 H, ArH), 7.38–7.44 (m, 5 H, ArH), 7.85 (d, *J* = 8 Hz, 1 H, ArH), 8.34 (dd, *J* = 8 Hz, 1 H), 9.99 (d, *J* = 4 Hz, 1 H).

		THIEME
SynOpen	R. Panwar et al.	OPEN ACCESS

Paper

¹³C NMR (100 MHz, DMSO- d_6): δ = 55.4, 67.6, 112.6, 116.4, 120.4, 120.7, 121.3, 121.3, 128.3, 130.4, 130.6, 133.0, 137.3, 139.5, 139.6, 142.7, 144.4, 145.5, 160.6.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₃H₂₀ClN₄O₃: 435.1218; found: 435.1219

5-(4-Bromophenyl)-9-nitro-4-(piperidin-1-yl)benzo[*h*]quinolin-2-amine (3k)

Yield: 82% (195 mg); red solid; mp 150-152 °C.

IR (KBr): 3366, 1333 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 1.15–1.27 (m, 4 H), 1.41–1.47 (m, 2 H), 2.85–2.90 (m, 2 H, CH₂-N-CH₂), 3.01–3.08 (m, 2 H, CH₂-N-CH₂), 4.87 (s, 2 H, -NH₂), 7.19–7.20 (m, 3 H, ArH), 7.27 (d, *J* = 12 Hz, 1 H, ArH), 7.46 (d, *J* = 8 Hz, 2 H), 7.80–7.81 (m, 1 H, ArH), 8.18 (dd, *J* = 8 Hz, 1 H), 8.80 (s, 1 H, ArH).

 ^{13}C NMR (100 MHz, CDCl₃): δ = 23.5, 26.0, 66.3, 109.0, 118.8, 119.2, 120.6, 120.7, 121.6, 121.1, 130.0, 130.1, 131.0, 131.1, 136.7, 139.0, 142.0, 143.0, 144.6, 160.0.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₄H₂₂BrN₄O₂: 477.0921; found: 479.0901.

5-(4-Bromophenyl)-4-morpholino-9-nitrobenzo[*h*]quinolin-2-amine (3l)

Yield: 83% (199 mg); red solid; mp 210-212 °C.

IR (KBr): 3351, 1338 cm⁻¹.

¹H NMR (400 MHz, $CDCl_3$): $\delta = 2.58-2.70$ (m, 4 H), 2.85 (d, J = 12 Hz, 2 H, CH_2 -N- CH_2), 3.54 (d, J = 8 Hz, 2 H, CH_2 -N- CH_2), 4.91 (s, 2 H, $-NH_2$), 6.33 (s, 1 H, ArH), 7.36 (t, J = 16 Hz, 3 H, ArH), 7.53 (d, J = 8 Hz, 2 H), 7.86 (d, J = 12 Hz, 1 H, ArH), 8.33 (dd, J = 8 Hz, 1 H), 9.98 (d, J = 4 Hz, 1 H).

 ^{13}C NMR (100 MHz, DMSO- d_6): δ = 51.2, 66.3, 100.1, 111.7, 120.6, 121.0, 121.4, 123.5, 129.3, 129.7, 130.6, 131.9, 136.5, 140.2, 141.5, 145.0, 149.4, 158.0, 160.0.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₃H₂₀BrN₄O₃: 479.0713; found: 481.0689.

5-(2,4-Dichlorophenyl)-9-nitro-4-(piperidin-1-yl)benzo[*h*]quinolin-2-amine (3m)

Yield: 72% (168 mg); red solid; mp 240-242 °C.

IR (KBr): 3493, 13343 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 0.75–0.82 (m 2 H), 1.24–1.26 (m, 2 H), 2.10–2.18 (m, 2 H), 2.51–2.55 (m, 2 H), 2.74 (d, *J* = 12 Hz, 1 H), 3.03 (d, *J* = 12 Hz, 1 H, CH₂-N-CH₂), 4.81 (s, 2 H, -NH₂), 6.37 (s, 1 H, ArH), 7.25–7.30 (m, 2 H, ArH), 7.36–7.37 (m, 2 H, ArH), 7.4 (d, *J* = 8 Hz, 1 H, ArH), 7.78 (d, *J* = 12 Hz, 1 H, ArH), 8.26 (dd, *J* = 12 Hz, 1 H, ArH), 9.96–9.97 (m, 1 H, ArH).

¹³C NMR (100 MHz, DMSO- d_6): δ = 25.0, 29.0, 55.0, 101.2, 114.0, 120.7, 120.8, 121.0, 121.2, 127.4, 127.6, 129.1, 129.3, 132.0, 135.7, 136.8, 140.5, 145.0, 159.5, 160.2.

HRMS (ESI): m/z [MH]⁺ calcd. for $C_{24}H_{21}Cl_2N_4O_2$: 467.1036; found: 467.1034.

5-(2,4-Dichlorophenyl)-4-morpholino-9-nitrobenzo[*h*]quinolin-2-amine (3n)

Yield: 74% (173 mg); red solid; mp 180-182 °C.

IR (KBr): 3356, 1334 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 2.25–2.34 (m, 2 H), 2.47–2.52 (m, 1 H), 2.71 (d, *J* = 12 Hz, 1 H), 2.90–3.01 (m, 2 H), 3.46 (d, *J* = 12 Hz, 1 H), 3.75 (d, *J* = 8 Hz, 1 H), 5.29 (s, 2 H, -NH₂), 6.47 (s, 1 H, ArH), 7.36–7.39 (m, 2 H, ArH), 7.45 (s, 1 H, ArH), 7.47–7.48 (m, 1 H, ArH), 7.86 (d, *J* = 8 Hz, 1 H, ArH), 8.32 (dd, *J* = 8 Hz, 1 H, ArH), 10.02–10.03 (m, 1 H, ArH).

 ^{13}C NMR (100 MHz, CDCl₃): δ = 53.6, 66.0, 100.3, 114.6, 121.5, 122.0, 126.1, 126.6, 128.6, 128.7, 131.2, 133.6, 135.5, 136.0, 136.3, 140.2, 145.8, 158.0, 160.1.

HRMS (ESI): m/z [MH]⁺ calcd. for $C_{23}H_{19}Cl_2N_4O_3$: 469.0829; found: 469.0823.

9-Nitro-4-(piperidin-1-yl)-5-(thiophen-2-yl)benzo[*h*]quinolin-2amine (30)

Yield: 80% (162 mg); red solid; mp 130–132 °C.

IR (KBr): 3354, 1331 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 1.14–1.23 (m, 2 H), 1.37–1.48 (m, 2 H), 2.88–2.93 (m, 2 H), 3.16–3.22 (m, 2 H, ArH), 4.88 (s, 2 H, -NH₂), 7.05–7.07 (m, 1 H, ArH), 7.33–7.36 (m, 3 H, ArH), 7.83 (d, J = 8 Hz, ArH), 8.21 (dd, J = 8 Hz, 1 H, ArH), 8.82 (s, 1 H, ArH).

 ^{13}C NMR (100 MHz, CDCl_3): δ = 23.6, 65.8, 119.0, 119.2, 119.5, 120.6, 120.7, 121.3, 126.6, 127.1, 127.3, 127.4, 130.0, 135.5, 136.7, 140.8, 143.0, 144.6, 159.8.

HRMS (ESI): m/z [MH]⁺ calcd. for $C_{22}H_{21}N_4O_2S$: 405.1380; found: 405.1375.

4-Morpholino-9-nitro-5-(thiophen-2-yl)benzo[*h*]quinolin-2-amine (3p)

Yield: 84% (171 mg); red solid; mp 180–182 °C.

IR (KBr): 3380, 1372cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 2.69–2.75 (m, 2 H), 2.84–2.90 (m, 2 H), 3.00–3.04 (m, 2 H, CH₂-N-CH₂), 3.60–3.64 (m, 2 H, CH₂-N-CH₂), 4.91 (s, 2 H, -NH₂), 6.35 (s, 1 H, ArH), 7.09–7.11 (m, 1 H, ArH), 7.17–7.18 (m, 1 H, ArH), 7.38–7.40 (m, 1 H, ArH), 7.58 (s, 1 H, ArH), 7.85 (d, *J* = 8 Hz, 1 H, ArH), 8.34 (dd, *J* = 8 Hz, 1 H, ArH), 9.98 (d, *J* = 4 Hz, 1 H, ArH).

 ^{13}C NMR (100 MHz, CDCl₃): δ = 51.6, 65.7, 98.6, 113.1, 121.6, 122.0, 125.4, 125.6, 126.4, 127.1, 128.4, 131.0, 133.3, 136.2, 143.8, 145.4, 145.5, 157.7, 159.0.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₁H₁₉N₄O₃S: 407.1172; found: 407.1169.

4-(4-Benzylpiperazin-1-yl)-5-(4-methoxyphenyl)-9-nitro-4a,10bdihydrobenzo[*h*]quinolin-2-amine (3q)

Yield: 75% (195 mg); red solid; mp >250 °C.

IR (KBr): 3356, 1362 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 2.43 (s, 4 H), 3.40 (s, 2 H, -CH₂), 3.78 (s, 4 H), 3.85 (s, 3 H, -OMe), 4.85 (s, 2 H, -NH₂), 6.9 (d, *J* = 8 Hz, 2 H, ArH), 7.27 (s, 5 H, ArH), 7.36 (d, *J* = 8 Hz, 2 H, ArH), 7.82 (d, *J* = 8 Hz, 1 H, ArH), 8.85 (s, 1 H, ArH).

 ^{13}C NMR (100 MHz, DMSO- d_6): δ = 53.0, 55.7, 62.2, 67.0, 113.5, 113.7, 116.6, 120.1, 120.4, 121.4, 122.1, 127.5, 128.7, 129.3, 129.5, 130.0, 130.1, 133.0, 137.3, 143.8, 144.1, 145.3, 159.4, 160.7.

HRMS (ESI): m/z [MH]⁺ calcd. for C₃₁H₃₂N₅O₃: 522.2500; found: 522.2502.

R. Panwar et al.

5-(4-Methoxyphenyl)-4-(4-methylpiperazin-1-yl)-9-nitro-4a,10bdihydrobenzo[*h*]quinolin-2-amine (3r)

Yield: 72% (161 mg); red solid; mp 210-212 °C.

IR (KBr): 3361, 1381 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 1.23 (s, 2 H), 1.67 (s, 3 H), 2.16 (s, 3 H, N-CH₃), 2.98–3.00 (m, 3 H,), 3.83 (s, 3 H, -OMe), 4.88 (s, 2 H, -NH₂), 6.91 (d, *J* = 8 Hz, 2 H, ArH), 7.28 (S, 1 H, ArH), 7.35–7.39 (m, 3 H, ArH), 7.83 (d, *J* = 8 Hz, 1 H, ArH), 8.22 (dd, *J* = 8 Hz, 1 H, ArH), 8.85–8.85 (m, 1 H, ArH).

 ^{13}C NMR (100 MHz, CDCl₃): δ = 45.7, 49.0, 52.2, 67.7, 113.3, 113.4, 114.5, 119.1, 119.2, 120.5, 120.7, 120.8, 129.8, 129.9, 132.3, 137.0, 142.8, 143.0, 144.5, 159.5, 160.4.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₅H₂₈N₅O₃: 446.2187; found: 446.2185.

4-(2-Cyano-4-nitrobenzyl)-2-oxo-6-phenyl-2*H*-pyran-3-carbonitrile (5a)

Yield: 71% (127 mg); red solid; mp 220-222 °C.

IR (KBr): 2225, 1733, 1520 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 4.41 (s, 2 H, -CH₂), 6.83 (s, 1 H, ArH), 7.45–7.49 (m, 3 H, ArH), 7.75–7.79 (m, 2 H, ArH), 7.82 (d, *J* = 8 Hz, 1 H, ArH), 8.46 (dd, *J* = 8 Hz, 1 H, ArH), 8.58 (d, *J* = 4 Hz, 1 H, ArH).

 $^{13}\mathrm{C}$ NMR (100 MHz, DMSO- d_6): δ = 29.0, 98.7, 104.2, 114.1, 116.0, 116.4, 117.3, 128.1, 128.4, 129.3, 129.4, 131.3, 145.8, 145.9, 158.1, 161.8, 164.1.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₀H₂₁N₃O₄: 358.0822; found: 358.0820.

4-(2-Cyano-4-nitrobenzyl)-6-(4-methoxyphenyl)-2-oxo-2Hpyran-3-carbonitrile (5b)

Yield: 75% (134 mg); red solid; mp 205-207 °C.

IR (KBr): 2225, 1729, 1504 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 3.98 (s, 3 H, -OMe), 4.40 (s, 2 H, -CH₂), 6.75 (s, 1 H, ArH), 6.98–7.00 (m, 2 H, ArH), 7.81–7.85 (m, 3 H, ArH), 8.47 (dd, *J* = 4 Hz, 1 H, ArH), 8.59 (d, *J* = 4 Hz, 1 H).

 ^{13}C NMR (100 MHz, DMSO- d_6): δ = 29.0, 55.7, 96.6, 102.7, 113.8, 114.2, 114.9, 115.8, 121.7, 128.4, 128.7, 131.6, 146.1, 146.8, 158.4, 163.0, 163.2, 164.0.

HRMS (ESI): *m*/*z* calcd. for C₂₁H₁₄N₃O₅: 388.0928; found: 388.0927.

4-(2-Cyano-4-nitrobenzyl)-6-(4-fluorophenyl)-2-oxo-2H-pyran-3carbonitrile (5c)

Yield: 74% (139 mg); red solid; mp 225-227 °C.

IR (KBr): 2226, 1730, 1528 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 4.42 (s, 2 H, -CH₂), 7.15 (s, 1 H, ArH), 7.35 (t, J = 16 Hz, 2 H, ArH), 7.75 (d, J = 8 Hz, 1 H, ArH), 7.91–7.95 (m, 2 H, ArH), 8.42 (dd, J = 4 Hz, 1 H, ArH), 8.77 (d, J = 4 Hz, 1 H, ArH).

¹³C NMR (100 MHz, DMSO-*d*₆): δ = 29.0, 98.7, 104.1, 114.1, 115.8, 116.6 (d, J_{C-F} = 30 Hz), 128.2 (d, J_{C-F} = 30 Hz), 129.4 (d, J_{C-F} = 20 Hz), 131.8, 146.0, 145.8, 158.1, 161.8, 158.1, 161.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₀H₁₁FN₃O₄: 376.0728; found: 376.0726.

6-(4-Chlorophenyl)-4-(2-cyano-4-nitrobenzyl)-2-oxo-2H-pyran-3-carbonitrile (5d)

Yield: 74% (145 mg); red solid; mp 215-217 °C.

IR (KBr): 2228, 1729, 1526 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 4.43 (s, 2 H, -CH₂), 6.85 (s, 1 H, ArH), 7.48 (d, *J* = 8 Hz, 3 H, ArH), 7.78 (d, *J* = 8 Hz, 2 H, ArH), 8.48 (dd, *J* = 4 Hz, 1 H, ArH), 8.60 (d, *J* = 4 Hz, 1 H, ArH).

 ^{13}C NMR (100 MHz, DMSO- d_6): δ = 29.5, 99.6, 105.0, 114.3, 116.3, 128.7, 128.8, 129.0, 128.9, 130.0, 132.2, 138.0, 146.4, 147.3, 158.5, 162.0, 164.6.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₀H₁₁ClN₃O₄: 392.0433; found: 392.0438.

3-(1-Amino-7-nitro-3-(thiophen-2-yl)naphthalen-2-yl)-3-(piperidin-1-yl)acrylonitrile (7)

Yield: 80% (162 mg); red solid; mp 170-172 °C.

IR (KBr): 3356, 2214, 1334 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): δ = 1.14–1.23 (m, 2 H), 1.37–1.48 (m, 2 H), 2.88–2.93 (m, 2 H, CH₂-N-CH₂), 3.16–3.22 (m, 2 H, CH₂-N-CH₂), 4.33 (s, 1 H, =CH-CN), 4.88 (s, 2 H, -NH₂), 7.05–7.07 (m, 1 H, ArH), 7.33–7.36 (m, 3 H, ArH), 7.48 (s, 1 H, ArH), 7.83 (d, J = 8 Hz, 1 H, ArH), 8.21 (dd, J = 8 Hz, 1 H, ArH), 8.82 (s, 1 H, ArH).

 ^{13}C NMR (100 MHz, CDCl₃): δ = 23.6, 29.6, 65.8, 114.5, 119.0, 119.2, 120.6, 120.7, 121.3, 126.6, 127.1, 127.3, 127.4, 130.0, 135.5, 136.7, 140.8, 143.0, 144.6, 159.8.

HRMS (ESI): m/z [MH]⁺ calcd. for C₂₂H₂₁N₄O₂S: 405.1380; found: 405.1379.

Funding Information

RP thanks CSIR, New Delhi (No. 02(0286)/16/EMR-II) for research funding. RP and S thank the University Grants Commission (UGC, New Delhi) and RS thanks CSIR, New Delhi for Senior Research Fellow-ships. The authors thank the University of Delhi for providing research funding and USIC, Delhi University for providing the instrumentation facility

Supporting Information

Supporting information for this article is available online at https://doi.org/10.1055/s-0037-1610665.

References

- Tewey, K. M.; Chen, G. L.; Nelson, E. M.; Liu, L. F. J. Biol. Chem. 1984, 259, 9182.
- (2) Pommier, Y.; Covey, J. M.; Kerrigan, D.; Markovits, J.; Pharm, R. Nucleic Acids Res. **1987**, *15*, 6713.
- (3) Larsen, A. K.; Grondard, L.; Couprie, J.; Desoize, B.; Comoe, L.; Jardillier, J. C.; Riou, J. F. Biochem. Pharmacol. 1993, 46, 1403.
- (4) Makhey, D.; Gatto, B.; Yu, C.; Liu, A.; Liu, L. F.; LaVoie, E. J. Med. Chem. Res. 1996, 56, 2795.
- (5) Nabiev, I.; Chourpa, I.; Riou, J. F.; Nguyen, C. H.; Lavelle, F.; Manfait, M. Biochemistry **1994**, 33, 9013.
- (6) (a) Zishiri, V. K.; Joshi, M. C.; Hunter, R.; Chibale, K.; Smith, P. J.; Summers, R. L.; Martin, R. E.; Egan, T. J. J. Med. Chem. 2011, 54, 6959. (b) Bellot, F.; Coslédéric, F.; Vendier, L.; Brocard, J.; Meunier, B.; Robert, A. J. Med. Chem. 2010, 53, 4103. (c) Klingenstein, R.; Melnyk, P.; Leliveld, S. R.; Ryckebusch, A.; Korth, C. J. Med. Chem. 2006, 49, 5300.
- (7) Michael, J. P. Nat. Prod. Rep. 2008, 25, 166.

		THIEME	
SynOpen	R. Panwar et al.	OPEN ACCESS	Pap

- (8) Chen, Y. W.; Chen, Y. L.; Tseng, C. H.; Liang, C. C.; Yang, C. N.; Yao, Y. C.; Lu, P. J.; Tzeng, C. C. J. Med. Chem. 2011, 54, 4446.
- (9) Lilienkampf, A.; Mao, J.; Wan, B.; Wang, Y.; Franzblau, S. G.; Kozikowski, A. P. J. Med. Chem. 2009, 52, 2109.
- (10) (a) Nandhakumar, R.; Suresh, T.; Jude, A. L. C.; Kannan, V. R.; Mohan, P. S. *Eur. J. Med. Chem.* **2007**, *42*, 1128. (b) Kategaonkar, A. H.; Shinde, P. V.; Kategaonhar, A. H.; Pasale, S. K.; Shingate, B. B.; Shingare, M. S. *Eur. J. Med. Chem.* **2010**, *45*, 3142. (c) Sabatini, S.; Gosetto, F.; Manfroni, G.; Tabarrini, O.; Kaatz, G. W.; Patel, D.; Cecchetti, V. J. Med. Chem. **2011**, *54*, 5722.
- (11) Mahata, P. K.; Venkatesh, C.; Kumar, U. K. S.; Ila, H.; Junjappa, H. *J. Org. Chem.* **2003**, 68, 3966.
- (12) Piechowska, J.; Gryko, D. T. J. Org. Chem. 2011, 76, 10220.
- (13) Younes, L.; Vincent, H.; Chandrasekaran, Y.; Desce, M. B.; Acher, F. C.; Nicolas, P. *J. Org. Chem.* **2012**, 77, 8294.
- (14) Yadav, D. K.; Rai, R.; Kumar, N.; Singh, S.; Misra, S.; Sharma, P.; Shaw, P.; Pérez-Sanchez, H.; Mancera, R. L.; Choi, H. E.; Kim, M. H.; Pratap, R. Scientific Reports **2017**, 6, 38128.
- (15) (a) Jenekhe, S. A.; Lu, L.; Alam, M. M. Macromolecules 2001, 34, 7315. (b) Aggarwal, A. K.; Jenekhe, S. A. Macromolecules 1991, 24, 6806. (c) Zhang, X.; Shetty, A. S.; Jenekhe, S. A. Macromolecules 1999, 32, 7422.
- (16) Skraup, Z. H. Monatsh. Chem. 1881, 2, 139.
- (17) Doebner, O.; Miller, W. V. Ber. Dtsch. Chem. Ges. 1884, 17, 1698.
- (18) Cohn, B. E.; Gustavson, R. G. J. Am. Chem. Soc. 1928, 50, 2709.
- (19) Warren, F. L. J. Chem. Soc. 1936, 1366.
- (20) Peters, O.; Friedrichsen, W. Tetrahedron Lett. 1995, 36, 8581.
- (21) Riesgo, E. C.; Jin, X.; Thummel, R. P. J. Org. Chem. 1996, 61, 3017.
- (22) Luo, Y.; Pan, X.; Wu, J. Org. Lett. 2011, 13, 1150.
- (23) (a) Sakai, N.; Tamura, K.; Shimamura, K.; Ikeda, R.; Konakahara, T. Org. Lett. 2012, 14, 836. (b) Huang, H.; Jiang, H.; Chen, K.; Liu, H. J. Org. Chem. 2009, 74, 5476.
- (24) Korivi, R. P.; Cheng, C. H. J. Org. Chem. 2006, 71, 7079.
- (25) Jiang, B.; Si, Y. G. J. Org. Chem. 2002, 67, 9449.
- (26) Zhao, P.; Yan, X.; Yin, H.; Xi, C. Org. Lett. 2014, 16, 1120.
- (27) Al-Mutairi, T. M.; Al-Hazimi, H. M.; El-Baih, F. M. J. Saudi. Chem. Soc. **2009**, 13, 199.
- (28) Janin, Y. L.; Bisagni, E.; Carrez, D. J. Heterocycl. Chem. **1993**, 30, 1129.
- (29) Pratap, R.; Ram, V. J. Tetrahedron Lett. 2007, 48, 2755.

- (30) Singh, S.; Yadav, p.; Sahu, S. N.; Sharone, A.; Kumar, B.; Ram, V. J. *Synlett* **2014**, *25*, 2599.
- (31) Singh, S.; Panwar, R.; Althagafi; Sharma, V.; Chaudhary, S. Tetrahedron Lett. **2015**, 56, 5203.
- (32) Singh, S.; Panwar, R.; Yadav, P.; Althagafi, I.; Sahu, S. N.; Pratap, R. *RSC Adv.* **2015**, *5*, 18335.
- (33) (a) Tominaga, Y.; Ushirogochi, A.; Matsuda, Y. J. Heterocycl. Chem. 1987, 24, 1557. (b) Gompper, R.; Kutter, E.; Töpfl, W. Justus Liebigs Ann. Chem. 1962, 659, 90. (c) Ram, V. J.; Verma, M. Indian J. Chem. 1990, 29B, 624.
- (34) (a) Pratap, R.; Kumar, B.; Ram, V. J. *Tetrahedron* 2006, 34, 8158.
 (b) Pratap, R.; Kumar, B.; Ram, V. J. *Tetrahedron* 2006, 62, 8158.
 (c) Farhanullah, F.; Agarwal, N.; Goel, A.; Ram, V. J. *J. Org. Chem.* 2003, 68, 2983. (d) Tominaga, Y.; Ushirogouchi, A.; Matsuda, Y.; Kobayashi, G. *Chem. Pharm. Bull.* 1984, 32, 3384. (e) Ram, V. J.; Verma, M.; Hussaini, F. A.; Shoeb, A. *J. Chem. Res., Synop.* 1991, 98.
- (35) Crystal data for $C_{24}H_{23}FN_4O_2$: A red crystal (0.220 × 0.200 × 0.180 mm³) was mounted on a capillary tube for indexing and intensity data collection at 298 K with an Oxford Xcalibur Sapphire3 CCD single-crystal diffractometer (Mo K α radiation, λ = 0.71073 Å). Routine Lorentz and polarization corrections were applied, and an absorption correction was performed using the ABSCALE 3 program [CrysAlis Pro software system, Version 171.34; Oxford Diffraction Ltd., Oxford U. K., 2011]. Patterson methods were used to locate the heavy atoms (SHELXS-86), and the remaining atoms were located from successive Fourier maps (SHELXL-97). All the non-hydrogen atoms were refined anisotropically. All hydrogen atoms were calculated after each cycle of refinement using a Riding model, with C-H = 0.93 Å + $U_{iso}(H) = 1.2U_{eq}(C)$ for aromatic H atoms, and with C–H = 0.97 Å + $U_{iso}(H)$ = 1.2 $U_{eq}(C)$ for methylene H atoms. Crystal data: CCDC No 1862527 for $C_{24}H_{23}FN_4O_2$; M_r = 416.45, crystal system: monoclinic; space group C 2/c; *a* = 16.9960(14) Å, *b* = 8.9323(6) Å, $c = 26.3854(19 \text{ Å}, \alpha = 90, \beta = 101.583, \gamma = 90; V = 3924.1(5);$ Z = 8, ρ_{calcd} = 1.410 Mg/m³ g cm⁻³; μ = 0.099 mm⁻¹; F(000) = 1744.0, R1 = 0.0558 (3860) and wR2 = 01420 (5075) for $I > 2\sigma(I)$ and 226 parameters, R1= 0.0558 (3860) and wR2 = 01420 (5075), gof = 0.904 for all data.