

Conclusion: Co-existing HOCM and aneurysm warrant vigilance to avoid factors increasing LVOT obstruction and to maintain the cerebral perfusion pressure.

Keywords: intracranial aneurysms, hypertrophic obstructive cardiomyopathy, LVOT

References

1. Sahoo RK, Dash SK, Raut PS, Badole UR, Upasani CB. Perioperative anesthetic management of patients with hypertrophic cardiomyopathy for noncardiac surgery: a case series. *Ann Card Anaesth* 2010;13(3):253–256
2. Kundra S, Mahendru V, Gupta V, Choudhary AK. Principles of neuroanesthesia in aneurysmal subarachnoid hemorrhage. *J Anaesthesiol Clin Pharmacol* 2014;30(3):328–337

A056 Postoperative Sialadenitis: A Rare Complication of Park Bench Positioning for Neurosurgery

Dona Saha,¹ Rudrashish Halder,¹ Sashi Srivastava,¹ Devendra Gupta,¹ Anil Agarwal¹

¹Department of Anaesthesiology, S.G.P.G.I.M.S., Lucknow, Uttar Pradesh, India

Introduction: Neurosurgical procedures require specific and often unique positions to maximize anatomical exposure. Prolonged surgeries demand maintenance of these positions for long durations. Retrosigmoid–suboccipital craniotomy for cerebellopontine angle tumor is one such technique which requires placing the patient in Park Bench position. Various complications are attributed to this position.¹ A rare and serious complication is postoperative submandibular sialadenitis along with diffuse soft tissue swelling of the neck occurring following extreme degrees of head positioning.² This can cause severe respiratory distress postoperatively.

Methodology/Description: We report a case report of a 42-year-old, ASA-I, male patient diagnosed with right vestibular schwannoma, posted for retrosigmoid–suboccipital craniotomy and excision in a right Park Bench position. The surgery lasted for 8 hours and patient was extubated at the end of surgery uneventfully. After 4 hours, he started developing acute left sided hemi-facial swelling which was diagnosed as acute submandibular sialadenitis with diffuse soft tissue swelling of the same sided oropharyngeal mucosa. The patient consequently developed respiratory distress and stridor, requiring reintubation. The swelling gradually diminished and he could be extubated on the second postoperative day. Complete resolution took approximately 5 days with conservative treatment. Multitude of etiological factors, attributed to this occurrence, and the various remedial measures have been discussed.

Conclusion: Acute postoperative sialadenitis is a rare complication after Park Bench positioning in neurosurgery. Although it has a good prognosis with conservative management, utmost vigilance is required for possible development of respiratory compromise. Meticulous attention to

preoperative oral hygiene and intraoperative positioning probably can prevent its development.

Keywords: sialadenitis, patient positioning, neurosurgery, craniotomy

References

1. Shimizu S, Sato K, Mabuchi I, et al. Brachial plexopathy due to massive swelling of the neck associated with craniotomy in the park bench position. *Surg Neurol* 2009;71(4):504–508; discussion 508–509
2. Lahkar D, Kedia R. Acute post operative unilateral submandibular gland swelling following acoustic schwannoma excision under general anesthesia. *J Neurol Stroke* 2016;5(3):00175

A057 Anesthesia Management of Clival Chordoma with Cervical Body Erosion and Brain Stem Compression

Pratima Kothare¹

¹Department of Anaesthesiology, Bombay Hospital, Mumbai, Maharashtra, India

Introduction: Chordomas are slow growing tumors arising from vestigial notochordal cells. Upto 25% occur at base of skull arising from clivus involving the cerebrovertebral junction (CVJ)

Methodology/Description: A 28-year-old woman, operated case of clival chordoma, presented 2 years later with tingling in both upper and lower limbs, imbalance while walking, and occasional headaches. On examination, assessment of airway was normal, X-ray of cervical spine showed erosion of C2 body, and MRI of brain showed erosion of clivus, extending in the posterior fossa with severe compression of brain stem. Surgical plan was in two stages. Posterior occipital cervical fusion followed by endoscopic endonasal skull base (EENSBS) approach for the excision of chordoma. Anesthesia plan: Awake fiberoptic intubation, with the management of anesthesia suitable for endoscopic skull base surgeries. She was further operated for cerebrospinal fluid (CSF) leak repair and tracheostomy in view of long-standing ventilation. She was discharged with no neurologic sequelae after 21 days.

Conclusion: Plan intubation extubation strategy. Anticipate difficult intubation. Use cervical collar and awake fiberoptic intubation for minimal disturbance at C V junction. Understand the location of tumor, its proximity to vessels and brain stem and be prepared for delayed recovery and ventilator dependency. EENSBS demands lax brain strategies and optimal mean blood pressures. Cater anesthesia for lower cranial nerves neuromonitoring.

Keywords: clival chordoma with cervical body erosion, brain stem compression, occipitocervical fixation, endoscopic endonasal skull base surgery

References

1. Stippler M, Gardner PA, Snyderman CH, Carrau RL, Prevedello DM, Kassam AB. Endoscopic endonasal approach for clival chordomas. *Neurosurgery* 2009;64(2):268–277; discussion 277–278