

Cardiac Arrest and Death in Neurosurgery: An Analysis of Perioperative Anesthetic Adverse Events in Thailand

Phuping Akavipat¹ Pimwan Sookplung¹ Thanatporn Boonsombat¹ Manee Raksakietisak²
Cataleya Tongrong³ Surunchana Lerdsirisopon⁴ Varinee Lekprasert⁵ Yodying Punjasawasd Wong⁶
Vichai Ittichaikulon⁵

¹Department of Anesthesiology, Prasat Neurological Institute, Bangkok, Thailand

²Department of Anesthesiology, Siriraj Hospital, Mahidol University, Bangkok, Thailand

³Department of Anesthesiology, Srinakarind Hospital, Khon Kaen University, Khon Kaen, Thailand

⁴Department of Anesthesiology, King Chulalongkorn Memorial Hospital, Chulalongkorn University, Bangkok, Thailand

⁵Department of Anesthesiology, Ramathibodi Hospital, Mahidol University, Bangkok, Thailand

⁶Department of Anesthesiology, Maharaj Nakorn Chiangmai Hospital, Chiangmai University, Chiangmai, Thailand

Address for correspondence Phuping Akavipat, MD, FRCAT, PhD, Department of Anesthesiology, Prasat Neurological Institute, 312 Rajvithee Road, Bangkok 10400, Thailand (e-mail: ppakvp@hotmail.com).

J Neuroanaesthesiol Crit Care 2018;5:87–93.

Abstract

Background Complexities of pathological causes in cardiac arrest and death in neurosurgery require individualized management. To decrease the incidents, this study was performed to describe characteristics and factors reducing adverse outcomes together with potential corrective strategies of perioperative cardiac arrest and death in neurosurgical patients.

Methods An observational study was performed in 22 hospitals. Peer-reviewed consensus was formed using database from Perioperative Anesthetic Adverse Events in Thailand (PAAAd Thai) study. The data contain demography, anesthetic, surgical details, opinions on contributing factors, and factors that minimized incident as well as suggested corrective strategies.

Results From 2,000 incidents, 64 (3%) cardiac arrest events were reported with a 50% chance of return of spontaneous circulation. The most common cardiac rhythm documented was asystole. Essentially, electrocardiography was the most frequent early detector. Surgical-related factors (81.3%), mostly intraoperative bleeding, played a major role in potential causes of arrest. In addition, anesthesia, patient conditions, and system-related factors were found in 65.6%, 57.8%, and 8.3% of the incidents, respectively. Moreover, the severity of the patients (73.3%) was considered to be the most common anesthesia-related risk factor. The prevention of an incident included rule enforcement for patient safety, mandatory knowledge, and also anesthesia personnel's skills development.

Conclusion The optimum fluid assessment and resuscitation, cerebral protection protocols, clinical awareness, and quality assurance together with human resource management are all essential to eliminate the catastrophic cardiac arrest and death in neurosurgery.

Keywords

- complication
- risk
- cardiopulmonary resuscitation
- neuroanesthesia
- national audit

received
April 30, 2018
accepted
May 24, 2018
published online
June 26, 2018

DOI <https://doi.org/10.1055/s-0038-1666884>.
ISSN 2348-0548.

Copyright ©2018 Indian Society of Neuroanaesthesiology and Critical Care

License terms



Introduction

Although a cardiac arrest or a death in neurosurgery does not occur frequently, an impact of an occurrence is severe enough to alter an anesthetic practice and to affect the judgment of a practitioner. The altered practice and judgment are proceeded to avoid a cardiac arrest that leads to severe brain dysfunction from global-ischemic brain injury.¹ According to a survey in 2005, Thai Anesthesia Incidents Study (THAI Study), which was the first national statistical data of mortality and morbidity related to anesthesia ($n = 163,403$), reported an overall perioperative cardiopulmonary arrest within 24 hours of 30.8:10,000 and death of 28.3:10,000.² In addition, an incident report in 2008 showed 14% cardiopulmonary arrest from 2,537 incidents.³ Approximately a decade later, after risk management strategies initiated by the Royal College of Anesthesiologists of Thailand were deployed, the incidence decreased significantly to 15.6:10,000 for the perioperative cardiac arrest and 13.3:10,000 for perioperative death.⁴ Even though the incidence has reduced significantly, continuously available literature monitoring and updating warrant prevention of perioperative cardiac arrest strategy or recommendation is still essential.

Every cardiac arrest in neurosurgical patients is complex and unique. Neurosurgical patients have certain factors that influence cardiopulmonary resuscitation (CPR) especially on mechanisms of cardiac arrest, for example intracranial pathology, surgical procedure, and position of the patient.^{5,6} The individual management based on anesthetic knowledge to enhance safety and improve the outcomes should be integrated properly, for example patient monitoring, dealing with unexpected events, personal nontechnical skill and human resource allocation, etc. Therefore, with a collaboration of the Royal College of Anesthesiologists of Thailand, this study aimed to investigate incidences, characteristics, related factors that minimize adverse outcomes, and suggestive strategies for cardiac arrest and death in neurosurgery in 2015.

Materials and Methods

This study was approved by the Prasat Neurological Institutional Ethics Committee on February 20, 2017 (Number 014/2560; Chairman: Suchart Hanchaipiboonkul) and registered to the Thai Clinical Trials Registry (TCTR) with the identification number of TCTR20170503001.

With the collaboration of 22 hospitals resided in Thailand, a prospective and multicenter observational study of potential causes and corrective strategies of perioperative cardiac arrest and death in neurosurgical patients was conducted. Standard incident record forms were anonymously filled out by performing anesthesiologists or related anesthesia personnel as a part of Perioperative Anesthetic Adverse Events in Thailand (PAAAd Thai) study from January 1 to December 31, 2015. Perioperative cardiac arrest and death data were collected and reviewed. According to the definition, perioperative cardiac arrest and death were defined as cardiopulmonary arrest or death occurred in the hospital, which practically needs CPR, not including a temporary beating stop

from surgical stimuli during intraoperative period and the first 24 hours postoperatively. The records contained demographic data, anesthetic, surgical details, opinions on factors influencing an unfavorable event, and factors minimizing incident as well as suggested corrective strategies.

Thereafter, the standard record forms were thoroughly reviewed by three experienced anesthesiologists and clinical epidemiologist to create a consensus on root-cause analysis. The analysis was performed to identify the most likely etiology of the events categorized into characteristics of patients, methods of surgery, anesthetic and system factors, proper initial management, and also developments of suitable preventive strategies.

In the next session, a descriptive statistical analysis is presented. The continuous data are represented in means \pm standard deviation (SD) whereas categorical data are shown in numbers and percentage.

Results

From 2,000 records, there were 60 (3.0%) patients with 64 events of cardiac arrest during perioperative neurosurgery under anesthesia services. The average age was 49.65 ± 17.71 years with the operation time of 135.86 ± 115.46 minutes. The majority of patients (41; 68.3%) had cardiac arrest in the medical service-directed hospital whereas the rest (19; 31.7%) were in the academic-directed hospitals. Most of the patients arrested in supine position (59; 98.3%). Balanced anesthetic technique with inhalation agent either sevoflurane (33; 84.6%), isoflurane (5; 12.8%), or desflurane (1; 2.6%) was used in 39 (65.0%) patients. Moreover, total intravenous anesthesia was given in 21 (35.0%) patients with propofol (8; 38.1%), narcotics (6; 28.6%), thiopental (4; 19.1%), midazolam (2; 9.5%), and sole muscle relaxant (1; 4.8%). Nitrous oxide was added in seven (17.9%) cases of general anesthesia with inhalation agents. The demographic of the cardiac arrest patients is demonstrated in ►Table 1.

There were 24 (40.0%) incidents that could have been prevented. The prevention can include (1) rule enforcement in three (12.5%) cases, that is, prohibiting practices of supervised anesthetic personnel in patients with ASA (American Society of Anesthesiologists) physical status $> III$ and limiting the number of supervised cases to one anesthesiologist at a time; (2) supplementary knowledge in 18 (75.0%) cases, that is, knowledge of brain protection strategy, pathology of severe brain injury, massive blood transfusion management, and advanced neuromonitoring; and (3) improved administrative skills in 13 (54.2%) cases, that is, awareness of potential massive bleeding and anesthesiologists' nontechnical skills (ANTS). Among these preventions, only seven (11.7%) events were related to the World Health Organization (WHO) surgical safety checklist.

According to the CPR procedures, all the clinical treatments were acceptable. Forty (66.7%) medical staffs who detected these critical events were anesthetic personnel, 18 (28.1%) were registered nurses, and 6 (9.4%) were surgeons. The defibrillation was done in six (9.4%) patients who had fulfilled indication. From 64 arrests, the "do not attempt to resuscitate" (DNAR)

Table 1 Demographics and anesthetic details of the cardiac arrest patients undergoing neurosurgery

	Number	Percent
Sex		
Male	24	40.0
Female	36	60.0
ASA physical status		
I–II	7	11.7
III–V	53	88.3
Diagnosis		
Cerebral tumor	7	11.7
Cerebral aneurysm	2	3.3
Traumatic brain injury	48	80.0
Hydrocephalus	2	3.3
Spinal fracture	1	1.7
Operation		
Craniotomy with tumor removal	7	11.6
Craniotomy with aneurysm clipping	2	3.3
Craniotomy with remove blood clot	19	31.7
Craniectomy with remove blood clot	19	31.7
Ventriculostomy	9	15.0
Shunt	3	5.0
Spinal surgery	1	1.7
Emergency situation	48	80.0
Nonofficial hours	42	70.0
Anesthetic techniques		
Balanced technique with inhalation anesthesia	39	65.0
Balanced technique with total intravenous anesthesia	21	35.0
Monitoring		
Arterial blood pressure	20	33.3
Central venous pressure	12	20.0
End-tidal gas tension	29	48.3
End-tidal carbon dioxide tension	57	95.0
Anesthesia performers		
Certified anesthesiologists	35	58.3
Nurse anesthetists under supervision	17	28.3
Anesthetic residents under supervision	8	13.3

was preordered in the intensive care unit (ICU) for 11 (36.7%) arrests, in the surgical ward for 18 (60.0%) arrests, and in the operating theater for 1 (3.3%) arrest. Nevertheless, after excluding the events of cardiopulmonary arrest with DNAR order, the return of spontaneous circulation (ROSC) appeared in 17 (50.0%) patients, that is, in the operating theater for 11 (64.7%) patients, in the ICU for 5 (29.4%) patients, and in surgical ward for 1 (5.9%) patient. Adrenaline was used in all occurrences. Dopamine, nor-adrenaline, atropine, sodium bicarbonate, and calcium were given 11 (32.4%), 3 (8.8%), 7 (20.6%), 13 (38.2%), and 6 (17.6%) arrests, respectively. The mean \pm SD duration of the procedure was 18.5 ± 32.4 minutes, and particularly in the ROSC case it

was 5 ± 6.8 minutes. The characteristics of CPR procedures in total are shown in ►Table 2.

The Glasgow outcome scale at a hospital discharge shows score of 1 (nonsurvivor) in 55 (91.7%) patients, 2 (severe disability) in 1 (1.7%) patient, and 3–5 (moderate disability to no disability) in 4 (6.7%) patients. Three deaths (5.8%) were declared in the operating theater, 19 (36.5%) in the ICU, and 30 (57.7%) in the surgical ward.

The majority of cardiac arrest events (52; 81.3%) related to a surgery, the other possible causes were due to anesthesia (42; 65.6%), patient-related (37; 57.8%), and system-related (30; 46.9%), as demonstrated in ►Table 3.

Table 2 The characteristics of cardiopulmonary resuscitation procedures

	Number	Percent
Period		
Intraoperative periods	15	23.4
Postoperative periods	49	76.6
Primary method of recognition		
Clinical recognition prior to monitor	9	14.0
Electrocardiography	30	46.9
Noninvasive blood pressure monitor	18	28.1
Arterial blood pressure monitor	4	6.3
Plethysmography	3	4.7
Initial cardiac rhythms		
Asystole	23	35.9
Pulseless electrical activity	7	10.9
Bradycardia	3	4.7
Ventricular fibrillation	3	4.7
Others	1	1.6
Unknown	27	42.2
Witness arrest	64	100.0
“Do not attempt to resuscitate” order	30	46.9

Table 3 The possible causes of cardiac arrest in neurosurgical patients

	Number	Percent
Surgical related		
Intraoperative bleeding	21	32.8
Cerebral edema	17	26.6
Postoperative bleeding	12	18.8
Skull fracture	2	3.1
Anesthesia related		
Cardiovascular instability/hypotension	34	53.1
Airway problems ^a	4	6.3
Pulmonary problems ^b	4	6.3
Patient related		
Severe brain damage	21	32.8
Metabolic derangement	9	14.1
Lethal arrhythmia	4	6.3
Miscellaneous	3	4.7
Systemic related		
Incoordination	4	6.3
Intensive care unit utilization pitfall	26	40.6

^aAirway problems: airway injury, obstruction.

^bPulmonary problems: aspirated pneumonia, pneumonia, pulmonary embolism, traumatic lung injury.

Intraoperative cardiac arrests were intraoperative bleeding and led to severe hypotension (14: 93.3%), tight brain (14: 93.3%), severe brain damage (12: 80%), severe metabolic acidosis (4: 26.7%), lethal arrhythmia (1: 6.7%), and other (2: 13.3%). The emergency cross-matching for blood products

was documented during 5 (8.3%) surgical procedures. Additionally, one (1.7%) particular cause of death was referred to pulmonary embolism.

► **Table 4** presents contributing factors, factors minimizing the incidences, and suggestive corrective strategies.

Table 4 Contributing factors, factors minimizing the events, and suggestive corrective strategies for the cardiac arrest in neurosurgical patients

	Number	Percent
Contributing factors		
Noncompliance to WHO surgical safety checklist	3	5.0
Inappropriate decision making	25	41.7
Lack of knowledge	25	41.7
Inexperience	24	40.0
Emergency situation	48	80.0
Shortage of manpower	5	8.3
Improper patient evaluation and preparation	33	55.0
Lack of monitoring equipment	16	26.7
Mismatched blood transfusion	4	6.7
Communication mishaps	17	28.3
Improper postoperative care unit	29	48.3
Patient severity	44	73.3
Factor minimizing the event		
Comply to checklist and guideline	21	35.0
Previous experienced	41	68.3
Capable assistant	37	61.7
Clinical awareness	46	76.7
Effective consultation system	33	55.0
Competent training	7	11.7
Effective patient monitoring systems	17	28.3
Suggestive corrective strategies		
Develop treatment guideline and recommendations	36	60.0
Appropriated human resource management enterprise	46	76.7
Efficient supervision	37	61.7
Quality control of medical equipment	15	25.0
Quality assurance activity	56	93.3
Intensive care utilization analysis	12	20.0

Abbreviation: WHO, World Health Organization.

The recommendations to decrease intraoperative mortality included mainly promoting quality assurance program (15: 100%); developing clinical practice guidelines (14: 93.3%) such as massive blood transfusion protocol, cerebral protection regimen, and tight brain management paradigm; and intensifying effective medical personnel control (9: 60.0%). Appropriate human resource management is categorized into enhancement of performance and competency (43: 71.7%), educational achievements in clinical anesthesia (20: 33.3%), and personnel allocations (19: 31.7%) whereas enhancement of performance and competency is divided into vigilance to protocols (10: 23.3%) and effective communication (33: 76.7%).

Discussion

This article is a part of the PAA Thai study that is the third national closed-claim study of the eventful complications

perioperatively.^{4,7} The report is based on a voluntary and anonymous basis as a multicenter trial. The incidences were collected at a shared central database but analyzed separately according to individual working group interest. The definitions of each complication were specified in the prerequisite document, and the consensus method was used for the conclusion of the contributing factors, the factors minimizing the event, and the suggestive corrective strategies.

To the best of our knowledge, this is the most recent primary report concerning perioperative cardiac arrest in neurosurgery. Therefore, we do not have a benchmark to compare the occurrences. Even though patient's age, ASA physical status classification, duration of operation, hospital type, emergency situation, diagnosis, and activities during nonofficial hour seem to be common causes of cardiac arrests in general, ASA physical status III–V, emergency condition, and absence or no monitoring of capnometer can also be considered risk factors for perioperative death in intracranial

surgery. However, the anesthetic technique and anesthetic agents are not mentioned in Brovman et al,⁸ Zajac,⁹ Watters et al,¹⁰ Zhou,¹¹ and Akavipat.¹²

Among in-hospital cardiac arrest patients, pulseless electrical activity was the most frequent cause of death from the entire mechanism of fatal arrhythmias.¹³ Though this study result differed, cardiac standstill was observed as the first-rank cause particularly in neurosurgical patients. Pathology of autonomic nervous system in higher brain center accompanying with surgical stimulations is discussed.¹⁴ A critical issue is how to detect this lethal phenomenon as early as possible. Either clinical recognition or information from monitoring equipment would be the best for an incident warning system.¹⁵ The preceding electrocardiographic changes in either the rhythms or the morphology might alert the medical personnel before it converts to a cardiac arrest.¹³ However, how well are the observers aware? This question has no answer yet. The supervision system, presence of a board-certified anesthesiologist, continuing medical education, and training in the aspect of ANTS are proposed in Brovman et al,⁸ Chang,¹⁶ and Lee et al.¹⁷

In this report, the ROSC and the survived patients are higher than the other, reflected fast recognition and abrupt chest compression without any doubts in initial rhythm.¹⁸ Besides the initial cardiac rhythm, the amount of blood loss, the CPR duration and the episodes of hypotension are also demonstrated as the factors related to the ROSC.^{19,20}

The causes of death are principally severe brain damage and cardiovascular and hemodynamics mishaps. Nonetheless, a cerebral protection scheme is not acknowledged documentarily to decrease mortality as well as an intraoperative volume assessment and an effective fluid resuscitation.²¹ Most of the threats that have been discussed are delayed responsiveness. The intraoperative fluid resuscitation including the intravascular volume assessment should be performed continually to minimize this complication as well as the initiation of contingency diagram to deal with intraoperative brain swelling. The institutional massive blood transfusion policy should be declared and followed restrictively emphasizing the criteria to activate this emergency protocol at the appropriated point of time.^{22,23,24,25} Another interesting issue is referred to the ICU utilization. Approximately, one-third of the critically ill patients who have no DNAR order are shifted to the surgical ward instead of an ICU because of an error in judgment, an administrative mistake, or miscommunication. The way of monitoring, critical approaching, and therapeutic facilities might differ. Hence, deaths can be potentially prevented at this level.²⁶

Surprisingly, WHO surgical safety checklist seems to be less useful for the factors minimizing cardiac arrest in neurosurgery although it is found favorable in general surgeries.²⁷ Characteristics of neurosurgical patients and operative procedures are unique.²⁸ Clinical responsiveness following a treatment is extremely dynamic in these fragile patients. The possibility of decreasing cardiac arrest events does not depend predominantly on systematic problem recognition, risk management, or communication

skills that can be revealed by the WHO surgical safety checklist. However, the authors would still recommend accomplishing the WHO surgical safety checklist routinely to prevent unpredictability and other complications.

Structure of anesthesiology department, especially logistics and personnel, was remarked as a cofactor in perioperative mortality.¹⁷ Clinical awareness, reasonable judgment, working experience, treatment guideline/recommendation, and knowledge integration are identified as suggestive corrective strategies based on the current study. The scenarios to manage this dilemma were listed elsewhere as debriefing after simulation-based nontechnical skill training, operational awareness, and medical professionalism.^{29,30}

The limitation of this study is the methodology itself. Because the retrospective data are collected on voluntary and anonymous basis, some reports are probably questionable. The quality control of data management and the rigorous action of individual site managers would be beneficial. In conclusion, perioperative cardiopulmonary arrests during anesthesia in neurosurgery were found 3% from the overall 2,000 patients who had noteworthy adverse events. Pitfalls in intraoperative fluid assessment and resuscitation as well as cerebral protection strategies were remarkable in potentially preventable incidents. Encouragement of clinical awareness, quality assurance, and appropriate human resource management were the effective means for the risk reduction.

Funding Support

Royal College of Anesthesiologists of Thailand, Prasat Neurological Institute, Faculty of Medicine of Chiang Mai University, Chulalongkorn University, Khon Kaen University, Mahidol University (Siriraj Hospital and Ramathibodi Hospital), Prince of Songkla University, Health System Research Institute, and National Research Council of Thailand.

Conflict of Interest

None.

Acknowledgments

This multicenter study was accomplished by personal sacrifice, inspiration together with the cooperation among Thai anesthesiologists, the Anesthesiology Department directors of the participating sites, and the Thai Society for Neuroanesthesia. The Royal College of Anesthesiologists of Thailand (RCAT) and the PAAad Thai study group wish to express deep gratitude to project advisors Professor Thara Tritrakarn; Professor Somsri Paosawasdi; Associate Professor Khun Wanna Somboonwiboon; and Associate Professor Oranuch Kyokong for their exceptionally encouragement, suggestion, and advices.

Furthermore, the authors would like to extend the appreciation to Professor Somrat Charuluxananan, Associate Professor Yodying Punjasawadwong for their devotion and also for the language revision.

References

- 1 Ellis SJ, Newland MC, Simonson JA, et al. Anesthesia-related cardiac arrest. *Anesthesiology* 2014;120(4):829–838
- 2 Charuluxananan S, Punjasawadwong Y, Suraseranivongse S, et al. The Thai Anesthesia Incidents Study (THAI Study) of anesthetic outcomes: II. Anesthetic profiles and adverse events. *J Med Assoc Thai* 2005;88(Suppl 7):S14–S29
- 3 Charuluxananan S, Suraseranivongse S, Jantorn P, et al. Multicentered study of model of anesthesia related adverse events in Thailand by incident report (The Thai Anesthesia Incidents Monitoring Study): results. *J Med Assoc Thai* 2008;91(7):1011–1019
- 4 Charuluxananan S, Sriraj W, Punjasawadwong Y, et al. Perioperative and anesthetic adverse events in Thailand (PAAD THAI) incident reporting study: anesthetic profiles and outcomes. *Asian Biomed* 2017;11(1):21–32
- 5 McClain CD, Soriano SG. Anesthesia for intracranial surgery in infants and children. *Curr Opin Anaesthesiol* 2014;27(5):465–469
- 6 Chowdhury T, Prabhakar H, Bithal PK, Schaller B, Dash HH. Immediate postoperative complications in transsphenoidal pituitary surgery: A prospective study. *Saudi J Anaesth* 2014;8(3):335–341
- 7 Punjasawadwong Y, Sriraj W, Charuluxananan S, et al. Perioperative and anesthetic adverse events in Thailand (PAAD THAI) incident reporting study: hospital characteristics and methods. *Asian Biomed* 2017;11(1):33–39
- 8 Brovman EY, Gabriel RA, Lekowski RW, Dutton RP, Urman RD. Rate of major anesthetic-related outcomes in the intraoperative and immediate postoperative period after cardiac surgery. *J Cardiothorac Vasc Anesth* 2016;30(2):338–344
- 9 Zajac K, Zajac M. Perioperative mortality. *Przegl Lek* 2005;62(3):173–180
- 10 Watters DA, Babidge WJ, Kiermeier A, McCulloch GA, Maddern GJ. Perioperative mortality rates in Australian public hospitals: the influence of age, gender and urgency. *World J Surg* 2016;40(11):2591–2597
- 11 Zhou Y, Li W, Herath C, et al. Off-hour admission and mortality risk for 28 specific diseases: a systematic review and meta-analysis of 251 cohorts. *J Am Heart Assoc* 2016;5(3):e003102
- 12 Akavipat P, Ittichaikulthol W, Tuchinda L, Sothikarnmanee T, Klanarong S, Pranootnarabhal T. The Thai Anesthesia Incidents (THAI Study) of anesthetic risk factors related to perioperative death and perioperative cardiovascular complications in intracranial surgery. *J Med Assoc Thai* 2007;90(8):1565–1572
- 13 Do DH, Hayase J, Tiecher RD, Bai Y, Hu X, Boyle NG. ECG changes on continuous telemetry preceding in-hospital cardiac arrests. *J Electrocardiol* 2015;48(6):1062–1068
- 14 Biering-Sørensen F, Biering-Sørensen T, Liu N, Malmqvist L, Wecht JM, Krassioukov A. Alterations in cardiac autonomic control in spinal cord injury. *Auton Neurosci* 2018;209:4–18
- 15 Bein B, Seewald S, Gräsner JT. How to avoid catastrophic events on the ward. *Best Pract Res Clin Anaesthesiol* 2016;30(2):237–245
- 16 Chang CH. Medical simulation is needed in anesthesia training to achieve patient's safety. *Korean J Anesthesiol* 2013;64(3):204–211
- 17 Lee JH, Kim EK, Song IK, et al. Critical incidents, including cardiac arrest, associated with pediatric anesthesia at a tertiary teaching children's hospital. *Paediatr Anaesth* 2016;26(4):409–417
- 18 Amnuaypattanapon K, Udomsubpayakul U. Evaluation of related factors and the outcome in cardiac arrest resuscitation at Thammasat Emergency Department. *J Med Assoc Thai* 2010;93(Suppl 7):S26–S34
- 19 Fei Y, Wang L, Zhong T. Factor analysis for the prognosis of perioperative cardiopulmonary resuscitation [in Chinese] *Zhonghua Yi Xue Za Zhi* 2014;94(1):18–21
- 20 Yi HJ, Kim YS, Ko Y, Oh SJ, Kim KM, Oh SH. Factors associated with survival and neurological outcome after cardiopulmonary resuscitation of neurosurgical intensive care unit patients. *Neurosurgery* 2006;59(4):838–845, discussion 845–846
- 21 Germans MR, Boogaarts HD, Macdonald RL. Neuroprotection in critical care neurology. *Semin Neurol* 2016;36(6):642–648
- 22 Cleland S, Corredor C, Ye JJ, Srinivas C, McCluskey SA. Massive haemorrhage in liver transplantation: consequences, prediction and management. *World J Transplant* 2016;6(2):291–305
- 23 Tran A, Matar M, Steyerberg EW, Lampron J, Taljaard M, Vailancourt C. Early identification of patients requiring massive transfusion, embolization, or hemostatic surgery for traumatic hemorrhage: a systematic review protocol. *Syst Rev* 2017;6(1):80
- 24 Wijdicks EF, Sheth KN, Carter BS, et al; American Heart Association Stroke Council. Recommendations for the management of cerebral and cerebellar infarction with swelling: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2014;45(4):1222–1238
- 25 Trembl AB, Gorlin JB, Dutton RP, Scavone BM. Massive transfusion protocols: a survey of Academic Medical Centers in the United States. *Anesth Analg* 2017;124(1):277–281
- 26 Teixeira PG, Inaba K, Hadjizacharia P, et al. Preventable or potentially preventable mortality at a mature trauma center. *J Trauma* 2007;63(6):1338–1346, discussion 1346–1347
- 27 Haugen AS, Sjøteland E, Almeland SK, et al. Effect of the World Health Organization checklist on patient outcomes: a stepped wedge cluster randomized controlled trial. *Ann Surg* 2015;261(5):821–828
- 28 Ibrahim GM, Morgan BR, Macdonald RL. Patient phenotypes associated with outcomes after aneurysmal subarachnoid hemorrhage: a principal component analysis. *Stroke* 2014;45(3):670–676
- 29 Garden AL, Le Fevre DM, Waddington HL, Weller JM. Debriefing after simulation-based non-technical skill training in healthcare: a systematic review of effective practice. *Anaesth Intensive Care* 2015;43(3):300–308
- 30 Matveevskii A, Moore DL, Samuels PJ. Competency and professionalism in medicine. *Clin Teach* 2012;9(2):75–79