

Relationship of Handedness with Second to Fourth Digit (2D: 4D) Ratio and its Role in Sexual Dimorphism in Tibeto-Nepalese and Indo-Nepalese Adult Population of the Dharan Municipality, Sunsari District of Eastern Nepal: An Anthropometric Study

Sandip Shah¹ Sarun Koirala¹ Laxman Khanal¹

¹Department of Human Anatomy, B.P Koirala Institute of Health Sciences, Nepal

J Morphol Sci 2019;36:33–38.

Address for correspondence Dr Sandip Shah, Associate Professor, Department of Human Anatomy, B.P Koirala Institute of Health Sciences, Nepal, Postal code:56700 (e-mail: sandip.shah@bпкиhs.edu; san230dip@yahoo.com).

Abstract

Introduction The importance of handedness lies in its relationship with the lateralization of the brain function, especially of language. The aim of the present study was to provide an authentic database on right and left second to fourth digit (2D:4D) ratios in 2 different ethnic groups of a particular age and gender, and to study its correlation with handedness and sexual dimorphism.

Materials and Methods The present study was conducted among 400 Tibeto-Nepalese (TN) subjects (200 of each gender) and 400 Indo-Nepalese (IN) subjects (200 of each gender) aged ≥ 18 years old. The values for the 2D:4D ratio were calculated for both hands. Hand preference was established according to the Edinburgh Handedness Inventory, and five hand preference determination groups were constituted after the calculation of the laterality score. The results were tabulated and subjected to statistical analysis. A p -value < 0.05 was considered as statistically significant.

Results The mean value of right sided 2D:4D ratio of TN female was higher as compared to IN female; and for left sided 2D:4D ratio, the value was lower in TN female as compared to IN female. When the values of the 2D:4D ratios were assessed by gender, the values were found to be statistically significant ($p < 0.001$), whereas when they were assessed by ethnicity, only the right hand 2D:4D ratio was found to be statistically significant ($p < 0.001$).

Conclusion When the relationships between the laterality score and the 2D:4D ratios were examined, the values were found to be statistically significant in both ethnic groups only on the left side. The 2D:4D ratio shows significant ethnic and population differences in the present study.

Keywords

- ▶ handedness
- ▶ lateralization
- ▶ tibeto-nepalese
- ▶ indo-nepalese

Introduction

The political scientists Joshi and Rose broadly classify the Nepalese population into three major ethnic groups in terms of their origin: Indo-Nepalese (IN), Tibeto-Nepalese (TN), and

Indigenous (IND). The first group, comprising those of IN origin, inhabits the more fertile lower hills, the river valleys, and the Terai plains adjoining the borders of India. The second major group consists of communities of TN origin occupying the higher hills from the west to the east.¹ The distribution of

received
August 17, 2018
accepted
January 9, 2019
published online
February 15, 2019

DOI <https://doi.org/10.1055/s-0039-1678715>.
ISSN 2177-0298.

Copyright © 2019 by Thieme Revinter Publicações Ltda, Rio de Janeiro, Brazil

License terms



hand preference is J-shaped, reflecting the predominant use of the right hand.² The importance of handedness lies in its relationship with the lateralization of the brain function, especially of language.³ The Geschwind-Behan-Galaburda theory states that the influence on early cell loss and prenatal levels of testosterone are related to cerebral lateralization and handedness.⁴ The sexual dimorphism in the 2D:4D ratio is established as early in as the 9th week of fetal life. It is found in children, is little affected by puberty, and appears to be universal in human populations.⁵⁻⁸

The main objectives of the present study were:

- To calculate the 2D:4D ratios of both hands of adult individuals from 2 different Nepalese ethnic groups (TN and IN) of both genders.
- To study the correlation of the 2D:4D ratios with handedness or hand preference.

Materials and Methods

A population-based cross-sectional study was conducted among TN and IN subjects aged ≥ 18 years old of both genders from the Dharan municipality, in the Sunsari District of Eastern Nepal, using the Edinburgh Handedness Inventory⁹ hand preference determination questionnaire and a separate pre-designed questionnaire for recording socio-demographic variables and length of the second and fourth digits of both hands. The sample size was calculated with an allowable error of 20% (95% confidence interval [CI]) by using the following formula:

$$\text{Sample size } (n) = 4pq/L^2$$

In which:

p = population proportion of positive character,

q = 1-p and

L = allowable error.

Hence:

$$\text{Sample size } (n) = 4pq/L^2 = 780 \text{ individuals}$$

Thus, a total of 800 adults (≥ 18 years old) were recruited for the study of both ethnic groups, TN and IN (400 each), including both genders (200 males and 200 females in each ethnic group) by a systematic random sampling technique. The cutoff point for the age of the subjects was the end of January, so all of the participants had to be of a certain age on February 1st.

Inclusion Criteria

- Residence in the Dharan Municipality.

Exclusion Criteria

- Subjects with any apparent physical hand anomalies, inflammation, trauma, deformities and hand surgery
- Subjects with any genetic, psychological, neurological or chronic diseases affecting hand parameters
- Individuals who disagreed to take part in the study

Digit lengths (–Figs. 1, 2) were measured on the ventral surface of the hand from the basal crease of the digit to the tip of the finger using a vernier caliper (Cescorf Equipamentos para Esportes, Porto Alegre, RS, Brazil) with a precision of 0.01




Fig. 1 Measurement of the right 2nd digit length (a= basal crease; b= the tip of the finger and c= vernier caliper (Cescorf Equipamentos para Esportes, Porto Alegre, RS, Brazil))




Fig. 2 Measurement of the right 4th digit length (a= basal crease; b= the tip of the finger and c= vernier caliper (Cescorf Equipamentos para Esportes, Porto Alegre, RS, Brazil))

mm. Then, they were reported on the questionnaire and the 2D:4D ratio was calculated. This technique of measuring digit length has been reported to have a high degree of repeatability.^{10,11} Hand preference or handedness was determined according to the Edinburgh Handedness Inventory, which evaluates the direction and the degree of hand preference.⁹

Preinformed written consents were obtained from each respondent, and we have also assured them that their anonymity would be preserved during and after the present study. Ethical clearance for the performance of the present study was taken from the institutional ethical and review board. Participation in the present study was voluntary and the purpose of the study was explained to the participants prior to the distribution of the questionnaires. Permission from the Office of the Dharan municipality was also obtained.

Questions regarding the hand preferences of the participants in: 1) writing; 2) drawing; 3) throwing balls; 4) using scissors; 5) using toothbrush; 6) knife without fork; 7) using

Table 1 Hand preference were evaluated in 5 groups depending on the value of the Geschwind laterality score

Hand Preference*	Geschwind Score**	
	Minimum value	Maximum value
Right hand		
Strong	+80	+100
Weak	+20	+75
Ambidextrous	-15	+15
Left hand		
Weak	-75	-20
Strong	-100	-80

*Hand preference was determined according to the Edinburgh Inventory, which evaluates the direction and the degree of hand preference (Oldfield Rc,1971).

**The resultant sum of these points were used to determine the Geschwind (laterality) score, an indicator of the direction and of the degree of hand preference.

a spoon 8); using a broom 9); striking matches; and 10) opening boxes were asked to the subjects by providing the questionnaire. Subjects had to put a “ + ” in the column associated with the hand that they was used to carry out each activity. They had to put a “ + + ” in the associated column if their preference for one hand was very strong; and to put a “ + ” in both columns if they used both hand equally. A “ + + ” in the right column was assigned 10 points, a “ + ” in the right column 5 points, a “ + + ” in the left column - 10 points, and, a “ + ” in the left column - 5 points. The resultant sum of these points was used to determine the Geschwind (laterality) score, an indicator of the direction and of the degree of hand preference. The $-100 \leq$ Geschwind score $\leq +100$ (right hand preference decreases and left hand preference is quantified by Geschwind score, value of which ranges from -100 (strong left hand preference) to $+100$ (strong right hand preference). Hand preference was evaluated in 5 groups depending on the value of the Geschwind laterality score, as shown in **Table 1**.

The results were expressed as mean \pm standard deviation (SD). The gender differences in the 2D:4D ratios and in hand preference in each ethnic group were evaluated by an unpaired

Table 3 Interpretation of 2D:4D ratio by gender, hand and ethnic group

Ethnicity	Hand	Female	Male	p-value
TN	Right	1.007 \pm 0.039	1.005 \pm 0.042	<0.001
	Left	0.999 \pm 0.040	1.003 \pm 0.045	<0.001
IN	Right	1.005 \pm 0.024	1.005 \pm 0.044	0.686
	Left	1.001 \pm 0.031	1.003 \pm 0.045	0.04

Abbreviations: 2D:4D, second to fourth digit ration; IN, Indo-Nepalese; TN,Tibeto-Nepalese.

Results are shown as mean \pm standard deviation, with p-value.

t-test. The right-left differences in 2D:4D ratios (2D:4D right hand - 2D:4D left hand) in each gender and ethnic group were evaluated by an unpaired t-test. The differences of laterality in the 2D:4D ratios, taking into account the hand preference, were evaluated by the paired student t-test. A $p < 0.05$ was considered statistically significant. The correlation between the hand preference and the 2D:4D ratios were evaluated by the Pearson correlation coefficient (r) analysis. Statistical analyses were performed using SPSS Statistics for Windows, Version 11.4 (SPSS Inc., Chicago, IL, USA).

Results

In the present study, 82% of the TN males were found to have a strong right hand (SRH) whereas 82.5% of the TN females were found to have a SRH, followed by 10% and 12.5% strong left hand (SLH) in males and females, respectively, whereas in IN males and females, SRH were 86%, followed by 10% and 12.5% SLH in males and females, respectively (**Table 2**).The 2D:4D ratio of the TN subjects in both hands of both genders were found be statistically significant ($p < 0.001$). However, in the IN subjects of both genders, only the left side 2D:4D ratio was found be statistically significant ($p < 0.001$) (**Table 3**).The right-left difference by gender in each group were found be statistically significant ($p < 0.001$) (**Table 4**). The right-left 2D:4D ratios in the 5 groups evaluated by hand preference in the TN population were found to be not statistically significant ($p > 0.05$) (**Table 5**). The right-left 2D:4Ds ratios in SRH and SLH TN males were found be

Table 2 Distribution of subjects by hand preference and gender

Hand preference	TN male n (%)	TN female n (%)	IN Male n (%)	IN female n (%)	Total n (%)
Strong right hand	164 (82)	165 (82.5)	172 (86)	172 (86)	673 (84)
Weak right hand	13 (6.5)	9 (4.5)	8 (4)	2 (1)	32 (4)
Ambidextrous hand	0	1 (0.5)	0	0	1 (0.1)
Weak left hand	3 (1.5)	0	0	1 (0.5)	4 (0.5)
Strong left hand	20 (10)	25 (12.5)	20 (10)	25 (12.5)	90 (11.2)
Total	200 (25)	200 (25)	200 (25)	200 (25)	800 (100)

Abbreviations: IN, Indo-Nepalese; IND, Indigenous; TN,Tibeto-Nepalese. Results are shown as number and percentage of frequency.

Table 4 Right-Left difference by gender in each group

Ethnicity	Male	Female	p-value
TN	-0.0021 ± 0.044	0.0012 ± 0.053	<0.001
IN	0.0021 ± 0.036	0.0012 ± 0.029	<0.001

Abbreviations: 2D:4D, second to fourth digit ratio; IN, Indo-Nepalese; TN, Tibeto-Nepalese.

Results are shown as mean ± standard deviation, with p-value.

Table 5 Right-Left 2D:4D ratio in five groups evaluated by hand preference in Tibeto-Nepalese

Handedness	Right	Left	p-value
SRH	1.0032 ± 0.041	1.0033 ± 0.042	> 0.05
WRH	0.979 ± 0.52	0.997 ± 0.48	> 0.05
AMBH	0.984	0.857	---
WLH	0.998 ± 0.097	0.967 ± 0.41	> 0.05
SLH	1.0049 ± 0.043	1.0047 ± 0.033	> 0.05

Abbreviations: 2D:4D, second to fourth digit ratio; AMBH, ambidextrous; SLH, strong left hand; SRH, strong right hand; WLH, weak left hand; WRH, weak right hand.

Results are shown as mean ± standard deviation, with p-value.

Table 6 Right-Left 2D:4D ratio in 5 groups evaluated by hand preference in Tibeto-Nepalese by gender

Gender	Handedness	Right	Left	p-value
Male	SRH	1.0033 ± 0.0421	1.0056 ± 0.0416	< 0.001
	WRH	0.9914 ± 0.054	1.0014 ± 0.052	0.581
	WLH	0.9882 ± 0.097	0.9677 ± 0.041	0.71
	SLH	1.0121 ± 0.061	1.0132 ± 0.044	<0.001
Female	SRH	1.0032 ± 0.041	1.0013 ± 0.041	0.151
	WRH	0.9624 ± 0.046	0.9920 ± 0.043	0.601
	AMBH	0.9848	0.8571	---
	SLH	0.9992 ± 0.0228	0.9979 ± 0.0214	0.623

Abbreviations: 2D:4D, second to fourth digit ratio; AMBH, ambidextrous; SLH, strong left hand; SRH, strong right hand; WLH, weak left hand; WRH, weak right hand.

Results are shown as mean ± standard deviation, with p-value.

statistically significant ($p < 0.001$) (► **Table 6**). The right-left 2D:4D ratios in SRH and in weak right hand (WRH) of IN subjects were found be statistically significant ($p < 0.001$) (► **Table 7**). The right-left 2D:4D ratios in SRH IN males and in SRH, WRH, and SLH in IN females were found be statistically significant ($p < 0.001$) (► **Table 8**). The right 2D:4D ratio in WRH ($p = 0.007$) and in SLH ($p = 0.048$) groups in males in both ethnic groups were found be statistically significant (► **Table 9**). The right 2D:4D ratio in SRH ($p < 0.001$) and the

Table 7 Right-Left 2D:4D ratio in 5 groups evaluated by hand preference in Indo-Nepalese

Handedness	Right	Left	p-value
SRH	1.0034 ± 0.036	1.0023 ± 0.038	0.04
WRH	0.998 ± 0.023	1.027 ± 0.077	< 0.001
AMBH	0	0	---
WLH	1.109	0.939	---
SLH	0.9976 ± 0.031	0.998 ± 0.022	> 0.05

Abbreviations: 2D:4D, second to fourth digit ratio; AMBH, ambidextrous; SLH, strong left hand; SRH, strong right hand; WLH, weak left hand; WRH, weak right hand.

Results are shown as mean ± standard deviation, with p-value.

Table 8 Right-Left 2D:4D ratio in 5 groups evaluated by hand preference in Indo-Nepalese by gender

Gender	Handedness	Right	Left	p-value
Male	SRH	1.0069 ± 0.046	1.0034 ± 0.047	< 0.001
	WRH	1.0027 ± 0.021	1.0076 ± 0.047	0.875
	SLH	0.9909 ± 0.037	0.9977 ± 0.025	0.287
Female	SRH	0.9998 ± 0.023	1.0011 ± 0.028	< 0.001
	WRH	0.9797 ± 0.028	1.107 ± 0.151	< 0.001
	WLH	1.1093	0.9393	---
	SLH	1.002 ± 0.0255	0.999 ± 0.199	< 0.001

Abbreviations: 2D:4D, second to fourth digit ratio; SLH, strong left hand; SRH, strong right hand; WLH, weak left hand; WRH, weak right hand.

Results are shown as mean ± standard deviation, with p-value.

right 2D:4D ratio in WRH ($p = 0.011$) groups in females in both ethnic groups were found be statistically significant (► **Table 10**). A positive correlation between the Geschwind score and the 2D:4D ratio was found in both ethnic groups, but this correlation was significant only in the right 2D:4D ratio of both ethnic groups (< 0.001 level; 2-tailed) (► **Table 11**).

Discussion

The differences in hand preferences may be linked to the prenatal production of testosterone and estradiol and, in the case of testosterone, to interactions with the homeobox genes *Hoxa* and *Hoxd*, which control the differentiation of the urogenital system and the development of the digits.¹² It is generally agreed in previous studies that 2D: 4D ratios tend to be greater in females (closer to 1.0), and that gender differences tend to be larger for the right hand than for the left. However, in the present study, only IN females had a greater ratio than males.^{13,14} ► **Table 12** shows the comparative evaluation of the 2D:4D ratios between the present study and previous studies.

Table 9 Right-Left 2D:4D ratio in 5 groups evaluated by hand preference in males in both ethnic groups

Handedness	Right/Left	TN	IN	p-value
SRH	Right	1.0033 ± 0.0421	1.0069 ± 0.046	0.831
	Left	1.0056 ± 0.0416	1.0034 ± 0.047	0.821
WRH	Right	0.9914 ± 0.054	1.0027 ± 0.021	0.007
	Left	1.0014 ± 0.052	1.0076 ± 0.047	0.466
WLH	Right	0.9882 ± 0.097
	Left	0.9677 ± 0.041
SLH	Right	1.0121 ± 0.061	0.9909 ± 0.037	0.048
	Left	1.0132 ± 0.044	0.9977 ± 0.025	0.064

Abbreviations: 2D:4D, second to fourth digit ratio; IN, Indo-Nepalese; SLH, strong left hand; SRH, strong right hand; TN, Tibeto-Nepalese; WLH, weak left hand; WRH, weak right hand. Results are shown as mean ± standard deviation, with p-value.

Table 10 Right-Left 2D:4D ratio in 5 groups evaluated by hand preference in females in both ethnic groups

Handedness	Right/Left	TN	IN	p-value
SRH	Right	1.0032 ± 0.041	0.9998 ± 0.023	< 0.001
	Left	1.0013 ± 0.041	1.0011 ± 0.028	0.06
WRH	Right	0.9624 ± 0.046	0.9797 ± 0.028	0.282
	Left	0.9920 ± 0.043	1.107 ± 0.151	0.011
AMBHI	Right	0.9848
	Left	0.8571
WLH	Right	1.1093
	Left	0.9393
SLH	Right	0.9992 ± 0.0228	1.002 ± 0.0255	0.373
	Left	0.9979 ± 0.0214	0.999 ± 0.199	0.972

Abbreviations: 2D:4D, second to fourth digit ratio; IN, Indo-Nepalese; SLH, strong left hand; SRH, strong right hand; TN, Tibeto-Nepalese; WLH, weak left hand; WRH, weak right hand. Results are shown as mean ± standard deviation, with p-value.

Conclusion

In the present study, 82% of TN males were found to be SRH, whereas 82.5% of TN females were found to be SRH, followed by 10% and 12.5% SLH in males and females, respectively. The 2D:4D ratio showed significant ethnic and population differences.

Table 11 Relationship between G- score and 2D:4D ratio

Ethnic group	2D:4D ratio	Coefficient of correlation#
TN	Right	0.573**
	Left	0.093 (NS)
IN	Right	0.348**
	Left	0.084 (NS)

Abbreviations: 2D:4D, second to fourth digit ratio; IN, Indo-Nepalese; NS, Not significant; TN, Tibeto-Nepalese. #Correlation is significant at a level < 0.001 (2-tailed).

Table 12 Comparative evaluation of 2D:4D ratios in males and in females in different studies

Author	Population	n	Gender	2D:4D
Manning et al (1998) ¹⁰	English	400	M	0.98
			F	1.00
William et al (2000) ¹⁵	American	108 146	M	0.96
			F	0.97
Swami et al (2013) ¹⁶	Harayani Brahmins Kashmiris pandits	150 150	M	0.95
			F	0.98
			M	0.99
			F	1.001
Present study	Tibeto-Nepalese Indo-Nepalese	200 200	M	1.004
			F	0.99
			M	1.003
			F	1.004

Abbreviations: 2D:4D, second to fourth digit ratio; F, female; M, male.

Financial Support

The present study received a research grant from the B. P. Koirala Institute of Health Sciences.

Conflicts of Interest

The authors have no conflicts of interest to declare.

Acknowledgments

We would like to thank Dr. Bhawesh Koirala for helping us during the data collection. We would like to acknowledge the B. P. Koirala Institute of Health Sciences, the Office of the Dharan Municipality, and the Institutional review committee (IRC) for their support and for allowing us to perform the present research.

References

- JOSHI & ROSE. "Nepal-Caste and Ethnicity" [Internet]; 1991, Available from <http://countrystudies.us/nepal/31.htm>. Accessed on 7th April 2018.
- Annett M. Handedness and brain asymmetry: The Right Shift Theory. Hove: Psychology Press; 2002
- Beaton AA. The nature and determinants of handedness. In: Hugdahl K, Davidson RJ (Eds.). The asymmetrical brain. Cambridge, MA: MIT Press; 2003:105–158
- Geschwind N, Galaburda A. Cerebral lateralization. Cambridge, MA: MIT Press; 1986
- Malas MA, Dogan S, Evcil EH, Desdicioglu K. Fetal development of the hand, digits and digit ratio (2D:4D). Early Hum Dev 2006;82 (07):469–475

- 6 Manning JT, Stewart A, Bundred PE, Trivers RL. Sex and ethnic differences in 2nd to 4th digit ratio of children. *Early Hum Dev* 2004;80(02):161–168
- 7 Manning JT, Churchill AJ, Peters M. The effects of sex, ethnicity, and sexual orientation on self-measured digit ratio (2D:4D). *Arch Sex Behav* 2007;36(02):223–233
- 8 McIntyre MH, Ellison PT, Lieberman DE, Demerath E, Towne B. The development of sex differences in digital formula from infancy in the Fels Longitudinal Study. *Proc Biol Sci* 2005;272(1571):1473–1479
- 9 Oldfield RC. The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia* 1971;9(01):97–113
- 10 Manning JT, Scutt D, Wilson J, Lewis-Jones DI. The ratio of 2nd to 4th digit length: a predictor of sperm numbers and concentrations of testosterone, luteinizing hormone and oestrogen. *Hum Reprod* 1998;13(11):3000–3004
- 11 Danborn B, Danborn AM. The effect of the season of birth and fluctuating asymmetry on second and fourth digit lengths and digit ratio (2D:4D) in Nigerians. *Eur J Zool Res* 2015;4(01):7–11
- 12 Mortlock DP, Innis JW. Mutation of HOXA13 in hand-foot-genital syndrome. *Nat Genet* 1997;15(02):179–180
- 13 Brown WM, Finn CJ, Cooke B, Breedlove SM. Differences in finger length ratios between self-identified 'butch' and 'femme' lesbians. *Arch Sex Behav* 2002;31:123–127
- 14 McFadden D, Shubel E. Relative lengths of fingers and toes in human males and females. *Horm Behav* 2002;42(04):492–500
- 15 Williams TJ, Pepitone ME, Christensen SE, et al. Finger-length ratios and sexual orientation. *Nature* 2000;404(6777):455–456
- 16 Swami S, Kumar T, Sharma & Kausal S. Effect of hand preference on second to fourth digit ratio and its role in sexual dimorphism: a study in 300 Haryanvi Brahmins and 300 Kashmiri Pandits. *Eur J Anat* 2013;17(04):243–249