


J. A. PRESCHER, D. H. DUBE, C. R. BERTOZZI* (UNIVERSITY OF CALIFORNIA, BERKELEY, USA)


Chemical Remodelling of Cell Surfaces in Living Animals

Nature 2004, 430, 873–877.

Staudinger Ligation in Living Animals



Staudinger ligation:



Significance: Chemical modification of cell-surface glycans in living animals provides a means to study these biopolymers within their native environment. The Staudinger ligation is a bioorthogonal reaction that can be carried out in the whole organism. This reaction is frequently used for the study of lipids, steroids, and cofactors.

Comment: Bertozzi and co-workers demonstrated the Staudinger ligation of an azide (SiaNAz) and a functionalized phosphine (Phos-Flag) in living mice. The azide was delivered to cell-surface glycoconjugates by metabolism of ManNAz to SiaNAz. The ligation in mice tissue was detected using fluorescein isothiocyanate-labelled anti-Flag antibodies followed by flow cytometry analysis.

SYNFACTS Contributors: Dirk Trauner, Andrej Shemet
Synfacts 2019, 15(10), 1191 Published online: 17.09.2019
DOI: 10.1055/s-0039-1690934; Reg-No.: T08519SF

© 2019, Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Chemistry in
Medicine and
Biology

Key words

Staudinger ligation

metabolic
oligosaccharide
engineering

bioorthogonal
reaction

azides

phosphines

Synfact
Classic

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.