

Occlusal Characteristics of Deaf-Mute Individuals in the Turkish Population

Semra Ciger^a

Seden Akan^b

ABSTRACT

Objectives: To classify and determine the occlusal characteristics of deaf-mute individuals and its gender distribution in the Turkish population.

Methods: For this study, 213 deaf-mute individuals (155 boys and 58 girls) were evaluated. The age range was between 10–24 years, and the mean age was 16.37 ± 2.53 years. Measurements were divided into four groups: dental, intraarch, interarch, and, extra data.

Results: Of the participants, 75.0% had a Class I molar relationship, whereas 13.0% and 8.0% had Class II and Class III malocclusions, respectively. 23.9% of individuals had a normal overbite, 38.4% had a deepbite, and 23.4% had an openbite. One or more congenitally missing teeth were found in 6.0% of individuals; 81.0% expressed satisfaction with their esthetics, and 19.0% expressed the contrary.

Conclusions: Different characteristics and malocclusions are present in deaf-mute individuals. (Eur J Dent 2010;4:128-136)

Key words: Deaf mutism; Occlusal characteristics; Overbite; Openbite; Turkish population.

^a Professor, Department of Orthodontics, Faculty of Dentistry, Hacettepe University, Ankara, Turkey.

^b Research Assistant, Department of Orthodontics, Faculty of Dentistry, Hacettepe University, Ankara, Turkey.

■ Corresponding author: Dt. Seden Akan, Hacettepe Universitesi, Dis Hekimligi Fakultesi, Ortodonti Anabilim Dalı, Sıhhiye, 06100 Ankara, Turkey.
Phone: +90 535 5113760
Fax: +90 312 3091138
E-mail: sedenakandt@hotmail.com

INTRODUCTION

Deaf mutism is a major health problem and limits appropriate communication and learning that is necessary for the normal development and maturation of a child.¹ Its etiology is generally due to hearing loss before the age of 2 – 3 years, which hinders the child being able to learn how to speak; congenital deaf mutism is also a possible etiology. Most cases of deaf mutism result from acute infectious diseases such as measles, epidemic

meningitis, encephalitis, typhoid, otitis media, or toxic effects of drugs.²

Deaf-Mute Individuals (DMI) constitute one of the largest groups of disabled individuals. The World Health Organization (WHO)³ estimates that in 2001, 250 million people worldwide had a disabling hearing impairment (moderate or severe hearing impairment in the better ear). In Turkey, it is estimated that 500 children with hearing impairment are born each year;⁴ there were 158,000 deaf children in 1990.⁵ According to the Turkish Ministry of Education,⁶ there are currently 63 schools for the deaf in Turkey that enrolled a total number of 6,268 students for the 2007–2008 school year. The number of deaf-mute subjects who receive special education accounts for only approximately 5.46% of all deaf-mute subjects in Turkey.

Studies^{7,8} report that prelingually deafened subjects establish open articulatory postures with excessive jaw displacement and minimal tongue movement, and do not contract and extend their tongues as do hearing talkers. A change in tongue muscle function can either cause morphologic variations in the normal configuration of the teeth and supporting bone, or it can exacerbate an existing malocclusion.⁹

There are studies^{1-8,10} that have evaluated abnormalities in DMI in Turkey: Egeli et al¹ found that, out of 162 cases of DMI, 7 individuals had jaw abnormalities and 11 had tooth abnormalities. Öztürk et al¹⁰ report 13 facial asymmetries, 3 cleft tongues, 2 cleft lips, 2 maxillary hypoplasias, 1 cleft palate, 1 prognathism, 11 long face, and 1 lobular tongue subject out of 840 deaf primary school children in Turkey. No study that examines occlusal characteristics of DMI in Turkish populations has been reported in the literature.

The aim of this study is to determine occlusal characteristics of deaf-mute individuals in the Turkish population.

MATERIALS AND METHODS

For this longitudinal study, 213 deaf-mute individuals - 155 boys (73.0%) and 58 girls (27.0%) - in the permanent dentition between the ages of 10-24 years who attend special needs schools in the city of Ankara were evaluated. The mean age of the individuals was 16.37±2.53 years; none of the subjects had received orthodontic treatment before the study.

Measurements were recorded using the one recorder (SC) on the form (Figure 1) which was ap-

proved by the Federal Dentaire International (FDI). This form was primarily for study purposes only.¹¹ Measurements were divided into four groups and are shown in Table 1.

Statistical method

Data analysis was performed using the Statistical Package for Social Sciences (SPSS) version 11.5 software (SPSS Inc., Chicago, IL, United States). Pearson Chi-square tests were used to assess the statistical significance of gender differences in the frequency distribution of categorical variables, unless the expected cell size was less than five or ten, when Fisher's Exact test or Continuity Corrected Chi-square tests were used. Whether or not the differences between gender groups in the subsets of number of missing, malformed, impacted, and extracted teeth were statistically significant was evaluated by using the Mann Whitney U test. A binomial test was applied in order to detect the differences in prevalence between male and female groups. A P value less than .05 was considered statistically significant.

RESULTS

The prevalence of male individuals in this study group was found to be statistically significant higher than girls (P<.001).

Interarch measurements

Anteroposterior molar relationship

The molar relationship was evaluated as Class I, Class II, and Class III according to Angle classification. Subjects whose molar relationship could not be determined because of molar extractions and subjects with different molar relationships on the right and left sides were not classified. Table 2 shows that 160 subjects (75.0%) had Class I malocclusions, 27 (13.0%) had Class II, 18 (8.0%) had Class III, and 8 (4.0%) individuals were not classified.

Vertical relationship

In this research, a 2 mm overbite is accepted as normal, negative values are classified as an openbite, and more than 2 mm is classified as a deepbite. According to this classification system, 51 (23.9%) DMI had a normal overbite, 82 (38.4%) had a deepbite, and 50 (23.4%) had an openbite. The distribution of overbite between genders is presented in Table 2; a statistically significant difference was found (P<.05).

According to the localization, openbite is divided into three groups: anterior, posterior, and anterior-posterior. According to this classification, 35 (16.4%) DMI had anterior openbites, 9 (4.2%) had posterior openbites, and 6 (2.8%) had anterior-posterior openbites (Table 2). Based on the severity, openbite was evaluated in 3 groups; slight (1-2 mm) openbite, moderate openbite (2-4 mm), and severe openbite (more than 4 mm).

According to this classification, 60 (28.0%) DMI had slight, 10 (4.6%) had moderate, and 8 (3.7%) had severe openbites (Table 2).

Transverse relationship

According to transverse relationship evaluation, 8 (2.3%) individuals had a posterior crossbite, 6 (1.9%) had a brodiebite (buccal nonocclusion) (Table 2).

Dental measurements

Congenitally missing teeth

In this group of DMI, congenitally missing teeth were found in 13 subjects (6.0%), but none of the subjects had more than 3 congenitally missing teeth (Table 3). Most frequently, the missing teeth

RECORD FORM

School:
No:
Name and Surname:
Birthday and Place:
Sex:

Dental Number System

	RIGHT										LEFT						
UPPER	17	16	15	14	13	12	11	21	22	23	24	25	26	27			
LOWER	47	46	45	44	43	42	41	31	32	33	34	35	36	37			

Dental Measurements

1-Development Anomalies:
 a) Congenitally Missing Teeth:
 b) Supernumerary Teeth:
 c) Malformed Teeth:
 d) Impacted Teeth:
 e) Transposed Teeth:

2- Teeth Extracted Because Of Trauma:

Intraarch Measurements

a) Crowding:
 b) Diastemas:
 c) Normal:
 d) Can not recording:

3- Midline Diastema:

Interarch Measurements

4-Lateral Segments:
 a)Anteroposterior (Molar Relationship)
 b)Vertically (Posterior Openbite)
 c)Transverse (Posterior Crossbite)

5- Anterior Segments:
 a)Anteroposterior (Overjet)
 b)Vertically (Openbite)
 c)Transverse (Crossbite)

Extra Data

6- Oral Hygiene:
 7- Pleasure of Esthetic:
 8-Previous Orthodontic Treatment:

Figure 1. The form was used in the present study.

Table 1. Measurement groups.

Interarch Measurements	Dental Measurements	Intraarch Measurements	Extra Data
Anteroposterior molar relationship	Congenitally missing teeth	Diastemas	Muscle tone
Vertical relationship	Supernumerary teeth	Crowding	Oral hygiene
Transverse relationship	Malformed teeth		Esthetic satisfaction
	Impacted teeth		
	Teeth extracted because of trauma		

were lateral maxillary incisors (n=11; prevalence: 50.0%; Table 4), or mandibular second premolars (n= 6; prevalence: 26.0%; Table 4). There were no missing canines, first premolars, or first and second molars. No statistically significant gender difference was found regarding missing teeth.

Supernumerary teeth

No supernumerary teeth were found in this group of DMI.

Malformed teeth

In this group, 37 malformed teeth were seen in 16 individuals; its prevalence was 7.5%. One male subject had 8 malformed teeth; this was the highest number of malformed teeth per subject in this group. Malformed teeth were found to be 6 times more frequent in boys than in girls (Table 3), but this difference was not statistically significant (P=0.84). The most frequently malformed teeth were the maxillary lateral incisors (n=16; preva-

Table 2. Distributions of occlusal characteristics between genders (*: P<.05, **: P<.01).

		Male	Female	Total	%	P
Malocclusion	Cl I	116	44	160	75	0.878
	Cl II	19	8	27	13	0.945
	Cl III	14	4	18	8	0.785
	Not grouped	6	2	8	4	1.000
	Total	155	58	213	100	
Overbite	normal	32	19	51	23.9	0.035*
	deepbite	67	15	82	38.4	0.028*
	openbite	36	14	50	23.4	0.738
	Total	135	48	183	85.9	
Openbite	anterior openbite	30	5	35	16.4	0.143
	posterior openbite	5	4	9	4.2	0.065
	ant+post openbite	5	1	6	2.8	1.000
	Total	40	10	50	23.4	
Severity of openbite	1-2 mm	51	9	60	28	0.483
	2-4mm	8	2	10	4.6	0.670
	4mm-	6	2	8	3.7	0.614
	Total	65	13	78	36.6	
Transverse relationship	normal	110	38	148	69.4	0.208
	crossbite	5	3	8	2.3	0.449
	brodiebite	3	3	6	1.9	0.346
	Total	118	44	162	76	
Crowding	slight	13	6	19	8.9	1.000
	moderate	6	2	8	3.7	0.695
	severe	10	7	17	18.3	0.645
	Total	29	15	44	20.6	

lence: 44.0%; Table 4), followed by the maxillary and mandibular central incisors and maxillary second molars (n=4; prevalence: 11.0%; Table 4).

Impacted teeth

In DMI, 22 impacted teeth were seen in 15 individuals; its prevalence was 7.0% (Table 3). The most frequently impacted teeth were maxillary canines (n=20; prevalence: 44.0%; Table 4), followed by mandibular second premolars and canines (n=1; prevalence: 4.5%; Table 4). A statistically significant gender difference for impacted teeth was found (P<.05).

Extracted teeth

A total of 107 teeth were missing due to extraction in 52 male and 12 female DMI; the total prevalence was 30.0% (Table 3). The most frequently extracted teeth were mandibular first molars (n=54; prevalence: 51.0%; Table 4), followed by maxillary first molars (n=36; prevalence: 34.0%; Table 4). Table 3 illustrates the fact that male individuals

presented with extracted teeth approximately 4 times more frequently than females, although this difference was not statistically significant (P=0.05). No canine extraction was seen in this group.

Intraarch measurements

Diastemas and crowding

In DMI, 44 subjects (20.6%) had crowding and 40 (18.7%) had diastemas in their dental arches. There were 29 males and only 15 females who had crowding (Table 2). In this research, crowding was divided into three groups: slight (0-2 mm crowding in any dental arch), moderate (2-4 mm), and severe (more than 4 mm). The distribution of subjects according to gender and severity of crowding is presented in Table 2.

Extra data

Oral hygiene

Of all the participants, 49 (30.0%) DMI had good oral hygiene, 48 (29.0%) had moderate hygiene, and 67 (41.0%) had poor oral hygiene (Table 5). There were statistically significant differences be-

Table 3. Number and gender distribution of dental characteristics (*: P<.05, **: P<.01).

	Per subject	Male	Female	Subject Number	%	Teeth Number	%	P
Missing teeth	1	3	2	5	2	5	23	0.349
	2	5	2	7	3	14	63	
	3	0	1	1	0.5	3	14	
	Total	8	5	13	6	22	100	
Malformed teeth	1	4	1	5	2.3	5	13.5	0.840
	2	5	3	8	3.7	16	43.3	
	4	2	-	2	1	8	21.6	
	8	1	-	1	0.5	8	21.6	
	Total	12	2	16	7.5	37	100	
Impacted teeth	1	3	5	8	3.7	8	36	0.021*
	2	4	3	7	3.3	14	64	
	Total	7	8	15	7	22	100	
Extracted teeth	1	26	9	35	16	35	33	0.050
	2	18	1	19	9	38	35	
	3	4	2	6	3	18	17	
	4	4	0	4	4	16	15	
	Total	52	12	64	30	107	100	

Table 4. Distribution of dental characteristics for the jaw.

		Maxillary teeth				Mandibular teeth			
		Right	Left	No	%	Right	Left	No	%
Congenitally missing teeth	Central incisor	1	1	2	9	1	1	2	9
	Lateral incisor	6	5	11	50				
	Canine								
	First premolar								
	Second premolar		1	1	5	4	2	6	27
	First molar								
	Second molar								
Malformed teeth	Central incisor	2	2	4	11	2	2	4	11
	Lateral incisor	8	8	16	44	1	1	2	5
	Canine	1	1	2	5				
	First premolar	1	1	2	5	1		1	3
	Second premolar								
	First molar								
	Second molar	2	2	4	11	1	1	2	5
Impacted teeth	Central incisor								
	Lateral incisor								
	Canine	12	8	20	91	1		1	4.5
	First premolar								
	Second premolar					1		1	4.5
	First molar								
	Second molar								
Extracted teeth	Central incisor	2	1	3	2.5				
	Lateral incisor					1		1	1
	Canine								
	First premolar		1	1	1				
	Second premolar	2	4	6	5	2		2	2
	First molar	20	17	37	34	31	25	56	51
	Second molar						1	1	1

Table 5. Distribution of additional data between genders (*: P<.05, **: P<.01).

		Male	Female	Total	%	P
Oral hygiene	Good	26	23	49	30	P<.001**
	Moderate	41	7	48	29	0.048*
	Poor	56	11	67	41	0.035*
	Total	123	41	164	100	
Pleasure of esthetic	Yes	58	22	80	81	0.386
	No	16	3	19	19	0.386
	Total	74	25	99	100	

tween genders who had good ($P<.001$), moderate ($P<.05$), and poor oral hygiene ($P<.05$).

Esthetic satisfaction

Table 5 shows that 80 (81.0%) DMI were satisfied with their esthetics, whereas 19 (19.0%) were not.

DISCUSSION

Teeth are positioned in equilibrium, in the dental arch between the opposing forces of the lips and tongue. However, additional forces or tonus changes applied to the teeth over extended periods of time will upset the alignment of the dental arch, creating a malocclusion.⁹

There is research proposing that prelingually deafened talkers do not displace the tongue in order to establish vowel steady-state postures and excessively displace the jaw, and that deaf subjects appear to have a less flexible tongue during speech production than do hearing subjects.⁸ Considering this difference, the present study evaluates the occlusal characteristics of deaf-mute individuals in the city of Ankara.

Because the study group contains students from various parts of Anatolia, it is assumed to be a good representation of the Turkish population.

Because of the large and widely scattered sample group, we used malocclusion record forms approved by the FDI for this study. This form was initially used for normal individuals in a study¹¹ and we used it as an opportunity to compare those results with our own.

Interarch and intraarch measurements are evaluated separately because malocclusions have different features¹¹ which do not permit grouping.

The prevalence of different types of malocclusions may show great variability even in a population of the same origin; Thilander et al¹² reports that malocclusion prevalence varies from 39.0% to 93.0%. According to our results, 75.0% of subjects had a Class I molar relationship, 13.0% had a Class II, 8.0% had a Class III malocclusion, and 4.0% could not be grouped because of molar extractions and different molar relationships on the right and left sides.

There are several researches that examine the orthodontic needs of the population and the response of several different orthodontic treatment centers to this need.¹⁰⁻²⁴ Güray et al,¹³ using

the Treatment Priority Index (TPI), reports that 72.3% of 483 primary school students were in need of orthodontic treatment. Ugur et al¹⁴ studied 483 high society primary school students, also using TPI, and found that 40.3% had normal occlusions, 21.5% had slight malocclusions, 25.2% had malocclusions that required treatment, and 5.0% had severe malocclusions. Sari et al¹⁵ evaluated 1602 patients that had accepted treatment and reports that 61.7% of patients had Class I; 25.1% had Class II, division 1; 3.0% had Class II, division 2; and 10.2% had Class III malocclusions. Sayın and Turkkahraman¹⁶ evaluated 1356 untreated patients and report that 64.0% patients had Class I, 24.0% had Class II, and 12.0% had Class III malocclusions. Aytan¹¹ used the same measurement method and evaluated the occlusal features of 1510 high school students. He reports that 56.4% had Class I, 6.9% had Class II, 3.5% had Class III molar relationships, and 33.0% could not be grouped. No significant differences were observed between the prevalence of malocclusions our study and those in previous studies that investigated Turkish, or other populations. In all, Class I malocclusions were found to be the most frequent malocclusion and Class III was the least frequent. Variant results may be explained primarily by sampling technique, local environmental influences (trauma, mouth breathing, or sucking habits), and nutrition.

Previous research¹⁰⁻²⁷ has also evaluated the prevalence of malocclusions in different populations. Jones¹⁷ reports 53.8% Class I malocclusions, 33.2% Class II, and 12.9% Class III malocclusions in a Saudi Arabian population comprised of 132 subjects. Helm¹⁸ investigated Danish population in groups of 1700 subjects and found 14.0% normal occlusion, 58.0% Class I, 18.8% Class II div 1, 2.7% Class II div 2 and 4.0% Class III malocclusion. Lew et al¹⁹ examined 1,050 Chinese subjects and reported that 7.1% had normal occlusions; 58.8% had Class I malocclusions; 18.8% had Class II, div 1; 2.7% had Class II, div 2; and 12.6% had Class III malocclusions.

There is no difference between normal individuals' results and the results of the present study except regarding congenital missing teeth. In the deaf-mute group, 6.0% of individuals had congenitally missing teeth, whereas 2.6% had missing teeth in the normal high school group. This differ-

ence is most likely due to hereditary factors, but no study has been done to investigate the association between deafness and congenitally absent teeth.

In a study²⁰ evaluating the prevalence of congenitally missing teeth in the Turkish population, it was reported that 20.6% of 3,043 people had missing teeth. Similar to our study, it was found that the most frequently missing teeth were maxillary lateral incisors, followed by mandibular second premolars.

According to intraarch measurements of our study, 20.6% individuals had crowding and 18.7% had diastema; in Aytan's study¹¹ these values were 92.0% and 5.6%, respectively. We consider that there are many congenitally missing teeth in the deaf-mute group; it appears that this reflects other intraarch measurements as well. Thus, the incidences of missing teeth and diastemas were high compared to that of crowding in the deaf-mute group. However, this is the first study that has evaluated a correlation between these variables.

In interarch measurements, the rates of deepbites and posterior crossbites were high in DMI, who appear to have a less flexible tongue during speech production than do hearing subjects.⁸ This difference could disrupt tongue function just as abnormal habits would. With changes in the tongue, cheek, and lip muscle functions, the overall effect is a significant narrowing of the maxillary arch, similar to that of open mouth syndrome or mouth breathing subjects.^{28,29} Because the tongue cannot exert the necessary pressure on the incisor segment, deepbites and crossbites may occur.⁹

According to additional data, twice as many DMI subjects exercise good oral hygiene compared to the normal group.¹¹ This shows that on average, DMI are more conscious of their oral hygiene than normal persons. When esthetic satisfaction was evaluated, 81.0% of DMI were satisfied with their appearance, compared to 73.0%¹¹ of normal individuals.

Al-Sarheed et al³⁰ evaluated the parents of 77 visually impaired, 210 hearing impaired, and 494 control children, and finds that 56.7% of the hearing impaired group needed orthodontic treatment, compared to 55.0% in the control group. The authors further report that only 17.9% of parents of hearing impaired children believe that their children are not concerned about their dental ap-

pearance. We believe that the high rate (81.0%) of satisfaction in our study group indicates that the subjects are not overly concerned with their dental appearance.

CONCLUSIONS

In this study of deaf-mute individuals, the most common occlusal relationship was Class I, and the least common was Class III. We also found that the percentage of congenitally missing teeth, deepbites, posterior crossbites, and diastemas are higher in this population, whereas and the incidence of crowding is lower than that of normal individuals. Finally, oral hygiene and esthetic satisfaction rates are very high in deaf-mute individuals.

REFERENCES

1. Egeli E, Çicekci G, Silan F, Ozturk O, Harputluoglu U, Onur A, Egeli A, Yildiz A. Etiology of deafness at the Yeditepe School for the deaf in Istanbul. *Int J Pediatr Otorhinolaryngol* 2003;67:467-471.
2. Janzen VD, Schafer D. Etiology of deafness in Robarts School for the deaf. *J Otolaryngol* 1984;13:47-48.
3. WHO, WHO calls on private sector to provide affordable hearing aids in developing world, WHO/34, 11 July, 2001.
4. Belgin E, Akdas F, Boke B, Çağlar A. The children population with sensorineural hearing loss in Turkey. International meeting in Auditory for the Mediterranean Countries, 1991, p. 181.
5. 1. Specific Education Council: Guide Handbook of specific education institutions, Ministry of Education pres, Ankara, 1991; pp. 12-13.
6. National Education Statistics Formal Education 2007-2008. Ministry of National Education Strategy Development Presidency. A publication of official statistics programme. pp.26. <http://sgb.meb.gov.tr>
7. Obserger MJ. Training effects on vowel production by two profoundly hearing-impaired speakers. *J Speech Hear Res* 1987;30:241-251.
8. Tye-Murray N. The establishment of open articulatory postures by deaf-mute and hearing talkers. *J Speech Hear Res* 1991;34:453-459.
9. Graber TM. The "three M's": Muscles, malformation and malocclusion. *Am J Orthod* 1963;49:418-450.
10. Ozturk O, Silan F, Oghan F, Egeli E, Belli S, Tokmak A, Egeli A, Harputluoglu U, Onder HI, Zafer C. Evaluation of deaf children in a large series in Turkey. *Int J Pediatr Otorhinolaryngol* 2005;69:367-373.
11. Aytan S. Ankara ili ve çevresi lise öğrencilerinde oklüzal özellikler. Hacettepe University, Ankara, Turkey-1978. Thesis.

12. Thilander B, Pena L, Infante C, Parada SS, de Mayorga C. Prevalence of malocclusion and orthodontic treatment need in children and adolescents in Bogota, Colombia. An epidemiological study related to different stages of dental development. *Eur J Orthod* 2001;23:153–167.
13. Güray E, Orhan M, Ertaş E, Doruk C. Konya yöresi ilkokul çocuklarında Treatment Priority Index (TPI) uygulaması (epidemiolojik çalışma). *Türk Ortodonti Dergisi* 1994;7:195–200.
14. Uğur T, Ciğer S, Aksoy A, Telli AE. An epidemiological survey using the Treatment Priority Index(TPI). *Eur J Orthod* 1998;20:189–193.
15. Sari Z, Uysal T, Karaman AI, Basciftci FA, Usumez S, Demir A. Orthodontic malocclusions and evaluation of treatment alternatives: an epidemiologic study. *Turkish J Orthod* 2003;16:119–126.
16. Sayin MO, Turkkahraman H. Malocclusion and crowding in an orthodontically referred Turkish population. *Angle Orthod* 2004;74:635–639.
17. Jones WB. Malocclusion and facial types in a group of Saudi Arabian patients referred for orthodontic treatment: a preliminary study. *Br J Orthod* 1987;14:143–146.
18. Helm S. Malocclusion in Danish children with adolescent dentition: an epidemiologic study. *Am J Orthod* 1968;54:352–366.
19. Lew KK, Foong WC, Loh E. Malocclusion prevalence in an ethnic Chinese population. *Aust Dent J* 1993;38:442–449.
20. Altug-Atac AT, Erdem D. Prevalence and distribution of dental anomalies in orthodontic patients. *Am J Orthod Dentofacial Orthop* 2007;131:510–514.
21. Järvinen S. Indexes for orthodontic treatment need. *Am J Orthod Dentofacial Orthop* 2001;120:237–239.
22. Harrison RL, Davis DW. Dental malocclusion in native children of British Columbia, Canada. *Community Dent Oral Epidemiol* 1996;24:217–221.
23. Ng'ang'a PM, Ohito F, Ogaard B, Valderhaug J. The prevalence of malocclusion in 15 year-old children in Nairobi, Kenya. *Acta Odontol Scand* 1996;54:126–130.
24. Lauc T. Orofacial analysis on the Adriatic islands: an epidemiological study of malocclusions on Hvar Island. *Eur J Orthod* 2003;25:273–278.
25. Silva RG, Kang DS. Prevalence of malocclusion among Latino adolescents. *Am J Orthod Dentofacial Orthop* 2001;119:313–315.
26. Diagne F, Ba I, Ba-Diop K, Yam AA, Ba-Tamba A. Prevalence of malocclusion in Senegal. *Community Dent Oral Epidemiol* 1993;21:325–326.
27. Liepa A, Urtane I, Richmond S, Dunstan F. Orthodontic treatment need in Latvia. *Eur J Orthod* 2003;25:279–284.
28. Aznar T, Galan AF, Marin I, Dominguez A. Dental arch diameters and relationships to oral habits. *Angle Orthod* 2006;76:441–445.
29. Gross AM, Kellum GD, Franz D, Michaz K, Walker M, Foster M, Bishop W. A longitudinal evaluation of open mouth posture and maxillary arch width in children. *Angle Orthod* 1994;64:419–424.
30. Al-Sarheed M, Bedi R, Hunt NP; The views and attitudes of parents of children with a sensory impairment towards orthodontic care. *Eur J Orthod* 2004;26:87–91.