



A. K. MISHRA, G. PARVARI, S. K. SANTRA, A. BAZYLEVICH, O. DORFMAN, J. RAHAMIM, Y. EICHEN*, A. M. SZPILMAN* (TECHNION – ISRAEL INSTITUTE OF TECHNOLOGY, HAIFA AND ARIEL UNIVERSITY, ISRAEL)

Solar and Visible Light Assisted Peptide Coupling

Angew. Chem. Int. Ed. 2021, 60, 12406–12412, DOI: 10.1002/anie.202011510.

Solar Radiation-Assisted Peptide-Bond Formation



Significance: Amide bond formation is extremely important in peptide chemistry. The authors have developed a sunlight-assisted method for generating active coupling reagent in situ from readily available DMAP and bromotrichloromethane to form peptide bonds.

Comment: Various dipeptides can be synthesized by using DMAP and bromotrichloromethane to generate the active coupling reagent. The reactions proceed rapidly, and the yields of the desired peptides are good to excellent.

SYNFACTS Contributors: Hisashi Yamamoto, An Wu
Synfacts 2021, 17(09), 1057 Published online: 18.08.2021
DOI: 10.1055/s-0040-17119953; Reg-No.: H07521SF

© 2021, Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Peptide Chemistry

Key words

peptide bond formation

photocatalysis

solar radiation

bromotrichloromethane

radical reaction

Synfact
of the
Month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.