# Real 3D Visualization of the Circumflex Artery Surrounding the Mitral Annulus

Bálint Fábián<sup>1</sup> András Osadczuk<sup>2</sup> László Bárány<sup>1</sup> Gábor Baksa<sup>1</sup> Gergely Rácz<sup>3</sup> Tamás Ruttkay<sup>1</sup>

Address for correspondence Tamás Ruttkay, MD, PhD, Department

of Anatomy, Histology and Embryology, Laboratory for Applied and

Clinical Anatomy, Semmelweis University, 58 Tűzoltó Street, H-1094

Budapest, Hungary (e-mail: ruttkaytom@gmail.com).

<sup>1</sup> Department of Anatomy, Histology and Embryology, Laboratory for Applied and Clinical Anatomy, Semmelweis University, Budapest, Hungary

<sup>2</sup>Department of Cardiology, Szent Rafael Hospital, Zala County, Zalaegerszeg, Hungary

<sup>3</sup>1st Department of Pathology and Experimental Cancer Research, Semmelweis University, Budapest, Hungary

Thorac Cardiovasc Surg 2022;70:87-92.

Abstract	<b>Background</b> The circumflex coronary artery is located close to the mitral annulus. Consequently, it is not immune to iatrogenic damage during mitral valve procedures. Our objective was to visualize the circumflex artery from a surgeon's point of view, emphasizing its proximity. Furthermore, comparing it to coronary angiograms might support preoperative planning. <b>Methods</b> Ten adult human hearts preserved in 4% formaldehyde solution were investigated (left coronary artery injected with contrast agent). After performing coronary angiographies from LAO (left anterior oblique) 40/cranial 20, RAO (right anterior oblique) 10/cranial 40, and true lateral projections, anatomical preparations were performed. Images were captured throughout the full course of the circumflex coronary artery from multiple angles. Finally, the mean distances were measured in every 5 mm between the investigated artery and the annulus of the mitral valve. <b>Results</b> Three-dimensional model of the circumflex coronary artery and its surround- ings was successfully achieved from a left atrial surgical viewpoint. The main branches were identified on the coronary angiograms. The closest distance was measured under the region of the left auricle ( $2.02 \pm 0.69$ mm; 1–3.1 mm). Afterward, the circumflex artery was observed to make a loop away from the annular region. <b>Conclusion</b> Our observations show correlation with previous anatomical studies and case reports addressing iatrogenic lesions on the circumflex coronary artery. Based on
Keywords	all these, we could determine a "danger zone" on the vessel. The simultaneous
► anatomy	evaluation of the anatomical preparations and the angiograms might improve the
► mitral valve surgery	acknowledgment of this vulnerable region serving to avoid any damage to the coronary
► coronary artery	artery.

# Introduction

Due to its proximity to the mitral annulus, the circumflex coronary artery is susceptible to injury during mitral valve procedures. In functional and degenerative mitral regurgitation-which are the most frequent mitral valve diseases-the

received April 15, 2020 accepted after revision May 29, 2020 published online October 21, 2020

preferred choice of treatment is mitral valve repair or mitral valve replacement.<sup>1–3</sup> The surgical method during both operations consists of suturing around the mitral annulus to implant either the annuloplasty ring in mitral valve repair or the artificial valve in case of mitral valve replacement. The

© 2020. Thieme. All rights reserved. Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

DOI https://doi.org/ 10.1055/s-0040-1715181. ISSN 0171-6425.

iatrogenic damage on the circumflex coronary artery during these procedures has been reported by some authors to occur in 0.5%,<sup>4</sup> while others have revealed the prevalence in 1.8%.<sup>5</sup> There are new commonly used percutaneous approaches as well, such as implantation of the Cardioband device. During these interventional treatments, between 12 and 17 anchors are deployed in a C-shaped form around the posterior mitral annulus from the anterolateral to the posteromedial commissure.<sup>6</sup> Additionally, mitral valve diseases are frequently associated with atrial fibrillation, for which the methods of treatment are novel percutaneous left atrial appendage closure procedures which also threaten the mentioned vessel.<sup>7</sup> Despite the great number of procedures mentioned earlier, the occurrence of complications on the circumflex coronary artery is rare in numbers. However, there are several case reports discussing the issue, which highlights the importance of being aware of this severe lifethreatening risk.<sup>8–12</sup>

In the review of literature, there are various studies addressing this topic. Some of them stressed the proximity of the examined arteries by measuring the mean distances at fix points on anatomic preparations,<sup>13–15</sup> while others evaluated intra- and perioperative imaging technologies to avoid iatrogenic complications.<sup>16–18</sup> Our study is aiming at emphasizing the proximity of the circumflex coronary artery to the annulus of the mitral valve by creating a three-dimensional (3D) model of the investigated region under exact anatomical conditions. Moreover, comparing coronary angiograms to anatomical preparations—as a radioanatomic correlation study—can help prevent any devastating injuries to the circumflex artery.

## **Materials and Methods**

In our research, 10 fresh adult human hearts (weight ranging from 290 to 670 g) removed from the chest by sectioning through the large vessels were studied. All the investigated hearts were free from any anamnestic or macroscopically observable pathological lesions that could have had any effect on our results. The cadavers (six males and four females; age ranging from 53 to 90 years) had been donated to the institution of the first author with the purpose of medical education and research.

#### **Coronary Angiographies**

Prior to the anatomical fixation, the left coronary artery was cannulated and subsequently injected with pure esophageal barium sulfate contrast agent (Microtrast, Guerbet, Roissy CdG Cedex, France). After 5 weeks of fixation in 4% formal-dehyde solution, coronary angiographies were performed (Innova 2100IQ, General Electric, Boston, United States). All the hearts were examined with the same three image exposure modes (LAO 40/cranial 20, RAO 10/cranial 40, and true lateral) as demonstrated in **~ Figs. 1–2** to **3**, "C–E."

#### **Anatomical Preparation**

In purpose of the proper anatomical preparation and visualization of the circumflex coronary artery, irrelevant parts of the hearts were removed. Thus, sectioning in an apicobasal direction with a plane going through the borders of the A1– A2 scallops and the P2–P3 scallops of the mitral valve, left atriotomy and ventriculotomy were performed. The anatomical preparation was performed in a rectangle shape following the rules of layering anatomical preparation. The bottom of the oblong was by the line of the mitral annulus, while the top border was the inferior edge of the left auricle. The left and the right borders of the rectangle were determined based on the coronary angiograms adjusted to reveal the full course of the investigated vessel.

Following, regardless of the level differences and the enclosed angles between the investigated structures, the closest distances were measured in every 5 mm between the circumflex coronary artery and the annular tissue starting from the anterolateral commissure of the mitral valve to the cleft between the P2–P3 scallops on each studied heart.

#### 3D Imaging

Colored photographs were taken (EOS 5D Mark II, Canon, Tokyo, Japan) from a surgical point of view from multiple angles, which allowed to create exact 3D modeling. Additionally, we integrated and paired the coronary angiograms with the anatomic preparations.

#### Results

The circumflex coronary artery was successfully made visible from a surgeon's point of view. 3D imaging models based on real anatomical preparations were successfully created to demonstrate exact anatomical conditions of hearts with each dominance pattern, such as right coronary dominance (**-Appendix A**), balanced coronary dominance (**-Appendix C**). (*PDF viewer which is able to open interactive 3D content is required. Instructions for opening in Adobe Acrobat Reader DC: 1. Download and open the PDF file. 2. Enable 3D content on the yellow pop-up bar by Options – Trust this document. 3. Clicking on the question mark in the upper left corner opens the interactive 3D model.*)

The main branches of the left coronary system with special regard on the circumflex coronary artery were evaluated as well as identified on both the coronary angiograms and the anatomical preparations on a heart with right (**Fig. 1B–E**), balanced (**Fig. 2B–E**), and left coronary dominance (**Fig. 3B–E**).

In each preparation—independently from the dominance pattern—the proximity of the circumflex artery to the mitral annulus was measured to be the closest accordingly to the P1 scallop  $(2.02 \pm 0.69 \text{ mm}; 1-3.1 \text{ mm})$ . This certain area was observed to concur anatomically with the orifice of the left auricle, as illustrated in **~Fig. 1-3**, "B." Following, the circumflex artery tended to make a significant loop away from the mitral annulus (8.97 ± 4.1 mm; 4.2–15.5 mm). Throughout the full course of the examined vessel, significant intermediate atrial or any other branches (more than 2 mm diameter) were not observed between the circumflex artery and the annular tissue.



**Fig. 1** Anatomical preparation of the circumflex artery compared with coronary angiograms on a heart with right coronary dominance. (A) The intact anatomical structure of the mitral valve and the mitral annulus in the left atrium. The vulnerable region is highlighted with a red rectangle. (B) The anatomical surrounding of the circumflex coronary artery. The main course of the artery is highlighted with white arrows. The mitral annulus is indicated by an interrupted line. (C) Coronary angiogram by RAO 10/cranial 40 projection. (D) Coronary angiogram by LAO 40/ cranial 20 projection. (E) Coronary angiogram by true lateral projection. 1, left auricle; 2, coronary sinus; 3, circumflex coronary artery; 4, left marginal coronary artery; 5, left anterior descending coronary artery; 6, posterior cusp of the mitral valve; 7, anterior cusp of the mitral valve.

Finally, regarding the coronary dominance, five hearts presented with right coronary dominance (50%), three hearts with balanced dominance (30%), and two hearts presented with left coronary dominance (20%).

# Discussion

Coronary angiography is part of the preoperative care serving the operator team to diagnose the coronary status beside the given mitral valve disease.<sup>19</sup> Though 3D imaging of the coronary vessels is achievable by switching between different viewpoints, the exact environmental relationship to the mitral valve cannot be revealed. Coronary computed tomography angiography is available for preoperative diagnostics, too. This method can solve the previous limitation by being able to visualize not just the full course of the coronary system but the anatomical surrounding as well, which might result more precise preoperative planning. With regard to the intraoperative screening, coronary angiography is performed as a standard routine while using Cardioband delivery system. Following the implantation of the first anchor, coronary angiography is performed to preclude any devastating vascular injuries to the circumflex coronary artery.<sup>20,21</sup> Despite the exact intraoperative screening during this catheter-based mitral annuloplasty, direct eye control is not available. On the contrary, open heart surgery allows direct eye control; however, intraoperative angiography is not accessible. Consequently, iatrogenic damage to the questionable vessels might be revealed with a delayed diagnosis in the postoperative terms. Though coronary angiography is not accessible during open heart surgery with median sternotomy, performing transesophageal echocardiography with modified midesophageal long axis might be an adequate method to preclude vascular injury to the circumflex artery.<sup>22,23</sup> In our study, we chose coronary angiograms as a reference to identify the examined coronary system. In addition, by having the anatomical preparations performed, real surgical point of view was achieved. Thus, we could extend the gold standard angiograms with a complete disclosure of the vicinity. Hereby, the simultaneous evaluation can result another viewpoint in risk assessment for the purpose of helping the surgeon to avoid any iatrogenic complications.



**Fig. 2** Anatomical preparation of the circumflex artery compared with coronary angiograms on a heart with balanced coronary dominance. (A) The intact anatomical structure of the mitral valve and the mitral annulus in the left atrium. The vulnerable region is highlighted with a red rectangle. (B) The anatomical surrounding of the circumflex coronary artery. The main course of the artery is highlighted with white arrows. The mitral annulus is indicated by an interrupted line. (C) Coronary angiogram by RAO 10/cranial 40 projection. (D) Coronary angiogram by LAO 40/ cranial 20 projection. (E) Coronary angiogram by true lateral projection. 1, left auricle; 2, coronary sinus; 3, left marginal coronary artery; 4, circumflex coronary artery; 5, posterolateral coronary artery; 6, left anterior descending coronary artery; 7, posterior cusp of the mitral valve; 8, anterior cusp of the mitral valve.

Regarding the proximity of the circumflex coronary artery to the annulus of the mitral valve, our data correlated with the results presented in previous articles.<sup>13–15,24,25</sup> These publications emphasize the risk of iatrogenic injury to the circumflex artery by measuring the mean distances at several points of the mitral annulus on the examined hearts. The measurement technique, we performed between the investigated structures, was similar to the methods described in previous articles. However, the importance of being aware of the proximity and mean distances in numbers is essential, we believe visualizing it from a surgeon's point of view might stress the proximity even more.

Taking into consideration the review of literature in a particular segment of the circumflex artery, special attention is advised. Previous case reports suggest that iatrogenic lesions secondary to mitral valve surgery might occur in the mid-portion,<sup>26,27</sup> but concordantly to our results mainly in the proximal segment of the investigated coronary artery.<sup>28–30</sup> This

certain proximal critical region is marked in **– Figs. 1–2** to **3**, "A" with a red rectangle as "danger zone." We observed a retraction on the left atrial wall under the orifice area of the left auricle. We believe this condition might play an important role in the proximity and vulnerability of the circumflex artery under this critical area.

The limitations of our study were, first of all, the small number of cases. Therefore, we could not draw exact conclusion on pattern of coronary dominance. However, 3D visualization of anatomical preparations of each dominance pattern was performed from a surgical viewpoint. Thus, it might increase the awareness of the susceptibility of the circumflex artery more effectively than quantitative results. Second, despite the fact that the full course of the circumflex artery was made visible on anatomical preparations, this method cannot be performed in vivo. Finally, having the examined hearts ex situ removed from the cadavers, moreover, the cannulation of the left coronary artery from the aortic root



**Fig. 3** Anatomical preparation of the circumflex artery compared with coronary angiograms on a heart with left coronary dominance. (A) The intact anatomical structure of the mitral valve and the mitral annulus in the left atrium. The vulnerable region is highlighted with a red rectangle. (B) The anatomical surrounding of the circumflex coronary artery. The main course of the artery is highlighted with white arrows. The mitral annulus is indicated by an interrupted line. (C) Coronary angiogram by RAO 10/cranial 40 projection. (D) Coronary angiogram by LAO 40/ cranial 20 projection. (E) Coronary angiogram by true lateral projection. 1, left auricle; 2, left marginal coronary artery; 3, circumflex coronary artery; 4, posterolateral coronary artery; 5, posterior descending coronary artery; 6, left anterior descending coronary artery; 7, posterior cusp of the mitral valve.

hardened the application of the standard projections for the circumflex artery on the coronary angiograms. Thus, in purpose of more exact and more efficient identification of the main arteries in concern, other projections were chosen.

# Conclusion

In conclusion, we believe this additional point of view and 3D visualization might help understand the anatomical vicinity of the affected annular region below the endocardial layer with special regard to the nearby circumflex coronary artery. Therefore, this research might contribute to avoid and reduce the incidence of iatrogenic complications resulting improved postoperative life expectancy.

#### Disclosures

The authors declare outside interests were not involved in neither the collection, analysis, and interpretation of the data nor the design of the study. Furthermore, the authors had freedom to fully disclose all results. Finally, no grants were received from any funding agencies in the public, commercials, or not-for-profit sectors.

#### Authors' Contribution

B.F.: project development, data collection, data analysis, manuscript writing; A.O.: data collection, data analysis; L.B.: data collection, project development; G.B.: project development; G.R.: data collection; T.R.: project development, data collection, data analysis, manuscript writing.

Funding None.

Conflict of Interest None declared.

## References

1 Kron IL, LaPar DJ, Acker MA, et al; American Association For Thoracic Surgery Ischemic Mitral Regurgitation Consensus Guidelines Writing Committee. 2016 update to The American Association for Thoracic Surgery consensus guidelines: ischemic mitral valve regurgitation. J Thorac Cardiovasc Surg 2017;153 (05):1076–1079

- 2 Chatterjee S, Rankin JS, Gammie JS, et al. Isolated mitral valve surgery risk in 77,836 patients from the Society of Thoracic Surgeons database. Ann Thorac Surg 2013;96(05):1587–1594, discussion 1594–1595
- 3 Nappi F, Avatar Singh SS, Santana O, Mihos CG. Functional mitral regurgitation: an overview for surgical management framework. J Thorac Dis 2018;10(07):4540–4555
- 4 Husain A, Alsanei A, Tahir M, Dahdouh Z, AlHalees Z, AlMasood A. Left circumflex artery injury postmitral valve surgery, single center experience. J Saudi Heart Assoc 2019;31(02):94–99
- 5 Aybek T, Risteski P, Miskovic A, et al. Seven years' experience with suture annuloplasty for mitral valve repair. J Thorac Cardiovasc Surg 2006;131(01):99–106
- <sup>6</sup> Nickenig G, Hammerstingl C, Schueler R, et al. Transcatheter mitral annuloplasty in chronic functional mitral regurgitation:
  <sup>6</sup>-month results with the Cardioband percutaneous mitral repair system. JACC Cardiovasc Interv 2016;9(19):2039–2047
- 7 Malakouti-Nejad B, Winkler E, Johnson M, et al. Evaluation of two methods of left atrial appendage (LAA) closure and its proximity to the circumflex artery. FASEB J 2015;29(1 Suppl);552.6
- 8 Banayan J, Dhawan R, Vernick WJ, McCarthy PM. CASE 3–2012. Iatrogenic circumflex artery injury during minimally invasive mitral valve surgery. J Cardiothorac Vasc Anesth 2012;26(03):512–519
- 9 Coutinho GF, Leite F, Antunes MJ. Circumflex artery injury during mitral valve repair: not well known, perhaps not so infrequentlessons learned from a 6-case experience. J Thorac Cardiovasc Surg 2017;154(05):1613–1620
- 10 Danielson GK, Cooper E, Tweeddale DN. Circumflex coronary artery injury during mitral valve replacement. Ann Thorac Surg 1967;4:53–59
- 11 Grande AM, Fiore A, Massetti M, Viganò M. latrogenic circumflex coronary lesion in mitral valve surgery: case report and review of the literature. Tex Heart Inst J 2008;35(02):179–183
- 12 Hiltrop N, Bennett J, Desmet W. Circumflex coronary artery injury after mitral valve surgery: a report of four cases and comprehensive review of the literature. Catheter Cardiovasc Interv 2017;89(01): 78–92
- 13 Cornu E, Lacroix PH, Christides C, Laskar M. Coronary artery damage during mitral valve replacement. J Cardiovasc Surg (Torino) 1995;36(03):261–264
- 14 Kaklikkaya I, Yeginoglu G. Damage to coronary arteries during mitral valve surgery. Heart Surg Forum 2003;6(06):E138–E142
- 15 Pessa CJN, Gomes WJ, Catani R, et al. Anatomical relationship between the posterior mitral valve annulus and the coronary arteries. Implications to operative treatment. Rev Bras Cir Cardiovasc 2004;19:372–377
- 16 Bevilacqua C, Eibel S, Foldyna B, et al. Analysis of circumflex artery anatomy by real time 3D transesophageal echocardiography compared to cardiac computed tomography. Int J Cardiovasc Imaging 2017;33(11):1703–1710

- 17 Ghersin N, Abadi S, Sabbag A, et al. The three-dimensional geometric relationship between the mitral valvar annulus and the coronary arteries as seen from the perspective of the cardiac surgeon using cardiac computed tomography. Eur J Cardiothorac Surg 2013;44(06):1123–1130
- 18 Wykrzykowska J, Cohen D, Zimetabum P. Mitral annuloplasty causing left circumflex injury and infarction: novel use of intravascular ultrasound to diagnose suture injury. J Invasive Cardiol 2006;18(10):505–508
- 19 Falk V, Baumgartner H, Bax JJ, et al. Corrigendum to '2017 ESC/EACTS Guidelines for the management of valvular heart disease' [*Eur J Cardiothorac Surg* 2017;52:616-664]. Eur J Cardiothorac Surg 2017;52(04):832
- 20 Kreusser MM, Pleger ST, Abu Sharar H, et al. Transient elevation of high-sensitive troponin T after Cardioband implantation. Herz 2019;44(06):546–552
- 21 Miura M, Zuber M, Gavazzoni M, et al. Possible left circumflex artery obstruction in a Cardioband transcatheter mitral annuloplasty caused by coronary kinking during cinching. JACC Cardiovasc Interv 2019;12(06):600–601
- 22 Ender J, Selbach M, Borger MA, et al. Echocardiographic identification of iatrogenic injury of the circumflex artery during minimally invasive mitral valve repair. Ann Thorac Surg 2010;89(06): 1866–1872
- 23 Murugesan C, Raghu B, Rao PV. Transesophageal echocardiographic diagnosis and management of circumflex artery injury following mitral valve repair. Cardiol Res 2011;2(02):90–92
- 24 Virmani R, Chun PK, Parker J, McAllister HA Jr. Suture obliteration of the circumflex coronary artery in three patients undergoing mitral valve operation. Role of left dominant or codominant coronary artery. J Thorac Cardiovasc Surg 1982;84 (05):773–778
- 25 Winkler E, Malakouti-Nejad B, Johnson M, et al. Establishing anatomical proximity between the coronary circumflex artery and the mitral valve annulus: implications for mitral valve surgery. FASEB J 2015;29(1 Suppl);552.7
- 26 Gentry JL III, Summers MR, Harb S, Jaber W, Pettersson G, Griffin B. Iatrogenic left circumflex coronary artery fistula after mitral valve replacement. CASE (Phila) 2018;2(05):218–221
- 27 Mantilla R, Legarra JJ, Pradas G, Bravo M, Sanmartín M, Goicolea J. [Percutaneous coronary intervention for iatrogenic occlusion of the circumflex artery after mitral anuloplasty]. Rev Esp Cardiol 2004;57(07):702–704
- 28 Aubert S, Barthélémy O, Landi M, Acar C. Circumflex coronary artery injury following mitral annuloplasty treated by emergency angioplasty. Eur J Cardiothorac Surg 2008;34(04):922–924
- 29 Calafiore AM, Iacò AL, Varone E, Bosco P, Di Mauro M. Distortion of the proximal circumflex artery during mitral valve repair. J Card Surg 2010;25(02):163–165
- 30 Scarsini R, De Maria GL, Ciofani J, et al. Coronary injury in mitral and aortic valve surgery: a case series using optical coherence tomography. Ann Thorac Surg 2020;109(03):e171–e174