

# Coagulopathy and Thrombosis as a Result of Severe COVID-19 Infection: A Microvascular Focus

Upendra K. Katneni<sup>1</sup> Aikaterini Alexaki<sup>2</sup> Ryan C. Hunt<sup>2</sup> Tal Schiller<sup>3</sup> Michael DiCuccio<sup>4</sup>  
Paul W. Buehler<sup>1</sup> Juan C. Ibla<sup>5</sup> Chava Kimchi-Sarfaty<sup>2</sup>

<sup>1</sup>Department of Pediatrics, The Center for Blood Oxygen Transport and Hemostasis, University of Maryland School of Medicine, Baltimore, Maryland, United States

<sup>2</sup>Hemostasis Branch, Division of Plasma Protein Therapeutics, Office of Tissues and Advanced Therapies, Center for Biologics Evaluation & Research, U.S. FDA, Silver Spring, Maryland, United States

<sup>3</sup>Diabetes, Endocrinology and Metabolic Disease Unit, Kaplan Medical Center, Rehovot, Israel

<sup>4</sup>National Center of Biotechnology Information, National Institutes of Health, Bethesda, Maryland, United States

<sup>5</sup>Division of Cardiac Anesthesia, Department of Anesthesiology, Perioperative and Pain Medicine, Boston Children's Hospital and Harvard Medical School, Boston, Massachusetts, United States

**Address for correspondence** Chava Kimchi-Sarfaty, Hemostasis Branch, Division of Plasma Protein Therapeutics, Office of Tissues and Advanced Therapies, Center for Biologics Evaluation & Research, US FDA, Silver Spring, MD 20993, United States (e-mail: Chava.kimchi-sarfaty@fda.hhs.gov).

Thromb Haemost 2020;120:1668–1679.

## Abstract

Coronavirus disease of 2019 (COVID-19) is the clinical manifestation of the respiratory infection caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). While primarily recognized as a respiratory disease, it is clear that COVID-19 is systemic illness impacting multiple organ systems. One defining clinical feature of COVID-19 has been the high incidence of thrombotic events. The underlying processes and risk factors for the occurrence of thrombotic events in COVID-19 remain inadequately understood. While severe bacterial, viral, or fungal infections are well recognized to activate the coagulation system, COVID-19-associated coagulopathy is likely to have unique mechanistic features. Inflammatory-driven processes are likely primary drivers of coagulopathy in COVID-19, but the exact mechanisms linking inflammation to dysregulated hemostasis and thrombosis are yet to be delineated. Cumulative findings of microvascular thrombosis has raised question if the endothelium and microvasculature should be a point of investigative focus. von Willebrand factor (VWF) and its protease, a disintegrin and metalloproteinase with a thrombospondin type 1 motif, member 13 (ADAMTS-13), play important role in the maintenance of microvascular hemostasis. In inflammatory conditions, imbalanced VWF-ADAMTS-13 characterized by elevated VWF levels and inhibited and/or reduced activity of ADAMTS-13 has been reported. Also, an imbalance between ADAMTS-13 activity and VWF antigen is associated with organ dysfunction and death in patients with systemic inflammation. A thorough understanding of VWF-ADAMTS-13 interactions during early and advanced phases of COVID-19 could help better define the pathophysiology, guide thromboprophylaxis and treatment, and improve clinical prognosis.

## Keywords

- ▶ COVID-19
- ▶ thrombosis
- ▶ inflammation
- ▶ ADAMTS-13
- ▶ von Willebrand factor

received  
May 14, 2020  
accepted after revision  
July 14, 2020

© 2020. Thieme. All rights reserved.  
Georg Thieme Verlag KG,  
Rüdigerstraße 14,  
70469 Stuttgart, Germany

DOI <https://doi.org/10.1055/s-0040-1715841>.  
ISSN 0340-6245.

## Introduction

Coronavirus disease of 2019 (COVID-19) is a respiratory illness caused by a novel coronavirus, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). SARS-CoV-2 is an enveloped, positive-sense single-stranded ribonucleic acid virus belonging to the *Coronaviridae* family.<sup>1</sup> The COVID-19 outbreak started in Wuhan, China, in late 2019 and rapidly spread to rest of the world. On March 11, 2020, the World Health Organization declared COVID-19 outbreak as pandemic. As of June 24, 2020, the global number of COVID-19 cases stood at 9.26 million with 478,000 deaths (Source: Johns Hopkins Coronavirus Resource Center, <https://coronavirus.jhu.edu/>). Disease course is markedly different between individuals while some are completely asymptomatic, others develop mild symptoms including mild fever, loss of taste or smell, dry cough, sore throat, shortness of breath, and myalgia.<sup>2-4</sup> In susceptible individuals, the disease progresses to pneumonia, hypoxemia, acute respiratory distress, and multiorgan dysfunction that may lead to death.<sup>3</sup> The predominance of asymptomatic or mild infections has contributed to the rapid spread of COVID-19 compared with earlier coronavirus outbreaks of SARS and Middle East respiratory syndrome in 2002 and 2012, respectively.<sup>4,5</sup>

## Consumptive Coagulopathy and the High Incidence of Thrombosis in COVID-19 Patients

Altered coagulation is a common feature of acute systemic diseases, specifically to those affecting primarily the respiratory system. Based on studies in patients with acute respiratory distress syndrome (ARDS), the coexistence of disseminated intravascular coagulation (DIC) with subsequent consumption of procoagulation proteins and platelets has been consistently described.<sup>6</sup> This in turn leads to the formation of microthrombi in the vascular bed of organs resulting from excess coagulation byproducts and suppression of endogenous anticoagulation factors.<sup>7</sup> The coexistence of consumptive coagulopathy and thrombosis are the result of a common pathologic pathway; however, the exact mechanisms that tilts the balance toward thrombosis in COVID-19 are less well understood.<sup>8</sup> In this sense, some features of the coagulopathy associated with COVID-19 may be not unique to this disease; however, the magnitude of the thrombotic response and its impact on mortality suggests the presence of additional mechanisms, beyond what is known for similar respiratory acute inflammatory diseases.

Several studies have linked coagulation abnormalities to severe COVID-19 illness<sup>9,10</sup> (► **Table 1**). In a study evaluating 449 severe COVID-19 patients, Tang et al<sup>11</sup> reported positive correlation of 28-day mortality with fibrin degradation product (FDP), D-dimers and prothrombin time (PT), and negative correlation with platelet count. Laboratory parameters were recorded at the time of onset of severe COVID-19 in the study. In an earlier study comprising 183 patients, Tang et al<sup>12</sup> reported elevated D-dimer levels and FDP levels and prolonged PT and activated partial thromboplastin times (aPTTs) at the time of admission in nonsurvivors compared

with survivors. In the same study, significantly lower levels of fibrinogen and antithrombin levels were observed during the late hospitalization in nonsurvivors. Huang et al<sup>13</sup> reported higher D-dimers and prolonged PT at the time of admission in intensive care unit (ICU) patients compared with non-ICU patients in a study of 41 patients. Wang et al<sup>14</sup> reported elevated PT in a study of 138 patients. In the same study, elevated levels of D-dimers were found in ICU patients compared with non-ICU patients as well as in survivors compared with nonsurvivors in a subgroup of patients with a definitive outcome. In a study of 94 COVID-19 patients, Han et al<sup>15</sup> reported lower antithrombin and higher D-dimers, FDP, and fibrinogen levels compared with healthy controls. Zhou et al<sup>16</sup> reported an association of elevated D-dimers with in-hospital death in a study of 191 patients. Also, elevated PT and decreased platelet counts were observed in nonsurvivors compared with survivors. Elevated levels of D-dimers were reported by Richardson et al<sup>17</sup> among 5,700 patients in the New York City area. Ranucci et al<sup>18</sup> reported a procoagulant profile in 16 patients characterized by increased clot strength by viscoelastography, elevated D-dimer levels, and hyperfibrinogenemia. A meta-analysis of 9 studies encompassing 1,779 patients with severe disease has identified significantly lower platelet counts.<sup>19</sup> A subgroup analysis based on survival has identified even lower platelet counts in nonsurvivors in this study. Litjós et al<sup>20</sup> and Helms et al<sup>7</sup> reported elevated D-dimer and fibrinogen levels in 26 and 150 ICU-admitted patients, respectively. Overall, elevated PT, increased D-dimer and fibrinogen levels, and thrombocytopenia are frequently reported in COVID-19 patients. However, bleeding events requiring therapeutic intervention are not reported.

Multiple studies have reported a higher incidence of thrombotic events, particularly pulmonary embolism, as a frequent complication in COVID-19 patients (► **Table 1**). Litjós et al<sup>20</sup> reported overall rate of 69% venous thromboembolism (VTE) in severe COVID-19 patients admitted to ICU. In this study, VTE incidence was found to be significantly higher in patients treated with prophylactic anticoagulation compared with those treated with therapeutic anticoagulation. Helms et al<sup>7</sup> reported 64 clinically relevant thrombotic complications in 150 ICU-admitted patients. Importantly, the incidence of thrombotic complications in COVID-19 ARDS patients was significantly higher than non-COVID-19 ARDS patients in this study. Ackermann et al<sup>21</sup> compared lung sections of COVID-19 patients with those died from ARDS secondary to influenza A (H1N1) infection and found relatively higher: (1) endothelial cell injury, (2) alveolar microthrombi (ninefold), and (3) intussusceptive angiogenesis in COVID-19 lung sections. Similarly, higher incidence of thromboembolic complications in ICU-admitted COVID-19 patients was also reported by Klok et al (31%),<sup>22</sup> Lodigiani et al (27.6%),<sup>23</sup> Middeldorp et al (47%),<sup>24</sup> Nahum et al (79%),<sup>25</sup> and Cui et al (25%).<sup>26</sup> For comparison, in a study by Zhang et al, the reported cumulative incidence of VTE in ICU-admitted patients receiving guideline-recommended thromboprophylaxis was 9.55% (95% confidence interval: 6.55–13.81).<sup>27</sup>

**Table 1** Studies (multiple patients) reporting abnormal coagulopathy in COVID-19

Study	Type of study, number of patients	Findings/Significance
<b>Clinical features of COVID-19 patients, coagulation parameters included</b>		
Huang et al <sup>13</sup>	Prospective, 41 patients	Prothrombin time and D-dimer levels on admission were higher in patients that required ICU treatment
Zhou et al <sup>16</sup>	Retrospective, 191 COVID-19 patients	Increased D-dimer on admission is associated with poor prognosis
Guan et al <sup>44</sup>	Retrospective, 1,099 COVID-19 patients	Thrombocytopenia in 36.2%
Goyal et al <sup>36</sup>	Retrospective, 393 COVID-19 patients	Thrombocytopenia in 27%
Zhu et al <sup>45</sup>	Meta-analysis	Elevated D-dimer in ~37.2% of patients
<b>Studies on coagulation parameters</b>		
Ranucci et al <sup>18</sup>	Prospective, 16 ARDS COVID-19 patients	Patients showed a procoagulant profile (clot strength, platelet, fibrinogen, D-dimers, hyperfibrinogenemia)
Tang et al <sup>12</sup>	Retrospective, 183 COVID-19 patients	Nonsurvivors had significantly higher D-dimer and fibrin degradation product (FDP) levels, longer prothrombin time, and activated partial thromboplastin time compared with survivors on admission. 71.4% of nonsurvivors and 0.6% survivors met the criteria of DIC during their hospital stay
Lippi et al <sup>19</sup>	Meta-analysis	Low platelet count associated with increased risk of severe disease and mortality in patients with COVID-19
Zhang et al <sup>29</sup>	Retrospective, 343 COVID-19 patients	Patients with D-dimer levels $\geq 2.0$ $\mu\text{g/mL}$ had a higher incidence of mortality when comparing to those who with D-dimer levels $< 2.0$ $\mu\text{g/mL}$
Escher et al <sup>108,109</sup>	Case study, 1 patient and 3 more in the follow-up publication	Continual increase of D-dimers, elevated FVIII activity, and normal platelet counts
Bowles et al <sup>112</sup>	216 COVID-19 patients 34 tested for lupus anticoagulant	91% of patients tested positive for lupus anticoagulant. All lupus anticoagulant-positive specimens had a prolonged aPTT. Increased aPTT should not be a reason to withhold anticoagulation therapy
Lorenzo-Villalba et al <sup>115</sup>	Case reports, 3 patients	Severe thrombocytopenia during COVID-19 infection associated with either cutaneous purpura or mucosal bleeding
Yin et al <sup>116</sup>	Retrospective, 449 COVID-19 and 104 non-COVID severe pneumonia	Patients with severe pneumonia induced by SARS-CoV-2 had higher platelet count than those induced by non-SARS-CoV-2. Patients infected by SARS-CoV-2 may benefit from anticoagulant treatment, if they have markedly elevated D-dimer
Tabatabai et al <sup>48</sup>	Case series, 10 patients	Elevated FVIII activity and low normal antithrombin and functional protein C activity
<b>Thrombosis in the COVID-19 patients</b>		
Middeldorp et al <sup>24</sup>	Retrospective, 198 patients	The cumulative incidences of VTE at 7, 14, and 21 days were 16%, 33%, and 42%, respectively. VTE was higher in the ICU and was associated with death
Nahum et al <sup>25</sup>	Prospective, 34 patients	Deep vein thrombosis was found in 22 patients (65%) at admission and in 27 patients (79%) when the venous ultrasonograms performed 48 hours after ICU admission were included. D-dimers and fibrinogen were also increased
Cui et al <sup>26</sup>	Retrospective, 81 severe COVID-19 patients	Incidence of VTE at 25%. D-dimer increase has a predictive value
Klok et al <sup>22</sup>	Retrospective, 184 patients, no control group	31% cumulative incidence of symptomatic acute pulmonary embolism (PE), deep vein thrombosis, ischemic stroke, myocardial infarction, or systemic arterial embolism in COVID-19 patients

**Table 1** (Continued)

Study	Type of study, number of patients	Findings/Significance
Zhang et al <sup>27</sup>	Prospective, 281 ICU COVID-19 patients	Cumulative incidence of VTE at 28 days was 9.55%, despite all patients receiving thromboprophylaxis
Demelo-Rodríguez et al <sup>117</sup>	Prospective, 156 COVID-19 patients	D-dimer levels > 1,570 ng/mL were associated with asymptomatic DVT
Grandmaison et al <sup>118</sup>	Cross-sectional study, 58 COVID-19 patients, 29 in the ICU and 29 in the medicine ward	In the ICU, VTEs were found in 17 (58.6%) of the 29 patients In the medicine ward, VTEs were found in 6 (20.7%) patients
Fraissé et al <sup>119</sup>	Retrospective, 92 ICU COVID-19 patients	High rate of thrombotic events (TEs) in ICU COVID-19 patients highlighting the necessity for thromboprophylaxis and TE screening. Hemorrhagic events (HEs) were also observed in patients on full-dose anticoagulation
Jian et al <sup>114</sup>	Retrospective, 3,218 COVID-19 patients	Acute stroke was the most common neuroimaging finding, present in 1.1% of hospitalized COVID-19 patients
Desborough et al <sup>121</sup>	Retrospective, 66 patients	10 patients had at least one proven episode of thromboembolism. Major bleeding occurred in seven cases
Akel et al <sup>122</sup>	Case reports, 6 patients	Patients did not have any hypercoagulable risk factors yet presented with pulmonary embolism
Kashi et al <sup>123</sup>	Case reports, 7 patients	Arterial thrombosis
Lax et al <sup>124</sup>	Prospective autopsy study, 11 deceased COVID-19 patients	Death may be caused by the thrombosis observed in segmental and subsegmental pulmonary arterial vessels despite the use of prophylactic anticoagulation
Thomas et al <sup>125</sup>	Retrospective, 63 COVID-19 patients	High thrombotic risk in patients with COVID-19
Gomez-Arbelaez et al <sup>126</sup>	Case reports, 4 patients	Aortic thrombosis and associated ischemic complications in patients with severe SARS-CoV-2 infection
<b>Anticoagulation treatment in COVID-19 patients</b>		
Tang et al <sup>11</sup>	Retrospective, 449 severe COVID-19 patients, 99 received heparin	Anticoagulant therapy is associated with better prognosis in severe COVID-19 patients with sepsis induced coagulopathy or markedly elevated D-dimer
Wang et al <sup>28</sup>	3 case reports	Treatment with tissue plasminogen activator lead to improvement in the respiratory status
Ayerbe et al <sup>127</sup>	2,075 COVID-19 patients, admitted in 17 hospitals in Spain	Heparin had been used in 1,734 patients. Heparin was associated with lower mortality
Wang et al <sup>128</sup>	Retrospective, 1,099 COVID-19 patients	High risk of venous thromboembolism, also high risk of bleeding
Artifoni et al <sup>129</sup>	Retrospective, 62 patients	16 patients developed VTE, 7 patients developed PE Very high negative predictive value of baseline D-dimer level for VTE and PE
Russo et al <sup>130</sup>	Retrospective, 192 COVID-19 patients	Preadmission antithrombotic therapy, both antiplatelet and anticoagulant, does not seem to show a protective effect in severe forms of COVID-19 with ARDS at presentation and rapidly evolving toward death
<b>Link between SARS-CoV-2 and thrombosis</b>		
Ackermann et al <sup>21</sup>	7 lung autopsies from COVID-19 patients and 7 from ARDS	Vascular angiogenesis distinguished the pulmonary pathobiology of COVID-19 from that of equally severe influenza virus infection
Maier et al <sup>131</sup>	Case studies 15 COVID-19 patients with hyperviscosity	Possible causal relationship between hyperviscosity and thrombotic complications in COVID-19

(Continued)

**Table 1** (Continued)

Study	Type of study, number of patients	Findings/Significance
Huisman et al <sup>105</sup>	12 COVID-19 patients	Low ADAMTS-13 activity, increased VWF levels and factor VIII levels
Galeano-Valle et al <sup>111</sup>	Prospective study, 24 patients	Prevalence of antiphospholipid antibodies in COVID-19 and venous thrombosis was low
Magro et al <sup>132</sup>	Case reports, 5 severe COVID-19 cases	Procoagulant state is associated with systemic complement activation

Abbreviations: ADAMTS-13, a disintegrin and metalloproteinase with a thrombospondin type 1 motif, member 13; aPTT, activated partial thromboplastin time; ARDS, acute respiratory distress syndrome; COVID-19, coronavirus disease of 2019; DIC, disseminated intravascular coagulation; DVT, deep vein thrombosis; FVIII, factor VIII; ICU, intensive care unit; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; VTE, venous thromboembolism; VWF, von Willebrand factor.

A high incidence of DIC diagnosed by D-dimer, fibrinogen, and antithrombin III levels has become a focus for the initiation of anticoagulation therapy in severe COVID-19 patients,<sup>28</sup> with some studies relying on D-dimers alone.<sup>11,29</sup> A retrospective analysis of 183 patients performed by Tang et al<sup>12</sup> suggested that more than 70% of severe COVID-19 patients who succumb to the infection demonstrate increased risk of thrombosis, further this group suggests that all of these patients meet the International Society on Thrombosis and Haemostasis definition of DIC. Subsequently, Tang et al<sup>11</sup> reported an equivalent 28-day mortality rate (30%) in 99 patients receiving low molecular weight or unfractionated heparin for 7 days compared with 350 nonheparin treated patients or those receiving a less than 7-day course of therapy. A case series reported by Wang et al<sup>28</sup> detailed the use and outcome following tissue plasminogen activator (tPA) in three patients with ARDS and coagulopathy consistent with DIC. Intravenous dosing with tPA indicated a potential benefit in each of the three cases of COVID-19. However, this study also warns of both unrelated effects and high risk of severe bleeding secondary to off-label tPA use. Several of the studies in coagulopathic COVID-19 patients suspected of DIC rely heavily on analysis of fibrin degradation and D-dimer levels, which are expected to be increased during DIC, arterial and venous thromboses, strokes, and thrombotic microangiopathies.<sup>30</sup> However, D-dimers are a nonspecific indicator of thrombosis in severe COVID-19 patients with pulmonary injury. Fibrin accumulation and lysis continuously occur during nonthrombotic inflammation as well as tissue necrosis, and therefore, significant D-dimer elevations also accumulate during cancers<sup>31</sup> and infections, consistent with inflammatory processes that coincide with the progression of severe COVID-19-related macrophage activation syndrome.<sup>32</sup> Therefore, we suggest that more comprehensive and robust assays be used to evaluate changes in hemostasis. For example, to date the use of thrombin, plasmin, or simultaneous thrombin/plasmin generation assays have not been reported within the context of hemostasis management of COVID-19 patients. Since their introduction thrombin and plasmin generation assays have been highly informative regarding the assessment of hemorrhage, coagulation, and fibrinolysis.<sup>33,34</sup> Assessment of impairment of these systems would provide a useful and appropriate guidance needed for and monitoring of therapeutic interventions in the unique coagulopathies associated with COVID-19.<sup>33,34</sup> Because patients are often on unfractionated

or low molecular weight heparin and plasminogen activator inhibitor 1, von Willebrand factor (VWF), plasminogen, fibrinogen, and factor VIII are all reported to be elevated in SARS infection,<sup>35</sup> and therefore careful modification of these assays may be warranted to optimize the concentrations of added tPA, tissue factor, and thrombomodulin.

These studies present a heterogeneous picture that is difficult to evaluate in the aggregate. Inclusion criteria for patients varied across these studies, making direct comparisons between the studies difficult. Further, the studies used different regimens of thromboprophylaxis, which could impact outcomes. In some studies, a high proportion of patients were still hospitalized at the end of the reporting period; conclusions and clinical courses therefore were based on incomplete information, and completion of these patients' clinical course could alter the final conclusions. The picture of coagulopathy in COVID-19 is complex. Specific, sensitive, and temporal assessments of coagulation and fibrinolysis should be established and further work is needed to untangle the roles of the host inflammatory response, preexisting thrombotic risk, and prehospitalization pharmacologic regimens in the optimal management of coagulopathy in the setting of COVID-19.

### Inflammation, Liver Injury, and Hypoxia in COVID-19 Patients

The risk of hospitalization, morbidity, and mortality from COVID-19 is highest for older patients with preexisting conditions such as hypertension, diabetes, cardiovascular disease, and obesity.<sup>13,14,16,17,36,37</sup> A common theme of all these comorbidities is their association with vascular inflammation and endothelial dysfunction.<sup>38,39</sup> Proinflammatory conditions affect hemostasis by blocking of fibrinolysis and induction of prothrombotic conditions through activation of endothelial cells and innate immune cells via release of several factors including tissue factor, VWF, and neutrophil extracellular traps (NETs) that promote thrombosis.<sup>40</sup> Induction of proinflammatory conditions was reported in the pathophysiology of several viral diseases including influenza and SARS.<sup>41</sup> Increased inflammation is commonly observed in COVID-19 patients, while severe cases are characterized by immune dysregulation and hyperinflammation, with a markedly increased serum interleukin (IL)-6.<sup>42</sup> Cytokine release syndrome has also been reported in COVID-19 patients and correlates with adverse

**Table 2** Studies reporting elevated inflammatory markers in COVID-19

Study	Patient group (number of patients) comparison	Elevated inflammatory markers
Huang et al <sup>13</sup>	ICU (13) vs. non-ICU (28)	Procalcitonin, IL-1 $\beta$ , IFN- $\gamma$ , IP10, and MCP1
Wang et al <sup>14</sup>	ICU (36) vs. non-ICU (102)	Procalcitonin
Zhou et al <sup>16</sup>	Nonsurvivor (54) vs. survivor (137)	Procalcitonin, ferritin, and IL-6
Richardson et al <sup>17</sup>	Relative to reference range (3066)	Procalcitonin, ferritin, and CRP
Ruan et al <sup>37</sup>	Nonsurvivor (68) vs. survivor (82)	CRP and IL-6
Giamarellos-Bourboulis et al <sup>42</sup>	Dysregulated (21) vs. intermediate state (26) of immune activation	CRP and IL-6
Chen et al <sup>47</sup>	Severe ( $\geq 9$ ) vs. moderate ( $\geq 7$ )	CRP, ferritin, IL-6, and TNF- $\alpha$
Han et al <sup>49</sup>	COVID-19 patients (102) vs. controls (45)	CRP, IL-6, TNF- $\alpha$ , and IFN- $\gamma$
Du et al <sup>50</sup>	Mild pneumonia (124) vs. no pneumonia (54) (pediatric patients)	Procalcitonin, IL-6, TNF- $\alpha$ , and IFN- $\gamma$
Wang et al <sup>52</sup>	SpO <sub>2</sub> $\geq 90\%$ ( $\geq 36$ ) vs. SpO <sub>2</sub> $< 90\%$ ( $\geq 7$ )	CRP and IL-6
Tan et al <sup>53</sup>	Severe (25) vs. mild/moderate (31)	CRP and IL-6
Tabatabai et al <sup>48</sup>	Relative to reference range (10)	Fibrinogen, CRP, and ferritin

Abbreviations: COVID-19, coronavirus disease of 2019; CRP, C-reactive protein; ICU, intensive care unit; IFN- $\gamma$ , interferon- $\gamma$ ; IL-1 $\beta$ , interleukin-1 $\beta$ ; IL-6, Interleukin-6; IP-10, interferon- $\gamma$  induced protein 10; MCP-1, monocyte chemotactic protein-1; SpO<sub>2</sub>, blood oxygen saturation level; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ .

clinical outcomes.<sup>43</sup> The presence of several inflammatory markers such as C-reactive protein, procalcitonin, ferritin, and fibrinogen are often reported in COVID-19 patients<sup>13,14,16,17,36,37,44–48</sup> (► **Table 2**). Further, multiple studies reported elevated levels of the proinflammatory cytokine IL-6 in severe cases of COVID-19<sup>16,37,42,47,49–53</sup> (► **Table 2**). A concurrent increase in the levels of anti-inflammatory cytokine IL-10, probably in response to overwhelming systemic inflammation, was also observed in several studies. The role of IL-6, in particular, is considered central in the pathogenesis of COVID-19 complications,<sup>54</sup> and therefore tocilizumab, an IL-6 inhibitor, is being used in ongoing clinical trials to prevent catastrophic inflammation.<sup>55–58</sup>

Liver injury during COVID-19 infections was described in multiple studies, including elevated levels of alanine aminotransferase, aspartate aminotransferase, and bilirubin.<sup>14,16,17,36,44,47</sup> The liver is the primary source of plasma proteins, particularly those involved in hemostasis. Thus, the occurrence of liver injury may contribute further to derangements of key hemostasis proteins and contributes to coagulopathy.<sup>59</sup> Similarly, hypoxemia observed in COVID-19 patients induces prothrombotic conditions through upregulation of plasminogen activator inhibitor and stimulation of endothelial synthesis of procoagulants, including tissue factor and VWF.<sup>60–63</sup> Thus, multiple clinical characteristics observed in COVID-19 patients contribute to altered coagulation and lead to increased incidence of thrombosis. However, the early onset of coagulopathy—before systemic organic effects occur—suggests proinflammatory conditions as the primary driving cause of thrombotic events in COVID-19 patients.

## VWF-ADAMTS-13 in Hemostasis and Thrombosis

VWF and its cleaving protease, a disintegrin and metalloproteinase with a thrombospondin type 1 motif, member

13 (ADAMTS-13), play an important role in hemostasis particularly within the microvasculature.<sup>64</sup> VWF is a large multimeric glycoprotein primarily expressed by endothelial cells and platelets. Endothelial cells show both basal secretion and regulated release of VWF stored in Weibel–Palade bodies in response to various stimuli. On the other hand, platelets secrete VWF stored in  $\alpha$ -granules only upon activation.<sup>65</sup> ADAMTS-13 is expressed both by hepatic stellate cells and endothelial cells; the relative contribution of hepatic and microvascular expression is not clear.<sup>66</sup> ADAMTS-13 regulates the biological activity of VWF by cleaving prothrombotic ultra-large VWF multimers ( $> 10,000$  kDa) secreted from endothelial cells into hemostatically active high molecular weight multimers ( $< 10,000$  kDa) under shear stress conditions.<sup>67</sup> Severe deficiency of ADAMTS-13 results in accumulation of ultra-large VWF multimers leading to microvascular thrombosis and consumptive thrombocytopenia, a condition termed thrombotic thrombocytopenic purpura (TTP).<sup>64</sup> In the event of vascular injury, VWF facilitates binding of platelets to subendothelium through its interactions with glycoprotein Ib and collagen, thereby inducing thrombus formation.<sup>64</sup> A reciprocal relationship exists between VWF and ADAMTS-13 levels where elevated circulatory VWF antigen levels are associated with concomitant decrease in ADAMTS-13 activity and vice versa.<sup>68–70</sup> Abnormal VWF-ADAMTS-13 ratios are implicated in arterial thrombosis,<sup>71</sup> ischemic stroke,<sup>72,73</sup> pediatric stroke,<sup>74</sup> and perioperative thrombosis in infants.<sup>75</sup> In addition, abnormal VWF/ADAMTS-13 metabolism has been positively associated with myocardial infarction in young women.<sup>76</sup> It is worth highlighting that in the case of perioperative thrombosis, elevated VWF even in the absence of significant deficiency of ADAMTS-13 was associated with thrombosis.<sup>75</sup> Severe hypoxia and acidosis likely caused a higher increase in VWF during cardiac surgery and were at higher risk of thrombosis.<sup>75</sup>

Elevated levels of VWF are found in several inflammatory and metabolic disorders including diabetes, obesity, and sickle cell disease.<sup>77</sup> In patients with systemic inflammatory response syndrome, active VWF predicted 28-day mortality.<sup>78</sup> VWF is an acute-phase response protein released by activated endothelial cells in response to inflammatory stimuli.<sup>77</sup> Inflammatory cytokines, IL-8 and tumor necrosis factor- $\alpha$  induced the release of VWF from human umbilical vein endothelial cells.<sup>79</sup> VWF released in inflammation binds to NETs released from activated neutrophils and recruits platelets and leukocytes to promote thrombosis.<sup>77</sup> ADAMTS-13 deficiency in inflammatory conditions was demonstrated to promote VWF-dependent leukocyte adhesion and extravasation in mice.<sup>80</sup>

In patients with systemic inflammation, ADAMTS-13 activity decreases proportional to the inflammatory response; an imbalance between ADAMTS-13 activity and VWF antigen is associated with organ dysfunction and death.<sup>81,82</sup> Dysregulated host response to infection including inflammation can result in septic shock. In septic shock, ADAMTS-13 activity was significantly lower<sup>83–85</sup> and elevated ratio of VWF propeptide (VWFpp) that is secreted along with ultra-large VWF multimers in to blood stream and ADAMTS-13 was associated with disease severity.<sup>86</sup> In patients with DIC, ADAMTS-13 activity decreased with DIC score<sup>87</sup> and VWFpp/ADAMTS-13 ratio was significantly elevated in nonsurvivors compared with survivors.<sup>88</sup> An interesting observation is that smoking, which is associated with adverse outcomes in COVID-19 patients,<sup>89</sup> was also found to be associated with decreased plasma ADAMTS-13 levels in a study of 3,244 individuals.<sup>90</sup> Increased expression of angiotensin-converting enzyme 2, the entry receptor for SARS-CoV-2, in the small airway epithelia of smokers was suggested as the potential mechanism for increased risk of severe COVID-19 in smokers.<sup>91</sup> Smoking is also associated with increased inflammatory markers.<sup>92</sup>

The imbalance between ADAMTS-13 and VWF in heightened inflammation could be a result of inhibition and/or deficiency of ADAMTS-13 activity.<sup>93</sup> The inhibition of VWF cleavage by ADAMTS-13 in inflammatory conditions was suggested to be mediated by several mechanisms: (1) thrombospondin-1 released from  $\alpha$ -granules of activated platelets by binding to the A2–A3 domain of VWF<sup>94,95</sup>; (2)  $\alpha$ -defensins released from neutrophils by binding to the A2 domain of VWF<sup>96</sup>; and (3) oxidation of Met 1606 residue in the ADAMTS-13 cleavage site of VWF.<sup>97</sup> Moreover, nonphysiological high concentrations of IL-6 have been shown to inhibit cleavage of VWF by ADAMTS-13 in vitro under shear flow conditions.<sup>79</sup> Granulocyte elastases, plasmin, and thrombin that are elevated in inflammatory conditions lower ADAMTS-13 activity through its proteolytic cleavage.<sup>98,99</sup>

## VWF-ADAMTS-13 Interactions in COVID-19

Despite playing an important role in the maintenance of hemostasis and the occurrence of micro- and macrovascular thrombosis, VWF-ADAMTS-13 interactions have not received much investigative attention in the evaluation of COVID-19 pathophysiology, specifically in relation to elevated incidence of VTE. Importantly, reduced ADAMTS-13 activity has been shown to correlate with increased inflammation in multiple sys-

tems,<sup>100–102</sup> while IL-6 has been shown to inhibit the cleavage of ultra-large VWF strings by ADAMTS-13 under flowing conditions.<sup>79,103</sup> The authors could find only five studies evaluating both VWF and ADAMTS-13 levels in COVID-19 patients in literature<sup>104–108</sup> (► **Table 3**). Majority of these studies reported lower ADAMTS-13 activity concurrent with higher VWF in COVID-19 patients.<sup>104–107</sup> In one of these studies, Bazzan et al<sup>104</sup> reported lower ADAMTS-13 levels in 88 COVID-19 patients compared with healthy controls ( $48.71 \pm 18.7\%$  vs. healthy control,  $108 \pm 9.1\%$ ; normal value 60–130%). Within patient cohort, lower ADAMTS-13 and higher VWF levels were found in nonsurvivors (9/88) compared with survivors. Further, lower than 30% ADAMTS-13 activity were significantly associated with mortality in survivor analysis. Huisman et al<sup>105</sup> observed low ADAMTS-13 activity levels ( $0.48 \pm 0.14$  IU/mL against a reference range of 0.61–1.31) in parallel with elevated VWF antigen and activity ( $\sim$  fourfold) in 12 ICU-admitted patients. A similar reduction in ADAMTS-13 and increased VWF levels was also reported by Adam et al<sup>106</sup> and Latimer et al<sup>107</sup> in 4 adult and 1 pediatric patients, respectively. On the other hand, Escher et al<sup>108</sup> observed normal to lower-normal ADAMTS-13 levels concurrently with  $> 2.5$ -fold increase in VWF antigen and activity in 3 ICU-admitted patients. Two other studies<sup>7,109</sup> reported VWF measurements alone, observing  $>$  threefold increase in both VWF antigen and activity. From the limited number of studies so far, it appears that COVID-19 infection may be characterized by markedly elevated VWF levels and below normal ADAMTS-13 activity. However, the current literature is limited by the small number of studies and variable timing of VWF/ADAMTS-13 measurements in relation to disease onset. Further evaluation of VWF and ADAMTS-13 interactions in large patient cohorts are warranted to more confidently understand their contributions to COVID-19 pathogenesis.

A secondary mechanism potentially contributing to ADAMTS-13 deficiency relates to the antiphospholipid antibody generation during SARS-CoV-2 infection.<sup>7,110–112</sup> Antiphospholipid antibodies have been inconsistently reported in all cases of COVID 19,<sup>7,111,112</sup> but strongly associated to prolong aPTT as reported by Bowles et al.<sup>112</sup> Patients with antiphospholipid syndrome have been found to have abnormal ADAMTS-13 plasmatic activity further increasing the risk of thrombosis.<sup>113</sup> The exact mechanisms by which antiphospholipid antibodies interfere with ADAMTS-13 cleaving activity are unclear. We speculate that antiphospholipid antibodies generated during active SARS-CoV-2 infection can potentially bind the spacer domain of ADAMTS-13 interfering with the recognition and proteolysis of VWF. Such a mechanism is similar to the binding of autoantibodies against ADAMTS-13 present in TTP resulting in clinical thrombosis.<sup>114</sup>

Based on the limited available data, we propose a mechanistic model in which: (1) SARS-CoV-2 causes endothelial activation and damage leading to overwhelming VWF release and (2) proinflammatory mediators or antibodies during the severe phase of COVID-19 result in reduced cleavage of high molecular weight VWF by ADAMTS-13, ultimately leading to thrombosis, see ► **Fig. 1**. This concept should be confirmed by large patient cohorts that encompass mild and severe clinical courses of COVID-19 disease. A mechanistic understanding of thrombosis

**Table 3** Studies reporting ADAMTS-13 and VWF levels in COVID-19


Study	Patient group (number of patients) comparison	Findings/Significance
Bazzan et al <sup>104</sup>	Nonsurvivor (9) vs. survivor (79)	Lower ADAMTS-13 and elevated VWF levels in nonsurvivors compared with survivors. After survival analysis, lower than 30% ADAMTS-13 levels were significantly associated with higher mortality
Huisman et al <sup>105</sup>	Relative to reference range (12)	Lower ADAMTS-13 and elevated VWF levels
Adam et al <sup>106</sup>	Relative to reference range (4)	Lower ADAMTS-13 and elevated VWF levels
Latimer et al <sup>107</sup>	Relative to reference range (1 pediatric patient)	Lower ADAMTS-13 and elevated VWF levels
Escher et al <sup>108,109</sup>	Case study, 1 patient and 3 more in the follow-up publication	Massive elevation of VWF and normal to lower-normal ADAMTS-13 activity. COVID-19 coagulopathy may be a distinct entity of highly prothrombotic alterations most probably an endothelial disease
Helms et al <sup>7</sup>	Relative to reference range (150)	Elevated VWF levels

Abbreviations: ADAMTS-13, a disintegrin and metalloproteinase with a thrombospondin type 1 motif, member 13; COVID-19, coronavirus disease of 2019; VWF, von Willebrand factor.

### A) Hemostasis



### B) Inflammation



### C) Cleavage of VWF by ADAMTS13 is prevented by the following mechanisms

- 1) Binding of thrombospondin-1 released from  $\alpha$ -granules of activated platelets to A2-A3 domain of VWF harboring the proteolytic cleavage site of ADAMTS13
- 2) Binding of  $\alpha$ -defensins released from neutrophils to A2 domain of VWF
- 3) Oxidation of Met 1606 residue in the ADAMTS13 cleavage site of VWF by reactive oxygen species
- 4) High concentrations of IL-6
- 5) Proteolytic cleavage of ADAMTS13 by granulocyte elastases, plasmin, and thrombin that are elevated in inflammatory conditions

**Fig. 1** von Willebrand factor (VWF)-a disintegrin and metalloproteinase with a thrombospondin type 1 motif, member 13 (ADAMTS-13) metabolism in inflammation. (A) During normal homeostasis, ADAMTS-13 regulates the activity of VWF by cleaving prothrombotic ultra-large VWF multimers released from endothelial cells in to hemostatically active high molecular weight multimers. (B) In inflammatory disorders, proinflammatory cytokines (e.g., interleukin [IL]-8 and tumor necrosis factor [TNF]- $\alpha$ ) stimulate excess release of VWF stored in Weibel–Palade bodies of endothelial cells. VWF interacts with neutrophil extracellular traps (NETs) released from neutrophils to provide a scaffold for platelet adhesion and thrombus formation. (C) In inflammation, cleavage of VWF by ADAMTS-13 is prevented by multiple mechanisms that either inhibit or reduce the proteolytic activity of ADAMTS-13.

during COVID-19 infection is greatly needed to better guide thromboprophylaxis and treatment. The extent to which VWF-ADAMTS-13 interactions contribute to the pathophysiology of COVID-19 should be an important investigative focus.

#### Funding

This work was partly supported by funds from the Hemostasis Branch/Division of Plasma Protein Therapeutics/Office of Tissues and Advanced Therapies/Center for Biologics Evaluation and Research of the U.S. Food and Drug Administration. This research was also supported by the Intramural Research Program of the National Library of Medicine at the NIH.

#### Conflict of interest

None declared.

#### References

- 1 Lu R, Zhao X, Li J, et al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet* 2020;395(10224):565–574
- 2 Epidemiology Working Group for NCIP Epidemic Response, Chinese Center for Disease Control and Prevention. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China [in Chinese]. *Zhonghua Liu Xing Bing Xue Za Zhi* 2020;41(02):145–151
- 3 Singhal T. A review of coronavirus disease-2019 (COVID-19). *Indian J Pediatr* 2020;87(04):281–286

- 4 Rothe C, Schunk M, Sothmann P, et al. Transmission of 2019-nCoV infection from an asymptomatic contact in Germany. *N Engl J Med* 2020;382(10):970–971
- 5 Petrosillo N, Viceconte G, Ergonul O, Ippolito G, Petersen E. COVID-19, SARS and MERS: are they closely related? *Clin Microbiol Infect* 2020;26(06):729–734
- 6 Gando S, Fujishima S, Saitoh D. Japanese Association for Acute Medicine (JAAM) Focused Outcomes Research in Emergency Care in Acute Respiratory Distress Syndrome, Sepsis and Trauma (FORECAST) Study Group, et al; The significance of disseminated intravascular coagulation on multiple organ dysfunction during the early stage of acute respiratory distress syndrome. *Thromb Res* 2020;191:15–21
- 7 Helms J, Tacquard C, Severac F. CRICS TRIGGERSEP Group (Clinical Research in Intensive Care and Sepsis Trial Group for Global Evaluation and Research in Sepsis), et al; High risk of thrombosis in patients with severe SARS-CoV-2 infection: a multicenter prospective cohort study. *Intensive Care Med* 2020;46(06):1089–1098
- 8 Boral BM, Williams DJ, Boral LI. Disseminated intravascular coagulation. *Am J Clin Pathol* 2016;146(06):670–680
- 9 Willyard C. Coronavirus blood-clot mystery intensifies. *Nature* 2020;581(7808):250
- 10 Thachil J, Tang N, Gando S, et al. ISTH interim guidance on recognition and management of coagulopathy in COVID-19. *J Thromb Haemost* 2020;18(05):1023–1026
- 11 Tang N, Bai H, Chen X, Gong J, Li D, Sun Z. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. *J Thromb Haemost* 2020;18(05):1094–1099
- 12 Tang N, Li D, Wang X, Sun Z. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost* 2020;18(04):844–847
- 13 Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395(10223):497–506
- 14 Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020;323(11):1061–1069
- 15 Han H, Yang L, Liu R, et al. Prominent changes in blood coagulation of patients with SARS-CoV-2 infection. *Clin Chem Lab Med* 2020;58(07):1116–1120
- 16 Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395(10229):1054–1062
- 17 Richardson S, Hirsch JS, Narasimhan M, et al. The Northwell COVID-19 Research Consortium, et al; Presenting characteristics, comorbidities, and outcomes among 5700 patients hospitalized with COVID-19 in the New York City area. *JAMA* 2020;323(20):2052–2059
- 18 Ranucci M, Ballotta A, Di Dedda U, et al. The procoagulant pattern of patients with COVID-19 acute respiratory distress syndrome. *J Thromb Haemost* 2020;18(07):1747–1751
- 19 Lippi G, Plebani M, Henry BM. Thrombocytopenia is associated with severe coronavirus disease 2019 (COVID-19) infections: a meta-analysis. *Clin Chim Acta* 2020;506:145–148
- 20 Litjens JF, Leclerc M, Chochois C, et al. High incidence of venous thromboembolic events in anticoagulated severe COVID-19 patients. *J Thromb Haemost* 2020;18(07):1743–1746
- 21 Ackermann M, Verleden SE, Kuehnel M, et al. Pulmonary vascular endothelialitis, thrombosis, and angiogenesis in Covid-19. *N Engl J Med* 2020;383(02):120–128
- 22 Klok FA, Kruip MJHA, van der Meer NJM, et al. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. *Thromb Res* 2020;191:145–147
- 23 Lodigiani C, Iapichino G, Carenzo L. Humanitas COVID-19 Task Force, et al; Venous and arterial thromboembolic complications in COVID-19 patients admitted to an academic hospital in Milan, Italy. *Thromb Res* 2020;191:9–14
- 24 Middeldorp S, Coppens M, van Haaps TF, et al. Incidence of venous thromboembolism in hospitalized patients with COVID-19. *J Thromb Haemost* 2020
- 25 Nahum J, Morichau-Beauchant T, Daviaud F, et al. Venous thrombosis among critically ill patients with coronavirus disease 2019 (COVID-19). *JAMA Netw Open* 2020;3(05):e2010478
- 26 Cui S, Chen S, Li X, Liu S, Wang F. Prevalence of venous thromboembolism in patients with severe novel coronavirus pneumonia. *J Thromb Haemost* 2020;18(06):1421–1424
- 27 Zhang C, Zhang Z, Mi J, et al. The cumulative venous thromboembolism incidence and risk factors in intensive care patients receiving the guideline-recommended thromboprophylaxis. *Medicine (Baltimore)* 2019;98(23):e15833
- 28 Wang J, Hajizadeh N, Moore EE, et al. Tissue plasminogen activator (tPA) treatment for COVID-19 associated acute respiratory distress syndrome (ARDS): a case series. *J Thromb Haemost* 2020;18(07):1752–1755
- 29 Zhang L, Yan X, Fan Q, et al. D-dimer levels on admission to predict in-hospital mortality in patients with Covid-19. *J Thromb Haemost* 2020;18(06):1324–1329
- 30 Urban K, Kirley K, Stevermer JJ. PURLs: it's time to use an age-based approach to D-dimer. *J Fam Pract* 2014;63(03):155–158
- 31 Schäfer M, Werner S. Cancer as an overheating wound: an old hypothesis revisited. *Nat Rev Mol Cell Biol* 2008;9(08):628–638
- 32 Merad M, Martin JC. Pathological inflammation in patients with COVID-19: a key role for monocytes and macrophages. *Nat Rev Immunol* 2020;20(06):355–362
- 33 Hemker HC, Giesen P, Al Dieri R, et al. Calibrated automated thrombin generation measurement in clotting plasma. *Pathophysiol Haemost Thromb* 2003;33(01):4–15
- 34 Simpson ML, Goldenberg NA, Jacobson LJ, Bombardier CG, Hathaway WE, Manco-Johnson MJ. Simultaneous thrombin and plasmin generation capacities in normal and abnormal states of coagulation and fibrinolysis in children and adults. *Thromb Res* 2011;127(04):317–323
- 35 Wu YP, Wei R, Liu ZH, et al. Analysis of thrombotic factors in severe acute respiratory syndrome (SARS) patients. *Thromb Haemost* 2006;96(01):100–101
- 36 Goyal P, Choi JJ, Pinheiro LC, et al. Clinical characteristics of Covid-19 in New York City. *N Engl J Med* 2020;382(24):2372–2374
- 37 Ruan Q, Yang K, Wang W, Jiang L, Song J. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive Care Med* 2020;46(05):846–848
- 38 Petrie JR, Guzik TJ, Touyz RM. Diabetes, hypertension, and cardiovascular disease: clinical insights and vascular mechanisms. *Can J Cardiol* 2018;34(05):575–584
- 39 Milan-Mattos JC, Anibal FF, Perseguini NM, et al. Effects of natural aging and gender on pro-inflammatory markers. *Braz J Med Biol Res* 2019;52(09):e8392
- 40 Engelmann B, Massberg S. Thrombosis as an intravascular effector of innate immunity. *Nat Rev Immunol* 2013;13(01):34–45
- 41 Jamilloux Y, Henry T, Belot A, et al. Should we stimulate or suppress immune responses in COVID-19? Cytokine and anti-cytokine interventions. *Autoimmun Rev* 2020;19(07):102567
- 42 Giamarellos-Bourboulis EJ, Netea MG, Rovina N, et al. Complex immune dysregulation in COVID-19 patients with severe respiratory failure. *Cell Host Microbe* 2020;27(06):992–1000.e3
- 43 Zhang C, Wu Z, Li JW, Zhao H, Wang GQ. Cytokine release syndrome in severe COVID-19: interleukin-6 receptor antagonist tocilizumab may be the key to reduce mortality. *Int J Antimicrob Agents* 2020;55(05):105954
- 44 Guan WJ, Ni ZY, Hu Y. China Medical Treatment Expert Group for Covid-19, et al; Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020;382(18):1708–1720

- 45 Zhu J, Ji P, Pang J, et al. Clinical characteristics of 3,062 COVID-19 patients: a meta-analysis. *J Med Virol* 2020. Doi: 10.1002/jmv.25884
- 46 Connors JM, Levy JH. COVID-19 and its implications for thrombosis and anticoagulation. *Blood* 2020;135(23):2033–2040
- 47 Chen G, Wu D, Guo W, et al. Clinical and immunological features of severe and moderate coronavirus disease 2019. *J Clin Invest* 2020;130(05):2620–2629
- 48 Tabatabai A, Rabin J, Menaker J, et al. Factor VIII and functional protein C activity in critically ill patients with coronavirus disease 2019: a case series. *A A Pract* 2020;14(07):e01236
- 49 Han H, Ma Q, Li C, et al. Profiling serum cytokines in COVID-19 patients reveals IL-6 and IL-10 are disease severity predictors. *Emerg Microbes Infect* 2020;9(01):1123–1130
- 50 Du H, Dong X, Zhang JJ, et al. Clinical characteristics of 182 pediatric COVID-19 patients with different severities and allergic status. *Allergy* 2020
- 51 Liu J, Li S, Liu J, et al. Longitudinal characteristics of lymphocyte responses and cytokine profiles in the peripheral blood of SARS-CoV-2 infected patients. *EBioMedicine* 2020;55:102763
- 52 Wang Z, Yang B, Li Q, Wen L, Zhang R. Clinical features of 69 cases with coronavirus disease 2019 in Wuhan, China. *Clin Infect Dis* 2020:ciaa272
- 53 Tan M, Liu Y, Zhou R, et al. Immunopathological characteristics of coronavirus disease 2019 cases in Guangzhou, China. *Immunology* 2020;160(03):261–268
- 54 Chen X, Zhao B, Qu Y, et al. Detectable serum SARS-CoV-2 viral load (RNAemia) is closely correlated with drastically elevated interleukin 6 (IL-6) level in critically ill COVID-19 patients. *Clin Infect Dis* 2020:ciaa449
- 55 Tocilizumab in COVID-19 Pneumonia (TOCIDVID-19). Available at: <https://ClinicalTrials.gov/show/NCT04317092>. Accessed July 29, 2020
- 56 Tocilizumab for Prevention of Respiratory Failure in Patients With Severe COVID-19 Infection. Available at: <https://ClinicalTrials.gov/show/NCT04377659>. Accessed July 29, 2020
- 57 Efficacy of Early Administration of Tocilizumab in COVID-19 Patients. Available at: <https://ClinicalTrials.gov/show/NCT04346355>. Accessed July 29, 2020
- 58 Tocilizumab in the Treatment of Coronavirus Induced Disease (COVID-19). Available at: <https://ClinicalTrials.gov/show/NCT04335071>. Accessed July 29, 2020
- 59 Premkumar M, Saxena P, Rangegowda D, et al. Coagulation failure is associated with bleeding events and clinical outcome during systemic inflammatory response and sepsis in acute-on-chronic liver failure: an observational cohort study. *Liver Int* 2019;39(04):694–704
- 60 Mojiri A, Nakhaii-Nejad M, Phan WL, et al. Hypoxia results in upregulation and de novo activation of von Willebrand factor expression in lung endothelial cells. *Arterioscler Thromb Vasc Biol* 2013;33(06):1329–1338
- 61 Matsuura Y, Yamashita A, Iwakiri T, et al. Vascular wall hypoxia promotes arterial thrombus formation via augmentation of vascular thrombogenicity. *Thromb Haemost* 2015;114(01):158–172
- 62 Ogawa S, Clauss M, Kuwabara K, et al. Hypoxia induces endothelial cell synthesis of membrane-associated proteins. *Proc Natl Acad Sci U S A* 1991;88(21):9897–9901
- 63 Fearn C, Loskutoff DJ. Induction of plasminogen activator inhibitor 1 gene expression in murine liver by lipopolysaccharide. Cellular localization and role of endogenous tumor necrosis factor-alpha. *Am J Pathol* 1997;150(02):579–590
- 64 Katneni UK, Ibla JC, Hunt R, Schiller T, Kimchi-Sarfaty C. von Willebrand factor/ADAMTS-13 interactions at birth: implications for thrombosis in the neonatal period. *J Thromb Haemost* 2019;17(03):429–440
- 65 Bryckaert M, Rosa JP, Denis CV, Lenting PJ. Of von Willebrand factor and platelets. *Cell Mol Life Sci* 2015;72(02):307–326
- 66 Zheng XL. ADAMTS13 and von Willebrand factor in thrombotic thrombocytopenic purpura. *Annu Rev Med* 2015;66:211–225
- 67 Stocksclaeder M, Schneppenheim R, Budde U. Update on von Willebrand factor multimers: focus on high-molecular-weight multimers and their role in hemostasis. *Blood Coagul Fibrinolysis* 2014;25(03):206–216
- 68 Mannucci PM, Capoferri C, Canciani MT. Plasma levels of von Willebrand factor regulate ADAMTS-13, its major cleaving protease. *Br J Haematol* 2004;126(02):213–218
- 69 Reiter RA, Knöbl P, Varadi K, Turecek PL. Changes in von Willebrand factor-cleaving protease (ADAMTS13) activity after infusion of desmopressin. *Blood* 2003;101(03):946–948
- 70 Reiter RA, Varadi K, Turecek PL, Jilma B, Knöbl P. Changes in ADAMTS13 (von-Willebrand-factor-cleaving protease) activity after induced release of von Willebrand factor during acute systemic inflammation. *Thromb Haemost* 2005;93(03):554–558
- 71 Sonneveld MA, de Maat MP, Leebeek FW. Von Willebrand factor and ADAMTS13 in arterial thrombosis: a systematic review and meta-analysis. *Blood Rev* 2014;28(04):167–178
- 72 Bongers TN, de Maat MP, van Goor ML, et al. High von Willebrand factor levels increase the risk of first ischemic stroke: influence of ADAMTS13, inflammation, and genetic variability. *Stroke* 2006;37(11):2672–2677
- 73 Sonneveld MA, de Maat MP, Portegies ML, et al. Low ADAMTS13 activity is associated with an increased risk of ischemic stroke. *Blood* 2015;126(25):2739–2746
- 74 Lambers M, Goldenberg NA, Kenet G, et al. Role of reduced ADAMTS13 in arterial ischemic stroke: a pediatric cohort study. *Ann Neurol* 2013;73(01):58–64
- 75 Hunt R, Hoffman CM, Emani S, et al. Elevated preoperative von Willebrand factor is associated with perioperative thrombosis in infants and neonates with congenital heart disease. *J Thromb Haemost* 2017;15(12):2306–2316
- 76 Andersson HM, Siegerink B, Luken BM, et al. High VWF, low ADAMTS13, and oral contraceptives increase the risk of ischemic stroke and myocardial infarction in young women. *Blood* 2012;119(06):1555–1560
- 77 Gagnano F, Sperlongano S, Golia E, et al. The role of von Willebrand factor in vascular inflammation: from pathogenesis to targeted therapy. *Mediators Inflamm* 2017;2017:5620314
- 78 Hyseni A, Kemperman H, de Lange DW, Kesecioglu J, de Groot PG, Roest M. Active von Willebrand factor predicts 28-day mortality in patients with systemic inflammatory response syndrome. *Blood* 2014;123(14):2153–2156
- 79 Bernardo A, Ball C, Nolasco L, Moake JF, Dong JF. Effects of inflammatory cytokines on the release and cleavage of the endothelial cell-derived ultralarge von Willebrand factor multimers under flow. *Blood* 2004;104(01):100–106
- 80 Chauhan AK, Kisucka J, Brill A, Walsh MT, Scheiflinger F, Wagner DD. ADAMTS13: a new link between thrombosis and inflammation. *J Exp Med* 2008;205(09):2065–2074
- 81 Bockmeyer CL, Claus RA, Budde U, et al. Inflammation-associated ADAMTS13 deficiency promotes formation of ultra-large von Willebrand factor. *Haematologica* 2008;93(01):137–140
- 82 Reuken PA, Kussmann A, Kiehntopf M, et al. Imbalance of von Willebrand factor and its cleaving protease ADAMTS13 during systemic inflammation superimposed on advanced cirrhosis. *Liver Int* 2015;35(01):37–45
- 83 Kremer Hovinga JA, Zeerleder S, Kessler P, et al. ADAMTS-13, von Willebrand factor and related parameters in severe sepsis and septic shock. *J Thromb Haemost* 2007;7(11):2284–2290
- 84 Bongers TN, Emonts M, de Maat MP, et al. Reduced ADAMTS13 in children with severe meningococcal sepsis is associated with severity and outcome. *Thromb Haemost* 2010;103(06):1181–1187
- 85 Karim F, Adil SN, Afaq B, Ul Haq A. Deficiency of ADAMTS-13 in pediatric patients with severe sepsis and impact on in-hospital mortality. *BMC Pediatr* 2013;13(01):44

- 86 Fukushima H, Nishio K, Asai H, et al. Ratio of von Willebrand factor propeptide to ADAMTS13 is associated with severity of sepsis. *Shock* 2013;39(05):409–414
- 87 Hyun J, Kim HK, Kim JE, et al. Correlation between plasma activity of ADAMTS-13 and coagulopathy, and prognosis in disseminated intravascular coagulation. *Thromb Res* 2009;124(01):75–79
- 88 Habe K, Wada H, Ito-Habe N, et al. Plasma ADAMTS13, von Willebrand factor (VWF) and VWF propeptide profiles in patients with DIC and related diseases. *Thromb Res* 2012;129(05):598–602
- 89 Vardavas CI, Nikitara K. COVID-19 and smoking: a systematic review of the evidence. *Tob Induc Dis* 2020;18:20
- 90 Ma Q, Jacobi PM, Emmer BT, et al. Genetic variants in *ADAMTS13* as well as smoking are major determinants of plasma ADAMTS13 levels. *Blood Adv* 2017;1(15):1037–1046
- 91 Leung JM, Yang CX, Tam A, et al. ACE-2 expression in the small airway epithelia of smokers and COPD patients: implications for COVID-19. *Eur Respir J* 2020;55(05):2000688
- 92 Barigye O. Smoking and inflammation. *PLoS Med* 2005;2(06):e198
- 93 Schwameis M, Schörgenhofer C, Assinger A, Steiner MM, Jilma B. VWF excess and ADAMTS13 deficiency: a unifying pathomechanism linking inflammation to thrombosis in DIC, malaria, and TTP. *Thromb Haemost* 2015;113(04):708–718
- 94 Wang A, Liu F, Dong N, et al. Thrombospondin-1 and ADAMTS13 competitively bind to VWF A2 and A3 domains in vitro. *Thromb Res* 2010;126(04):e260–e265
- 95 Bonnefoy A, Daenens K, Feys HB, et al. Thrombospondin-1 controls vascular platelet recruitment and thrombus adherence in mice by protecting (sub)endothelial VWF from cleavage by ADAMTS13. *Blood* 2006;107(03):955–964
- 96 Pillai VG, Bao J, Zander CB, et al. Human neutrophil peptides inhibit cleavage of von Willebrand factor by ADAMTS13: a potential link of inflammation to TTP. *Blood* 2016;128(01):110–119
- 97 Chen J, Fu X, Wang Y, et al. Oxidative modification of von Willebrand factor by neutrophil oxidants inhibits its cleavage by ADAMTS13. *Blood* 2010;115(03):706–712
- 98 Crawley JT, Lam JK, Rance JB, Mollica LR, O'Donnell JS, Lane DA. Proteolytic inactivation of ADAMTS13 by thrombin and plasmin. *Blood* 2005;105(03):1085–1093
- 99 Ono T, Mimuro J, Madoiwa S, et al. Severe secondary deficiency of von Willebrand factor-cleaving protease (ADAMTS13) in patients with sepsis-induced disseminated intravascular coagulation: its correlation with development of renal failure. *Blood* 2006;107(02):528–534
- 100 Liu C, Zhao L, Zhao J, Xu Q, Song Y, Wang H. Reduced ADAMTS-13 level negatively correlates with inflammation factors in plasma of acute myeloid leukemia patients. *Leuk Res* 2017;53:57–64
- 101 Takaya H, Kawaratani H, Kubo T, et al. Platelet hyperaggregability is associated with decreased ADAMTS13 activity and enhanced endotoxemia in patients with acute cholangitis. *Hepatol Res* 2018;48(03):E52–E60
- 102 Takaya H, Yoshiji H, Kawaratani H, et al. Decreased activity of plasma ADAMTS13 are related to enhanced cytokinemia and endotoxemia in patients with acute liver failure. *Biomed Rep* 2017;7(03):277–285
- 103 Chen J, Chung DW. Inflammation, von Willebrand factor, and ADAMTS13. *Blood* 2018;132(02):141–147
- 104 Bazzan M, Montaruli B, Sciascia S, Cosseddu D, Norbiato C, Roccatello D. Low ADAMTS 13 plasma levels are predictors of mortality in COVID-19 patients. *Intern Emerg Med* 2020
- 105 Huisman A, Beun R, Sikma M, Westerink J, Kusadasi N. Involvement of ADAMTS13 and von Willebrand factor in thromboembolic events in patients infected with SARS-CoV-2. *Int J Lab Hematol* 2020
- 106 Adam EH, Zacharowski K, Miesbach W. A comprehensive assessment of the coagulation profile in critically ill COVID-19 patients. *Thromb Res* 2020;194:42–44
- 107 Latimer G, Corriveau C, DeBiasi RL, et al. Cardiac dysfunction and thrombocytopenia-associated multiple organ failure inflammation phenotype in a severe paediatric case of COVID-19. *Lancet Child Adolesc Health* 2020;4(07):552–554
- 108 Escher R, Breakey N, Lämmle B. ADAMTS13 activity, von Willebrand factor, factor VIII and D-dimers in COVID-19 inpatients. *Thromb Res* 2020;192:174–175
- 109 Escher R, Breakey N, Lämmle B. Severe COVID-19 infection associated with endothelial activation. *Thromb Res* 2020;190:62
- 110 Lee SJ, Kim JE, Han KS, Kim HK. Thrombotic risk of reduced ADAMTS13 activity in patients with antiphospholipid antibodies. *Blood Coagul Fibrinolysis* 2016;27(08):907–912
- 111 Galeano-Valle F, Oblitas CM, Ferreiro-Mazón MM, et al. Antiphospholipid antibodies are not elevated in patients with severe COVID-19 pneumonia and venous thromboembolism. *Thromb Res* 2020;192:113–115
- 112 Bowles L, Platton S, Yartey N, et al. Lupus anticoagulant and abnormal coagulation tests in patients with Covid-19. *N Engl J Med* 2020;383(03):288–290
- 113 Austin SK, Starke RD, Lawrie AS, Cohen H, Machin SJ, Mackie IJ. The VWF/ADAMTS13 axis in the antiphospholipid syndrome: ADAMTS13 antibodies and ADAMTS13 dysfunction. *Br J Haematol* 2008;141(04):536–544
- 114 Jian C, Xiao J, Gong L, et al. Gain-of-function ADAMTS13 variants that are resistant to autoantibodies against ADAMTS13 in patients with acquired thrombotic thrombocytopenic purpura. *Blood* 2012;119(16):3836–3843
- 115 Lorenzo-Villalba N, Zulfiqar AA, Auburtin M, et al. Thrombocytopenia in the course of COVID-19 infection. *Eur J Case Rep Intern Med* 2020;7(06):001702
- 116 Yin S, Huang M, Li D, Tang N. Difference of coagulation features between severe pneumonia induced by SARS-CoV2 and non-SARS-CoV2. *J Thromb Thrombolysis* 2020. Doi: 10.1007/s11239-020-02105-8
- 117 Demelo-Rodríguez P, Cervilla-Muñoz E, Ordieres-Ortega L, et al. Incidence of asymptomatic deep vein thrombosis in patients with COVID-19 pneumonia and elevated D-dimer levels. *Thromb Res* 2020;192:23–26
- 118 Grandmaison G, Andrey A, Périard D, et al. Systematic screening for venous thromboembolic events in COVID-19 pneumonia. *TH Open* 2020;4(02):e113–e115
- 119 Fraissé M, Logre E, Pajot O, Mentec H, Plantefève G, Contou D. Thrombotic and hemorrhagic events in critically ill COVID-19 patients: a French monocenter retrospective study. *Crit Care* 2020;24(01):275
- 120 Bastiani G, Valle MT. Determination of factor XII in blood products and correction of its deficiency after plasma transfusion in a case [in Italian]. *Haematologica* 1979;64(05):635–640
- 121 Desborough MJR, Doyle AJ, Griffiths A, Retter A, Breen KA, Hunt BJ. Image-proven thromboembolism in patients with severe COVID-19 in a tertiary critical care unit in the United Kingdom. *Thromb Res* 2020;193:1–4
- 122 Akel T, Qaqa F, Abuarqoub A, Shamoof F. Pulmonary embolism: a complication of COVID 19 infection. *Thromb Res* 2020;193:79–82
- 123 Kashi M, Jacquin A, Dakhil B, et al. Severe arterial thrombosis associated with Covid-19 infection. *Thromb Res* 2020;192:75–77
- 124 Lax SF, Skok K, Zechner P, et al. Pulmonary arterial thrombosis in COVID-19 with fatal outcome: results from a prospective, single-center, clinicopathologic case series. *Ann Intern Med* 2020;173(05):350–361
- 125 Thomas W, Varley J, Johnston A, et al. Thrombotic complications of patients admitted to intensive care with COVID-19 at a teaching hospital in the United Kingdom. *Thromb Res* 2020;191:76–77
- 126 Gomez-Arbelaiz D, Ibarra-Sanchez G, Garcia-Gutierrez A, Comanges-Yeboles A, Ansuategui-Vicente M, Gonzalez-Fajardo

- JA. COVID-19-related aortic thrombosis: a report of four cases. *Ann Vasc Surg* 2020;S0890-5096(20)30438-6
- 127 Ayerbe L, Risco C, Ayis S. The association between treatment with heparin and survival in patients with Covid-19. *J Thromb Thrombolysis* 2020;50(02):298-301
- 128 Wang T, Chen R, Liu C, et al. Attention should be paid to venous thromboembolism prophylaxis in the management of COVID-19. *Lancet Haematol* 2020;7(05):e362-e363
- 129 Artifoni M, Danic G, Gautier G, et al. Systematic assessment of venous thromboembolism in COVID-19 patients receiving thromboprophylaxis: incidence and role of D-dimer as predictive factors. *J Thromb Thrombolysis* 2020;50(01):211-216
- 130 Russo V, Di Maio M, Attena E, et al. Clinical impact of pre-admission antithrombotic therapy in hospitalized patients with COVID-19: a multicenter observational study. *Pharmacol Res* 2020;159:104965
- 131 Maier CL, Truong AD, Auld SC, Polly DM, Tanksley CL, Duncan A. COVID-19-associated hyperviscosity: a link between inflammation and thrombophilia? *Lancet* 2020;395(10239):1758-1759
- 132 Magro C, Mulvey JJ, Berlin D, et al. Complement associated microvascular injury and thrombosis in the pathogenesis of severe COVID-19 infection: a report of five cases. *Transl Res* 2020;220:1-13