

Preprocedural Use of Povidone-Iodine Mouthwash during Dental Procedures in the COVID-19 Pandemic

Eisha Imran¹ Zohaib Khurshid^{2,✉} Adam Ahmed M. Al Qadhi³
Ahmed Adel A. Al-Quraini³ Khaloud Tariq⁴

¹Department of Dental Materials, Dental College, HITEC-Institute of Medical Sciences, Rawalpindi, Pakistan

²Department of Prosthodontics and Dental Implantology, College of Dentistry, King Faisal University, Al Ahsa, Saudi Arabia

³Department of Dental Hospital, College of Dentistry, King Faisal University, Al Ahsa, Saudi Arabia

⁴Department of Community Dentistry, University College of Dentistry, The University of Lahore, Lahore, Pakistan

Eur J Dent:2020;14(suppl S1):S182–S184

A bizarre situation has been created in the lives of individuals across the globe due to coronavirus disease (COVID-19)'s uncertain time and the severity of its consequences. Since December 2019, health care workers are endeavoring on effective treatment plans and implementing strategies to control the spread of the virus. Clinical symptoms of the disease vary from mild respiratory depression to severe organ failure.¹ A recently published global survey assessed the level of fear and practice modifications by dental practitioners worldwide, while it further emphasized on following guidelines by health policy departments.² Respiratory droplets generated during human-to-human interaction are acknowledged as the source of its transmission, hence creating an alarming situation for the health care workers as they are frontline workers, exposed to the highest viral load.³ Excessive viral load found in the nasal cavity, oropharynx, nasopharynx, and saliva is the reason behind the highest transmission risk in a dental care.^{4,5} Saliva can be an important diagnostic tool for rapid mass detection of patients based on the fact that it is an essential biomarker, used for diagnosing several viral and bacterial infections.⁶

Adversity in our dental setting during this pandemic is not an exception, thus creating an immense need to adopt precautionary measures during dental practices.³ A virus concentration of 10^7 /mL is found in the saliva during the early infectious period, thus augmenting the risk of cross-infection and instrument contamination.⁷ In contrast, asymptomatic person makes the spread even more challenging since they carry and shed millions of virus particles.

Harrel and Molinari proposed the use of three protective coatings for health care workers which include personal protective equipment, mouthwashes, and high-speed suction

Address for correspondence Zohaib Khurshid, MRes, FHEA, MDTFEd, FPFA, BDS, Department of Prosthodontics and Dental Implantology, College of Dentistry, King Faisal University, Al Ahsa 31982, Saudi Arabia
(e-mail: drzohaibkhurshid@gmail.com).

instruments.⁸ Preprocedural decontamination of the oral cavity is essential. The Centers for Disease Control (CDC) and Prevention guidelines acknowledged the utilization of chlorhexidine oral rinses to reduce aerosol production (<https://www.cdc.gov/mmwr/preview/mmwrhtml/rr5217a1.htm>).

However, preliminary *in vitro* investigations spotlighted the effectiveness of chlorhexidine only against bacterial particles while it showed negligible antiviral activity.⁹ Regrettably for us, little is covered in the literature regarding the antiviral properties of different available oral rinses, and the utilization of these mouthwashes is not considered in our dental practice.¹⁰

Antecedent *in vitro* investigations on the severe acute respiratory syndrome coronavirus (SARS-CoV) (2002–2003) and the Middle East respiratory syndrome coronavirus (MERS-CoV) (2012–2013), the causative agents accountable for the previous pandemics in history, revealed significantly higher antiviral properties of povidone-iodine (PVP-I) mouthwashes in contrast to other oral rinses which include chlorhexidine and benzalkonium.¹¹ All these coronaviruses strains are coated with a lipid membrane, thus having the same structure as SARS-CoV-2. Hence, PVP-I can serve as a potential candidate to destroy the virus membrane, thus limiting its transmission. Bidra et al, in a recently conducted *in vitro* investigation, concluded that a 0.5% lowest concentration of PVP-I was required with a contact time of 15 seconds to eradicate SARS-CoV-2 virus successfully.¹⁰ Eggers et al evaluated the efficacy of PVP-I formulation at 1% concentration against MERS-CoV and modified Vaccinia virus Ankara and concluded its excellent virucidal activity within 15 to 60 seconds.¹² In another investigation, they evaluated a reduction in the viral titers after 15 to 30 seconds of exposure to 0.7 and 0.23% concentration of PVP-I.¹³

DOI <https://doi.org/10.1055/s-0040-1717001>
ISSN 1305-7456.

© 2020. European Journal of Dentistry.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India




Fig. 1 Protocols for running dental practice in the COVID-19 pandemic.^{16,17}

Earlier in June 2020, Martínez Lamas et al conducted an *in vivo* investigation which aimed to determine the influence of PVP-I on COVID-19 positive patients.¹⁴ They evaluated and compared the viral loads through polymerase chain reaction testing before and after PVP-I mouth rinse. The results signified a decline in the viral load of the saliva of these COVID-19 patients after 1-minute use of PVP-I and the effects persisted for 3 hours. However, the study had limitations such as small sample size; second, there was no elimination of confounding factors which include determination of the immune response of the patient or the concentration of viral load. Hence, these factors need to be studied to evaluate a specific concentration which could be related to its clinical efficacy. Nonetheless, this study supports the fact that PVP-I can significantly aid in reducing the viral load of COVID-19 patients.¹⁴ Sarfaraz et al, lately in a cross-sectional study, concluded that majority of dental practitioners utilized chlorhexidine oral rinses in contrast to other mouthwashes during this pandemic, thus revealing a lack of knowledge regarding the antiviral properties of PVP-I oral rinses.¹⁵

Representation of reported protocols for the use of PVP-I is illustrated in ►Fig. 1 so that their use can practically be implemented in a dental setting.^{16,17} The American Dental Association (ADA) advised preprocedural utilization of 0.2% PVP-I to reduce transmission. Based on the literature search by Frank et al, 2.5% of PVP-I formulation can be used to decontaminate the oral cavity, keeping in mind the toxic adverse effects.¹⁸ However, care should be taken for unconscious patients as cases of aspiration pneumonia have been reported; thus, a small amount is recommended. Second, it should also be spotlighted that PVP-I is not recommended for pregnant women or patients undergoing radioactive therapy or suffering from thyroid problems.¹⁸

Based on these studies, we propose and urge the dental staff for preprocedural use of 0.2% PVP-I mouthwashes for 30 seconds. PVP-I is available with trade name “Betadine,” a small quantity of the mouthwash needs to be introduced

and distributed into the mouth for 30 seconds, and gargles for further 30 seconds are recommended in the throat and spit after 1 minute.

PVP-I oral antiseptic rinses may offer promising results as a decontaminating agent in present COVID-19 era because of their supercilious efficacy against SARS-CoV-2 virus as indicated in the literature. Their preprocedural use is highly recommended. However, its dose-dependent toxic effects and contraindications should be evaluated before the prescription. Along with this opinion, we hope to get a translational step toward clinical practice for the use of PVP-I oral rinses.

Conflict of Interest

None declared.

References

- Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020;323(11):1061–1069
- Ahmed MA, Jouhar R, Ahmed N, et al. Fear and practice modifications among dentists to combat novel coronavirus disease (COVID-19) outbreak. *Int J Environ Res Public Health* 2020;17(8):2821
- Peng X, Xu X, Li Y, Cheng L, Zhou X, Ren B. Transmission routes of 2019-nCoV and controls in dental practice. *Int J Oral Sci* 2020;12(1):9
- Zou L, Ruan F, Huang M, et al. SARS-CoV-2 viral load in upper respiratory specimens of infected patients. *N Engl J Med* 2020;382(12):1177–1179
- Khurshid Z, Asiri FYI, Al Wadaani H. Human saliva: non-invasive fluid for detecting novel coronavirus (2019-nCoV) *Int J Environ Res Public Health* 2020;17(7):2225
- Hamid H, Khurshid Z, Adanir N, Zafar MS, Zohaib S. COVID-19 pandemic and role of human saliva as a testing biofluid in point-of-care technology 2020. *Eur J Dent* 2020;14(suppl S1):S123–S129 doi:10.1055/s-0040-1713020
- To KK, Tsang OTY, Yip CC, et al. Consistent detection of 2019 novel coronavirus in saliva. *Clin Infect Dis* 2020; 71(15):841–843

- 8 Harrel SK, Molinari J. Aerosols and splatter in dentistry: a brief review of the literature and infection control implications. *J Am Dent Assoc* 2004;135(4):429–437
- 9 Marui VC, Souto MLS, Rovai ES, Romito GA, Chambrone L, Pannuti CM. Efficacy of preprocedural mouthrinses in the reduction of microorganisms in aerosol: a systematic review. *J Am Dent Assoc* 2019;150(12):1015–1026.e1
- 10 Bidra AS, Pelletier JS, Westover JB, Frank S, Brown SM, Tessema B. Rapid in-vitro inactivation of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) using povidone-iodine oral antiseptic rinse. *J Prosthodont* 2020;29(6):529–533
- 11 Kariwa H, Fujii N, Takashima I. Inactivation of SARS coronavirus by means of povidone-iodine, physical conditions and chemical reagents. *Dermatology* 2006;212(1, suppl 1):119–123
- 12 Eggers M, Eickmann M, Zorn J. Rapid and effective virucidal activity of povidone-iodine products against Middle East respiratory syndrome coronavirus (MERS-CoV) and modified vaccinia virus Ankara (MVA) *Infect Dis Ther* 2015;4(4):491–501
- 13 Eggers M, Koburger-Janssen T, Eickmann M, Zorn J. In vitro bactericidal and virucidal efficacy of povidone-iodine gargle/mouthwash against respiratory and oral tract pathogens. *Infect Dis Ther* 2018;7(2):249–259
- 14 Martínez Lamas L, Diz Dios P, Pérez Rodríguez MT, et al. Is povidone iodine mouthwash effective against SARS-CoV-2? *Oral Dis* 2020;10.1111/odi.13526 doi:10.1111/odi.13526
- 15 Sarfaraz S, Shabbir J, Mudasser MA, et al. Knowledge and attitude of dental practitioners related to disinfection during the COVID-19 pandemic. *Healthcare (Basel)* 2020;8(3):232
- 16 Kirk-Bayley J, Challacombe S, Sunkaraneni V, Combes J. The use of povidone iodine nasal spray and mouthwash during the current COVID-19 pandemic may protect healthcare workers and reduce cross infection. *SSRN Electron J* 2020. Doi: 10.2139/ssrn.3563092
- 17 Mady LJ, Kubik MW, Baddour K, Snyderman CH, Rowan NR. Consideration of povidone-iodine as a public health intervention for COVID-19: utilization as “Personal Protective Equipment” for frontline providers exposed in high-risk head and neck and skull base oncology care. *Oral Oncol* 2020;105:104724
- 18 Frank S, Capriotti J, Brown SM, Tessema B. Povidone-iodine use in sinonasal and oral cavities: a review of safety in the COVID-19 era. *Ear Nose Throat J* 2020;(June):145561320932318