


C. CHEN, J. C. PETERS\*, G. C. FU\* (CALIFORNIA INSTITUTE OF TECHNOLOGY, PASADENA, USA)


Photoinduced Copper-Catalyzed Asymmetric Amidation via Ligand Cooperativity

Nature 2021, 596, 250–256, DOI: 10.1038/s41586-021-03730-w.


# Enantioconvergent Photoinduced Copper-Catalyzed Synthesis of Chiral Secondary Amides



## Selected examples:



## Proposed mechanism:



**Significance:** Under blue-LED irradiation, the enantioconvergent C–N coupling of racemic alkyl electrophiles with primary amides using copper catalysis has been described. A variety of chiral secondary amides were formed, with a Lewis-basic functional group on the electrophile component to allow for differentiation of similar alkyl substituents. Through control studies, the stereochemistry of the amide product was shown to be determined by the diamine ligand, irrespective of the stereochemistry of the bisphosphine ligand or electrophile.

**Comment:** Extensive mechanistic studies were conducted to elucidate the role of the three ligands used in the catalyst system. Together, the bisphosphine and phenoxide ligands form the photocatalyst responsible for activating the C–Br bond, whereas the diamine ligand was found to be necessary for forming the C–N bond and controlling stereochemistry. The involvement of a radical intermediate was supported by trapping and cyclization experiments.

**SYNFACTS Contributors:** Mark Lautens, Rachel Ross  
Synfacts 2021, 17(09), 0995 Published online: 18.08.2021  
DOI: 10.1055/s-0040-1720804; Reg-No.: L10121SF

© 2021, Thieme. All rights reserved.  
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Metals in Synthesis

Key words

copper catalysis

amidation

enantioconvergence

visible light

Synfact  
of the  
Month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.